

Influences of Ammonia for Synthesis of 8-Hydroxyquinoline Copper(II)

by Sriatun Sriatun

Submission date: 25-Apr-2020 11:06PM (UTC+0700)

Submission ID: 1307518353

File name: es_of_Ammonia_for_Synthesis_of_8-Hydroxyquinoline_Copper_II.pdf (663.85K)

Word count: 2164

Character count: 12264

Influences of Ammonia for Synthesis of 8-Hydroxyquinoline Copper(II)

Suhartana, Laelatri Agustina, Sriatun

Abstract

Synthesis and characterization of complex compounds of 8-hydroxyquinoline copper (II) has been done. Synthesis of 8-hydroxyquinoline copper (II) is done by mixing $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ with different concentration of ammonia and 8-hydroxyquinoline ligand in methanol. Product performed with magnetic stirrer, filtered, washed and dried in a desiccator. The results is obtained precipitated 8-hydroxyquinoline is copper (II) green-yellow. Characterization of complex compound was done by analysis of UV-Vis, AAS and FTIR. UV-Vis analysis results of 8-hydroxyquinoline copper (II) have a maximum wavelength at 394 nm, FTIR analysis results indicate the presence of N atom and O atom of C-O clusters in 8-hydroxyquinoline ligand was bonding with Cu (II) atom. If concentration of ammonia increasing, complex 8-hydroxyquinoline copper (II) was obtained decreasing.

Keywords: Synthesis, complex compound Cu- 8-hydroxyquinoline

*Chemistry Department, Mathematics and Sciences Faculty, Diponegoro University

Corresponding author e-mail address: suhartanagi_putra@yahoo.com

Introduction

Oxine or 8-hydroxyquinoline ($\text{C}_9\text{H}_7\text{ON}$) forms sparingly soluble derivatives with metallic ions, which have the composition $\text{M}(\text{C}_9\text{H}_7\text{ON})_2$ if coordination number of the metal is four (e.g., magnesium, zinc, copper, cadmium, lead and indium), $\text{M}(\text{C}_9\text{H}_7\text{ON})_3$ if the coordination number is six (e.g., aluminium, iron, bismuth, and gallium), and $\text{M}(\text{C}_9\text{H}_7\text{ON})_4$ if the coordination number is eight (e.g., thorium and zirconium). There are, however, some exceptions, for examples, $\text{TiO}(\text{C}_9\text{H}_7\text{ON})_2$, $\text{MnO}_2(\text{C}_9\text{H}_7\text{ON})_2$, $\text{WO}_2(\text{C}_9\text{H}_7\text{ON})_2$, and $\text{UO}_2(\text{C}_9\text{H}_7\text{ON})_2$ (Vogel, 1978).

Copper (Cu)

Copper, silver and gold are in group 11 of the periodic table, and they share certain attributes. They have one s-orbital electron on top of filled d-electron shell and are characterized by high ductility and electrical conductivity. The filled d-shells in these elements do not contribute much to the interatomic interaction, which are dominated by the s-electrons through metallic bonds. Unlike in metals with incomplete d-shells, metallic bonds copper are lacking a covalent character and are relatively weak [Huheey, 1981].

The softness of copper partly explains its high electrical conductivity ($59.6 \times 10^6 \text{ s/m}$) and thus also high thermal conductivity, which are the second highest among pure metals at room temperature. This is because the resistivity to electron transport in metals at room temperature mostly originates from scattering of electrons on thermal vibrations of the lattice, which relatively weak for a soft metal [Huheey, 1981].

Copper has 29 atomic number, they are 29 isotopes of copper. ^{63}Cu and ^{65}Cu are stable, with ^{63}Cu comprising approximately 69% of naturally occurring copper. The other isotopes (^{62}Cu , ^{64}Cu and ^{67}Cu) are radioactive. Copper is present in the earth crust at a concentration of about 50 part per million (ppm), where occurs as native copper or in minerals such as the copper sulphides *chalcopyrite* or *chalcocite*, the copper carbonate *azurite* and *malachite*, and the copper (I) oxide mineral *cuprite* (Cotton and Wilkinson, 1988).

8-Hydroxyquinoline

The 8-hydroxyquinoline and its derivatives are widely used as analytical reagent [Raj, et al, 2001] and anti-amoebic agents, 8-hydroxyquinoline behaves as bidentate (N & O⁻) univalent ligand to form chelates with several metals ions [Basollo, 1973]. 8-Hydroxyquinoline and its derivatives are widely used as ligand. 8-Hydroxyquinoline has 145,16 g/mol mass relatively, white powder, not soluble with aquadest, but soluble at organic solvent and acids, acetic acid for example (Underwood, 1980).

8-hydroxyquinoline behaves as bidentate (N & O⁻) and used as ligand in synthesis complexes compounds. Structure of 8-hydroxyquinoline:




Figure 1. Structure of hydroxyquinoline

8-hydroxyquinoline has melting point 74-76 °C and has boiling point 276 °C, density 1.034 g/mol. The 8-hydroxyquinoline and its derivatives would be used as therapeutics as antibacterial, antifungal, as well as for treatment of tuberculosis, diabetes and malaria (Soekardjo, 1995).

Derivatives of 8-hydroxyquinoline have been used as tropical antiseptics and internal disinfectants that apparently exhibit low toxicity for humans (Underwood, 1998).

This research Cu metals used as the central atom and 8-hydroxyquinoline used as a ligand. 8-hydroxyquinoline ligand has O at CO bonding and N as an electron donor [Huheey, 1981]. So bonding Cu metals and 8-hydroxyquinoline ligand has been obtained. "Continue Variation" method has been done to form 8-hydroxyquinoline copper (II) (Sugiarto, 2009).

Synthesis of 8-hydroxyquinoline copper (II) in this research has been done with a mixing/reaction of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ with 8-hydroxyquinoline (in methanol as solvent) in different ammonia concentrations. Mixing by magnetic stirrer. Result obtained dried by desiccator, precipitate grey-yellow was obtained, this is 8-hydroxyquinoline copper (II) complex.

Result of 8-hydroxyquinoline copper (II) has been analysed by UV-Vis, spectrophotometry, AAS and FTIR. From UV-Vis spectra, wavelength maximum at 394 nm, FTIR analysis shows interaction of Cu metal with N atom and atom O at C-O bonding from 8-hydroxyquinoline as a coordination bond. If ammonia concentration increases, the complex of 8-hydroxyquinoline copper (II) was obtained decreasing.

Methodology

Materials

$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ p.a (Merck), Hydroxyquinoline p.a (Merck), methanol p.a (Merck), ammonia solution, aquabidest and aquadest.

Tool Volume pipet, beaker glass, analytical balance, magnetic stirrer, heating mantle, desiccator, funnel, reflux, FTIR spectrophotometer Shimadzu prestige 21, spectrophotometer UV-Vis Shimadzu 1601, atomic

absorption spectrophotometer (AAS), Whatmann paper.

Procedure

8-hydroxyquinoline Copper (II) synthesis

Becker glass I An aqueous solution contents of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (0.395 g) with ammonia concentration various (0,01M, 0,02M, 0,03M, 0,04M, 0,05 M and 0,07M) was soluble at 10 ml aquadest, Becker glass II Have contents 8-hydroxyquinoline soluble at methanol 10 ml. Becker glass I was added for Becker glass II dispensing drop by drop. Result obtained was reflux and mix by magnetic stirrer until 3 hour, and precipitate was obtained. Precipitate was filtered and wash by methanol and dried at desiccator until 3 days. Precipitate has grey-yellow colour, this is 8-hydroxyquinoline copper (II) complex. Base data, if concentration of ammonia increasing, Cu (II)- 8-hydroxyquinoline complex was decreasing obtained, indicated with the colour grey-yellow for the complex is less. Analysis the complex was done with UV-Vis spectrophotometry, AAS, and FTIR.

Result and Discussion

Synthesis of 8-hydroxyquinoline Copper (II) has been done with mixing/reaction of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ with 8-hydroxyquinoline (in methanol as solvent) in different ammonia concentrations. Mixing by magnetic stirrer. Result obtained dried by desiccator, precipitate grey-yellow was obtained, this is 8-hydroxyquinoline copper (II) complex.

Result of 8-hydroxyquinoline copper (II) has been analysed by UV-Vis, spectrophotometry, AAS and FTIR. From UV-Vis spectra, wavelength maximum at 394 nm, FTIR analysis shows interaction of Cu metal with N atomic and atomic O at C-O bonding from 8-hydroxyquinoline as a coordination bond.

Ammonia Variation

An aqueous CuSO_4 solution with ammonia concentration differently has result different. An aqueous CuSO_4 solution without ammonia the product is light blue. But if an aqueous CuSO_4 solution with ammonia the product is blue, this is tetramine copper (II) compound. If ammonia concentration increases, the result of 8-hydroxyquinoline copper (II) decreases. This indicates that 8-hydroxyquinoline copper (II) grey-yellow is less. If ammonia concentration increases, the system would be a reaction of Cu^{2+} with 8-hydroxyquinoline to form Tetramine Copper(II). So the 8-hydroxyquinoline copper (II) complex decreases.

Ammonia Effect for 8-hydroxyquinoline copper (II) synthesis

From AAS data, if ammonia concentration increases, the result of 8-hydroxyquinoline copper (II) is

decreasing. This data would prove Cu^{2+} ion remainder obtained increasing, like table 1. This data would showed, if Ammonia concentration is increase, so Tetramine Copper(II) complex in solution is increase too. Copper(II) complex would be contribute to blocked Cu^{2+} ions to react with 8-hydroxiquinoline.

Table 1. Concentration Cu^{2+} at Variation Ammonia concentration

[Cu^{2+}] total	[Ammonia]	[Cu^{2+}] remainder
100 ppm	0.01 M	3.4 ppm
100 ppm	0.02 M	4.2 ppm
100 ppm	0.03 M	6.8 ppm
100 ppm	0.04 M	9.6 ppm
100 ppm	0.05 M	12.8 ppm
100 ppm	0.07 M	18.2 ppm

UV-Vis characterization

The adsorption and assignments related the ligand and the complex showed in Figure 2. Analysis spectrophotometry UV-Vis has used for establish λ_{max} (nm) sample. This wavelength used to trace λ_{max} (nm) $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ and λ_{max} (nm) 8-hydroxyquinoline copper (II) complex, after reaction was occurred.

The spectra where the electronic configuration of the metal d^{10} continuous adsorption of any ($d-d$) transition. According the spectra data as well as those obtained from elemental analysis the chemical structure of the complex may be suggested for 8-hydroxyquinoline copper (II) complex. Change of wavelength, from 817 nm ($\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$) to 394 nm (8-hydroxyquinoline copper (II) complex) is indicated coordination bonding would be obtained. (Fessenden, 1986).




Figure 2 Spectra UV-Vis (a). $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, (b). 8-hydroxiquinolin copper (II) at methanol solvent

Transition electronic Energy 10Dq from $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ and 8-hydroxyquinoline copper (II) showed table 3. Complex compounds usually has colour. Colour obtained not colour adsorption, but reflection from complement colour. Complement colour is overall wavelength from 8-hydroxiquinoline copper (II) complex. This complex have λ_{max} 394 nm, at UV area, because peak was obtained from ultraviolet UV-Vis area and adsorption wavelength from violet to blue (Fessenden, 1986).

Table 3: Wavelength and 10Dq Energy for $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ and 8-hydroxyquinoline cuprum (II) complex

No.	Compound	λ_{max} (nm)	10Dq (KJ/mol)
1.	$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$	817	146,288
2.	8-hydroxyquinoline copper (II) complex	394	303,343

IR characterization

The characteristic vibration and assignments of ligand 8-hydroxyquinoline and complex compound are showed in figure 4 and analysed the IR spectra are described in table 5.




Figure 4 (a.) IR Spectra 8-hydroxyquinoline, (b.) 8 -hydroxyquinoline copper (II).

Table 5: IR spectra

Peak	8-hydroxyquinoline (cm ⁻¹)	8 -hydroxyquinoline copper (II). (cm ⁻¹)
5 v O-H stretch	3132.4	3448.7
v C=N stretch aromatic	1504.4	1496.7
v C-N stretch aromatic	1280.7	1327.0
v C-O stretch phenol	1056.9	1111.0
v Cu-N	-	478.3
v Cu-O	-	524.6

The spectrum of phenol exhibited the strong band of 1056,9 cm⁻¹ for 8-hydroxy-quinoline and 1111 cm⁻¹ for 8 -hydroxyquinoline copper (II). While another strong adsorption bands at 1280 cm⁻¹ for 8-hydroxyquinoline and 1327 cm⁻¹ for 8 -hydroxyquinoline copper (II).

The spectra of complex.

The spectra exhibited amarked difference the adsorption band (Figure: IV.3) belonging to be

stretching vibration of v (C-O) of the carbonyl group have been found in the range between (1100 cm⁻¹-1200 cm⁻¹). According data, suggesting the possibility of the bonding complex compound is coordination bond of the ligand 8-hydroxyquinoline through the oxygen atomic in carbonyl group.

Adsorption assigned for v (C-N) was noticed at the range 1300 cm⁻¹-1350 cm⁻¹ shifted to the higher frequencies. Adsorption assigned for v (Cu-N) was

noticed at the range 478,3 cm^{-1} and V (C-O) was noticed at the range 500 cm^{-1} – 550 cm^{-1} , which indicate the coordination of nitrogen atomic of the V (C-N) group and oxygen atomic of the V (C-O) group to the central copper ion.

Conclusion

In this study, 8-hydroxyquinoline (8HQ) copper (II) synthesis, were investigated:

1. $\text{CuSO}_4 \cdot 5 \text{H}_2\text{O}$ has UV-Vis characteristic adsorption at λ_{max} 817 nm, and 8-hydroxyquinoline (8HQ) copper (II) has UV-Vis characteristic adsorption at λ_{max} 394 nm.
2. 8-hydroxyquinoline would coordinate bond with Cu^{2+} ion, by N atomic and O atomic.

Acknowledgments

I would like to thank my student laelatri Agustina, also my friend Sri Atun for her support during my experiment. You have helped so much, there is no way I could possibly ever repay you.

Reference

Cotton, F.A. and Wilkinson, G., 1988, *Advanced Inorganic Chemistry*, Edisi ke 5, John Wiley & Sons Inc, New York

Huheey, J.E., and Keitr, R. L., 1993, *Inorganic Chemistry, Principle of Structure and Reactivity*, Edisi ke 2, Harper Collin Collage Publisher, New York

Jolly WL., 1978, *The Synthesis and Characterization of Inorganic Compounds*, Prentice hall Inc, New York.

Sukardjo, 1995, *Kimia Koordinasi*, Rineka Cipta, Jakarta

Sugiarto, 2006, *Sintesis dan Karakterisasi Kompleks Di (8- hidroksiquinolin) tembaga (II) trihidrat dan Kompleks Tri (8- hidroksiquinolin) besi (III) dihydrate*, Skripsi, FMIPA UNS, Surakarta

Underwood, A. L. & Day, R. A., 2002, *Analisis Kimia Kuantitatif*, Edisi Keenam, Penerbit Erlangga, Jakarta.

4
Vogel, A.I., 1978, *A Text Book of Quantitative Inorganic Analysis*, Fourth edition, The English Language Book Society and Longman, New York.

Influences of Ammonia for Synthesis of 8-Hydroxyquinoline Copper(II)

ORIGINALITY REPORT

10%

SIMILARITY INDEX

7%

INTERNET SOURCES

2%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Submitted to Alliance University 4%
Student Paper
- 2** Submitted to King Mongkut's University of Technology Thonburi 3%
Student Paper
- 3** oristar.vn 2%
Internet Source
- 4** S. C. Datta. "Threshold levels of release and fixation of phosphorus: Their nature and method of determination", Communications in Soil Science and Plant Analysis, 2006 1%
Publication
- 5** K. P. Latha, V. P. Vaidya, J. Keshavayya. "Comparative Studies on Metal Complexes of 2-Acetylnaphtho[2,1-]furan Oxime and 2-Benzoylnaphtho[2,1-]furan Oxime ", Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry, 2004 1%
Publication

6

Furqan Abdulfattah, Ibrahim A. Rafukka, Sunusi M. Manladan. "Chapter 5 Trends in Characterization and Beneficiation of Non-ferrous Metallic Ores in Nigeria", Springer Science and Business Media LLC, 2020
Publication

1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off