

Binahong Leaves (*Anredera Cordifolia* Tenore Steen) Extract as an Alternative Treatment for Perineal Wound Healing of Postpartum Mothers

by Suhartono Suhartono

Submission date: 27-Oct-2020 02:59PM (UTC+0700)

Submission ID: 1427932635

File name: 11.pdf (848.01K)

Word count: 3406

Character count: 18744

© 2017 The Author(s)

This is an Open Access article distributed under the terms of the [Creative Commons Attribution 4.0 International License](#) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORIGINAL RESEARCH

ISSN: 2477-4073

BINAHONG LEAVES (ANREDERA CORDIFOLIA TENORE STEEN) EXTRACT AS AN ALTERNATIVE TREATMENT FOR PERINEAL WOUND HEALING OF POSTPARTUM MOTHERS

Dita Selvia Aditia^{1*}, Syarief Thaufik Hidayat², Nur Khafidhoh¹, Suhartono³, Ari Suwondo⁴

¹Magister Applied Midwifery Program, Health Polytechnic of Semarang, Semarang, Indonesia

²Medical Staff Group of Obstetry Gynecology Department, Dr. Kariadi Hospital, Semarang, Indonesia

³Postgraduate Epidemiology Program, Diponegoro University of Semarang Indonesia

⁴Health and Work Safety Department, Faculty of Public Health, Diponegoro University, Semarang, Indonesia

*Corresponding author:

Dita Selvia Aditia

Magister Applied Midwifery Program, Health Polytechnic of Semarang, Semarang, Indonesia

Jl. Tirta Agung, Pedalangan, Banyumanik Kota Semarang, Jawa Tengah, Indonesia (50268)

E-mail: aditiaselvia@gmail.com

Abstract

Background: Perineal wounds are at risk for infection, without a good treatment, perineal wound can harbor sepsis and become a chronic, long-term problem. Binahong leaves is considered as an alternative to accelerate perineal wound healing process.

Objective: To determine the effect of Binahong leaf extract (anredera cordifolia (tenore) steen) on perineal wound healing process in postpartum mothers.

Methods: This study employed a quasi-experimental study with control group posttest-only design. The study was conducted in the working area of Telogosari Wetan Health Center and Health Center of Kulon Kota Semarang Indonesia in January-February 2017. There were 36 respondents selected using purposive sampling assigned in the intervention group (Binahong leaves extract) and the control group (Povidone Iodine 10%). REEDA scale was used for assessing perineal healing. Data were analyzed using Mann Whitney, Chi Square and Kruskal Wallis test.

Results: Findings showed that the average day of perineal wound healing process in the Binahong leaves group was 6.11 days and in the povidone iodine group was 7.66 days. There was a statistically significant difference in the duration of wound healing between the experiment and control group with p-value 0,001 (p value <0.05).

Conclusion: Binahong leaves extract is effective in healing perineal wound in postpartum mothers. Therefore, Binahong leaves are recommended to use as an alternative treatment for the mothers.

Keywords: Binahong leaves; perineal wound; postpartum mother

INTRODUCTION

The period of the birth of the placenta and the fetal membrane is a mark of the final period of intrapartum to the return of the female reproductive tract in nonpregnant conditions. This period is also called the puerperium period, and the woman with puerperium is called puerperal. Postpartum recovery lasts about six weeks (Mohamed & El-Nagger, 2012). Recent studies of morbidity

experienced by mothers after childbirth indicated that there are some mothers continue to have problems (Marshall & Raynor, 2014). The birth canal wound may occur due to the head of the fetus being born too soon, labor not being led properly, and the need for episiotomy in order to avoid a wide perineal rupture (Prawirohardjo, 2006).

Perineal injury from episiotomy, rupture, or laceration is an area that is not easily maintained. In postpartum period, a mother will be susceptible to infection. Infections of this perineal injury can occur because the mother has lack of understanding about how to care for the correct perineal wound, what the benefits and goals of perineal wound care, no pay attention to good personal hygiene which is one of the internal factors that inhibit wound healing ([Prawirohardjo, 2006](#)). Besides, the existence of 50% of an anaerobic *Streptococcus* bacteria is the cause of infection ([Mochtar, 2011](#)).

This condition if not treated promptly will impact to the maternal mortality rate. Based on the report of the Community Health Center in Semarang City, maternal mortality in 2014 occurred as many as 33 cases from 26,992 live births or about 122.25 per 100,000 live births. Maternal mortality due to infection was 3.03%, with the condition during the puerperium was 54.55%. Based on data from the Health Department in 2015, the maternal mortality rate as many as 35 cases increased from 2013, and the incidence due to infection in 2015 in the working area of the Community of Health Center of Telogosari Wetan was 1 mother and in the working area of the Community of Health Center of Telogosari Kulon was 1 mother ([Dinkes, 2014](#)).

The wound can heal through the primary process (primary intention) that occurs when the edge of the wound is approximated by sewing it. The second healing through secondary processes (secondary intention) as the formation of granulation tissue and wound contraction ([Mochtar, 2011](#)). However, the perineal wound itself will have an impact for the mother that is discomfort disorders ([Prawirohardjo, 2006](#)).

Wound healing process consists of three stages, namely inflammatory stage, proliferative stage, maturation stage. One of the factors that can accelerate the healing of the perineal wound includes vascularization, while factors that can slow wound healing include anemia, age, other diseases, nutrition and stress. Normally the wound can heal for 6-7 days when a mother can do well treatment, on the contrary, if the wound is not

treated properly, then wound healing process becomes longer and can cause infection ([Potter & Perry, 2005](#)). Perineal wound infections due to episiotomy, rupture, or laceration are areas that are not easy to keep clean and dry, the act of cleaning the vulva may provide an opportunity for careful inspection of the perineal area.

On the other hand, perineal wound infections are also common from birth attendant by a shaman who is less concerned with hygiene during labor. If this is not treated promptly, it will lead to cervicitis, endocervical, endometritis, parametritis and palveoperitonitis and abscesses. In response, mother care can be done with local antibiotics, betadine and hygiene to prevent infection, clean the genital area every time after urination and defecation ([Bahiyatun, 2009](#)).

Currently, herbal medicines become a trend in Indonesia, one of which is Binahong leaves. The chemical content contained in the Binahong Plant (*Anredera Cordifolia* (Tenore) Steen) consists of oleanolic acid, antimicrobial, ascorbic acid, saponin triterpenoid flavonoid, and protein ([Adriani et al., 2012](#)). The content in Binahong is also shown to have anti-inflammatory, anti-fever and anti-parasitic effects. This can be an alternative treatment, as povidone iodine 10% only has antiseptic but does not have anti-inflammatory content ([Adriani et al., 2012](#)).

Clinical test on the content of Binahong leaf extract has been done in several previous studies, with the content of Binahong leaf extract, namely flavonoid, saponin, phenol, terpenoid, oleanolic acid, protein, ascorbat acid, antimicrobe ([Adriani et al., 2012](#); [Darsana, Besung, & Mahatmi, 2012](#); [Widya, Max, & Gayatri, 2013](#)).

In addition, Adnyana, et al study has done a clinical trial on Binahong leaves in patients with diabetic injuries regarding the extent of minimum inhibitory concentrations (KHM) and minimum killing (KBM) levels against *staphylococcus aureus* and *pseudomonas aeruginosa* bacteria ([Adnyana & Sukandar, 2014](#)).

Therefore, this study aimed to examine the effect of Binahong leaf extract on perineal wound healing process.

METHODS

Study Design

A quasi-experimental study with control group posttest-only design

Setting

The research was conducted in the working area of Telogosari Wetan Health Center and Health Center of Kulon Kota Semarang in January-February 2017.

Population and Sample

There were 36 respondents selected using purposive sampling assigned in the intervention group (Binahong extract) and the control group (Povidone Iodine 10%).

The inclusion criteria of the sample were a mother with a normal delivery, parity > 1, postpartum 6-10 hours, postpartum with perineal wound of degree I with no hecting, willing to use Binahong leaf extract for perineal wound treatment (intervention group), and willing to use povidone iodine 10% for the treatment of perineal wound (control group). The exclusion criteria included a mother who had a history of diabetes, BMI <18.4, and had a food obstacle.

Intervention

Binahong leaf extract was administered by dropping 5-10 drops directly on the mother's perineal wound or by dropping 10 drops in sterile gauze and then gently rubbing on the perineal wound of the mother. The dosage given to each mother was 0.5 g (500 mg) of thick extract of Binahong leaf dissolved into using aquadest of 60 ml, then shaved with a special tool until the extract blended with the solvent to obtain 0.8 % of concentration of dose extract leaf Binahong liquid. The use of

Binahong leaf extract was given 2 times per day given during the healing period of the perineal wound. Because the perineal wound in this study was perineal wound level I with no hecting, then it was considered that the wound healing process will last for 6-7 days.

While the control group received perineal wound care using povidone iodine 10% in liquid form. The povidone iodine content of 10%, given 2 times per day. The 10% of iodine administration was performed during perineal wound healing for 7-8 days. At the initial treatment of Binahong leaf extract and povidone iodine 10% provision was assisted by the researcher so that the respondents can perform as recommended by the researcher for the next days.

Instruments

To assess postpartum perineal trauma in this study, the REEDA scale was used. The REEDA scale is a tool for assessing perineal healing that was primarily developed by Davidson (Davidson, 1974) and later reviewed by Carey (Alvarenga et al., 2015). It includes five items related to the healing process: hyperemia, edema, ecchymosis, discharge and coaptation of the wound edges (Redness, Edema, Ecchymosis, Discharge, Approximation - REEDA) (see Figure 1). The length of time and level of wound density were measured and observed one day after treatment.

Ethical Consideration

This study has been approved by the Health Research Ethics Committees (K.E.P.K) of the Health Polytechnic of Semarang with number: 033 / KEPK / Poltekkes-SMG / EC / 2017. Prior to data collection, informed consent in each respondent was performed.

Data Analysis

Data were analyzed using Mann Whitney, Chi Square and Kruskal Wallis test.

Points	Redness	Oedema	Ecchymosis	Discharge	Approximation
0	None	None	None	None	Close
1	Within 0.25 cm of the incision bilaterally	Perineal, less than 1 cm from incision	Within 0.25 cm bilaterally or 0.5 cm unilaterally	Serum	Skin separation 3 mm or less
2	Within 0.5 cm of the incision bilaterally	Perineal and/or between 1 to 2 cm from the incision	Between 0.25 cm to 1 cm bilaterally or between 0.5 to 2 cm unilaterally	Serosan-guinous	Skin and subcutaneous fat separation
3	Beyond 0.5 cm of the incision bilaterally	Perineal and/or vulvar, greater than 2 cm from incision	Greater than 1 cm bilaterally or 2 cm unilaterally	Bloody, purulent	Skin, subcutaneous fat and fascial layer separation
Score					
				Total	

Figure 1 REEDA Scale (Alvarenga et al., 2015)

RESULTS

Characteristics of the respondents based on age, education, changing pad, and Hb level of postpartum mothers

Figure 2 and 3 show that the average of the respondents' age in the experiment group was 25.56 years old and in the control group was

28.83 years old. The majority of the respondents had senior high school background. The hemoglobin level in the experiment group was 10.95 gr/dL and in the control group was 10.68 gr/dL. The majority of them between the two groups changed the pads 3-4 times per day.




Figure 2 Characteristics of the respondents based on age, changing pad, and Hb level of postpartum mothers




Figure 3 Characteristics of the respondents based on education level of postpartum mothers

The average day of perineal wound healing process

As shown in the figure 4, the average day of perineal wound healing process in the experiment group was 6.11 days and in the

control group was 7.66 days. Binahong leaves extract group shows a faster wound healing process compared with healing process in the control group.



Figure 4 The average day of perineal wound healing process

Table 1 Change analysis of REEDA (*Redness, Edema, Ecchymosis, Discharge dan Approximation*) in the experiment and control group in the fifth day using Kruskal Wallis

REEDA Dimension	Experiment group	Control group	P-value
Redness	0.61±0.502	1.11±0.232	0.000
Edema	0.06±0.236	0.89±0.323	0.000
Ecchymosis	0.11±0.323	0.50±0.514	0.013
Discharge	0.11±0.323	0.44±0.511	0.028
Approximation	1.00±0.000	1.39±0.502	0.004
REEDA	1.00±0.000	1.11±0.323	0.000

Mean difference of REEDA (*Redness, Edema, Ecchymosis, Discharge dan Approximation*) between the two groups as shown in the Table 1 indicated that REEDA in the experiment group has the lower scores compared with the score in the control group. It could be said the

experiment group with Binahong leaves extract shows a better perineal wound healing. Kruskal Wallis test shows that there was a significant difference of REEDA score between the experiment and control group with p-value in each dimension <0.05.

Table 2 Analysis of difference of perineal wound healing process using Mann Whitney test

Variable	Group	p-value
Duration of wound healing	Experiment group Control group	0.000

Mann Whitney test as shown in the table 2 showed p-value 0.000 (<0.05), which indicated that there was a statistically significant difference in the duration of wound healing process between the experiment and control group.

DISCUSSION

Perineal wound is the missing or damaged part of the body tissue located between the vulva and the anus with an average length of 4 cm (Prawirohardjo, 2006). Wound care in this study is done 3 times by dripping or rubbing the liquid extract of Binahong leaves to the perineal injury. This was done referring to a good time for perineal wound care, namely when bathing, urination, and defecation (at least 3 times per day) by washing the wound area cleanly from the front of the vagina towards the back of the vagina, wipe dry and give antibiotics (Lumbessy, Abidjulu, & Paendong, 2013).

The results of this study were in accordance with previous study indicated that basically perineal wound healing would last for 6-7 days if no infection occurs (Prawirohardjo, 2006). Wound is a process in which the damaged tissue is restored to normal and there will be an inflammatory reaction divided into several phases of wound healing, namely the inflammatory phase that occurs in the 1st-4th day characterized by erythema, warmth to the skin, edema and pain. The proliferative phase

in the 5th – 20th day is the formation of granulation tissue to cover defects or injured tissue. The maturation phase occurs on the 21st day until one month even reaching the year (Adriani et al., 2012).

Findings of this study revealed that there was a significant effect of Binahong leaf on the perineal wound healing process, and there was a significant difference of REEDA score between the use of Binahong leaves and povidone iodine 10% (one day difference). Thus, it could be said that Binahong leaf has a better result in wound healing process. Binahong leaves are considered having a flavonoid content that serves as anti-inflammatory, anti-septic, antioxidants and analgesic. This is in line with previous study proved that Binahong leaf extract accelerates wound healing in infection in mice (Firdausi, 2015). Supported by Darsana, et al indicated that the higher the Binahong content used, the greater the inhibition of bacterial growth. The mechanism of inhibition of the growth of microorganisms is by destroying the peptidoglycan components in bacterial cells by alkaloids, polyphenols, saponi, flavonoids (Darsana et al., 2012). This study provides the evidence that Binahong leaves are effective to use in the healing process of perineal wound. In addition, Binahong plants are very easy to obtain and planted in home gardening without any special cares.

2. CONCLUSION

It can be concluded that Binahong leaves has a significant effect on perineal wound healing process in postpartum mothers. Therefore Binahong leaves are recommended to use as an alternative treatment for the mothers. However, further research is needed to explore the effect of Binahong leaves for mothers who have hecting perineal wound.

Declaration of Conflicting Interest

None declared. All authors contributed equally in this study.

Funding

This study was supported by Magister Applied Midwifery Program, Health Polytechnic of Semarang, Semarang, Indonesia

References

- Adnyana, I. K., & Sukandar, E. Y. (2014). Kajian aktivitas penyembuhan luka dan antibakteri Binahong (*Anredera cordifolia* (Ten.) Steenis, Pegagan (*Centella asiatica* (L.) Urban) serta kombinasinya terhadap bakteri *staphylococcus aureus* dan *pseudomonas aeruginosa* dari pasien luka kaki diabetes [Study of wound healing and antibacterias of Binahong (*Anredera cordifolia* (Ten.) Steenis, Pegagan (*Centella asiatica* (L.) Urban) and their combination againts *staphylococcus aureus* dan *pseudomonas aeruginosa* in patients with diabetic foot]. *New Bionatura*, 16(2), 78-82.
- Adriani, W. P., Ardianingtiyas, I., Wulansari, N. H., Safitri, D. N. a., Ap, I. P., & Mahanani, E. S. (2012). Uji pemanfaatan daun Binahong (*anredera cardifolia* (tenore) steenis) pada proses penyembuhan luka gingiva tikus wistar (*rattus norvegicus*) melalui pengamatan kepadatan seratbut kolagen dan ketebalan epitel [Examination of the use of Binahong (*anredera cardifolia* (tenore) steenis) leaf in the process of gingiva wound healing in *rattus norvegicus* through observation of collagen fiber density and epithelia thickness]. *Insisiva Dental Journal*, 1(2), 10-16.
- Alvarenga, M. B., Francisco, A. A., Oliveira, S. M. J. V. d., Silva, F. M. B. d., Shimoda, G. T., & Damiani, L. P. (2015). Episiotomy healing assessment: Redness, Oedema, Ecchymosis, Discharge, Approximation (REEDA) scale reliability. *Revista Latino Americana de Enfermagem*, 23(1), 162-168.
- Bahiyatun, S. P. (2009). *Buku ajar asuhan kebidanan nifas normal [Teaching book of midwifery care of normal postpartum]*. Jakarta: EGC.
- Darsana, I. G. O., Besung, I. N. K., & Mahatmi, H. (2012). Potensi daun Binahong (*Anredera cordifolia* (Tenore) Steenis) dalam menghambat pertumbuhan bakteri *escherichia coli* secara in vitro [Potential of Binahong (*Anredera cordifolia* (Tenore) Steenis) leaves in inhibiting the growth of *escherichia coli* in vitro]. *Indonesia Medicus Veterinus*, 1(3), 337-351.
- Davidson, N. (1974). REEDA: evaluating postpartum healing. *Journal of Nurse-Midwifery*, 19(2), 6-8.
- Dinkes. (2014). Health profile of Semarang City. Semarang, Indonesia: Department of Health of Semarang.
- Firdausi, R. N. (2015). *Pengaruh ekstrak etanol Binahong (Anredera cordifolia (Ten.) Steenis) terhadap profil histopatologi penyembuhan luka tikus Wistar jantan yang diinduksi aloksan [Effect of etanol Binahong (Anredera cordifolia (Ten.) Steenis) extract on hystopatology profile of wound healing in alloxan-induced male Wistar rats]*. Jember, Indonesia: Fakultas Farmasi, Universitas Jember.
- Lumbessy, M., Abidjulu, J., & Paendong, J. J. E. (2013). Uji total flavonoid pada beberapa tanaman obat tradisional di Desa Waitina Kecamatan Mangoli Timur Kabupaten Kepulauan Sula Provinsi Maluku Utara [Total examination of flavonoid in several traditional herbal plants at Desa Waitina Kecamatan Mangoli Timur Kabupaten Kepulauan Sula Provinsi Maluku Utara]. *Jurnal Mipa Unsrat Online*, 2(1), 50-55.
- Marshall, J. E., & Raynor, M. D. (2014). *Myles' textbook for midwives*. Philadelphia: Elsevier Health Sciences.
- Mochtar, R. (2011). *Sinopsis obstetri: Obstetri fisiologi, obstetri patologi [Obstetric synopsis: Physiology obstetric, patology obstetric]*. Jakarta: EGC.
- Mohamed, H. A. E., & El-Nagger, N. S. (2012). Effect of self perineal care instructions on episiotomy pain and wound healing of postpartum women. *Journal of American Science*, 8(6), 640-650.
- Potter, P. A., & Perry, A. G. (2005). *Buku ajar fundamental keperawatan: Konsep, proses, dan praktik [Textbook of fundamental of nursing: Concept, process, and practice]*. Jakarta: EGC.
- Prawirohardjo, S. (2006). *Buku acuan nasional pelayanan kesehatan maternal dan neonatal [National reference book of maternal and neonatal health services]*. Jakarta: Yayasan Bina Pustaka.
- Widya, S., Max, R. J. R., & Gayatri, C. (2013). Kandungan flavonoid dan kapasitas antioksidan total ekstrak etanol daun Binahong (*Anredera cordifolia*). Steenis [Flavonoid content and antioxidant of total extract of ethanol Binahong (*Anredera cordifolia*). Steenis] leaf. *Pharmacon*, 2(1), 18-22.

Cite this article as: Aditia, D. S., Hidayat, S. T., Khafidhoh, N., Suhartono., Suwondo, A. (2017). Binahong leaves (*anredera cordifolia tenore steen*) extract as an alternative treatment for perineal wound healing of postpartum mothers. *Belitung Nursing Journal*, 3(6), 778-783. <https://doi.org/10.33546/bnj.290>

Binahong Leaves (Anredera Cordifolia Tenore Steen) Extract as an Alternative Treatment for Perineal Wound Healing of Postpartum Mothers

ORIGINALITY REPORT

10%

SIMILARITY INDEX

4%

INTERNET SOURCES

14%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1

Dewi Susanti, Onny Setiani, Umaroh Umaroh, Kun Aristiati, Imam Djamaluddin Mashoedi, Mardiyono Mardiyono. "EFFECT OF CONSUMING TAMARIND AND TURMERIC DRINK ON THE LENGTH OF PERINEAL WOUND HEALING IN POSTPARTUM MOTHERS", Belitung Nursing Journal, 2017

Publication

4%

2

Herlina Ofiwijayanti, Syarief Thaufik Hidayat, Nur Khafidhoh. "BIRD'S NEST EXTRACT CREAM: TREATMENT FOR PERINEAL WOUND IN RATTUS NORVEGICUS", Belitung Nursing Journal, 2017

Publication

3%

3

Surya Dayyana, Suryono Suryono, Melyana Nurul Widyawati, Syarief Thaufik Hidayat. "EFFECTIVENESS OF MUSIC THERAPY ON ANXIETY AND B-ENDORPHIN LEVELS IN PRIMIGRAVIDA DURING THE THIRD STAGE

2%

OF PREGNANCY", Belitung Nursing Journal, 2017

Publication

Exclude quotes On

Exclude bibliography On

Exclude matches < 2%

Binahong Leaves (Anredera Cordifolia Tenore Steen) Extract as an Alternative Treatment for Perineal Wound Healing of Postpartum Mothers

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6