

Correlation Between Fasting Blood Glucose and Hba1c with Vitamin D in Diabetes Mellitus

by Edward Kurnia Setiawan L

Submission date: 12-Nov-2020 09:44AM (UTC+0700)

Submission ID: 1443486476

File name: Correlation_Between_Fasting.pdf (197.33K)

Word count: 3391

Character count: 17267

Correlation Between Fasting Blood Glucose and HbA1c with Vitamin D in Diabetes Mellitus

AKHMAD RAUMULFARO AKBAR¹, EDWARD KURNIASETIAWANLIMIJADI², MEITA HENDRIANINGTYAS², K. HERINUGROHO H.S³

¹Faculty of Medicine, Diponegoro University, Semarang, Indonesia

²Department of Clinical Pathology, Faculty of Medicine, Diponegoro University, Semarang, Indonesia

³Department of Internal Medicine, Faculty of Medicine, Diponegoro University, Semarang, Indonesia

Jl. Prof. Soedharto SH, Tembalang, Semarang, Central Java, Indonesia, 50275

Correspondence to Edward Kurnia Setiawan Limijadi, Email :edwardksl@fk.undip.ac.id

ABSTRACT

Background: Diabetes Mellitus (DM) is one of the largest health problems in the world. The prevalence of DM in human population has increased globally. Fasting blood glucose and HbA1c is one of parameters for diagnosing and monitoring DM. Vitamin D is a fat-soluble vitamin that has a role in glucose tolerance through its effect on insulin secretion and sensitivity in DM and the vitamin D deficiency condition sometime happened. Further more, the relation between fasting blood glucose and HbA1c with vitamin D in DM need to be investigated

Aim: To determine the correlation of fasting blood glucose and HbA1c with vitamin D in DM.

Methods: An observational analytic study with cross sectional approach in 46 subjects suitable with the inclusion and exclusion criterias. This study was conducted from April to September 2019 in *Rumah Sakit Nasional Diponegoro* (RSND) Semarang. Variables of this study were fasting blood glucose (FBG), HbA1c levels, and vitamin D (25(OH)D). Data were analyzed by Spearman and Pearson correlation. Test was significant if $p < 0,05$.

Result: Mean \pm SD of fasting blood glucose, HbA1c and vitamin D (25(OH)D) were 168,78 \pm 86,21 mg/dL, 8,46 \pm 2,20 %, and 15,85 \pm 8,75 ng/mL. The correlation between fasting blood glucose and vitamin D were not significant with $p = 0,193$; $r = 0,195$ and neither the correlation between HbA1c and vitamin D with $p = 0,153$; $r = 0,214$. **Conclusion:** Assessment of vitamin D deficiency base on the measurement of the vitamin D in diabetes melitus and this parameter don't have correlation with FBG and HbA1c levels.

Keywords: Fasting blood glucose, HbA1c, Vitamin D, Diabetes Mellitus

INTRODUCTION

Diabetes Mellitus (DM) is one of the largest problems in the world. In recent years, prevalence of DM in human population has been increased globally¹. It has been reported in "*Riset Kesehatan Dasar*" that the prevalence of DM in Indonesia for population with aged ≥ 15 years has increased from 6,9% in 2013 to 8,5% in 2018².

Diabetes mellitus is defined as a group of metabolic diseases characterized by chronic hyperglycemia caused by damage of insulin secretion, insulin action or both, causing abnormalities in the metabolism of carbohydrates, lipids, and proteins³.

The condition of hyperglycemia in DM increases reactive oxygen species (ROS) which induces oxidative stress in pancreatic β cells. This oxidative stress increase the production and release of pro-inflammatory mediators (cytokines and chemokines) which activate CYP27B1 (1 α -hydroxylase), an enzyme that cause 25 (OH) D to be converted to 1,25 (OH) D, thereby reducing level of 25 (OH) D and increasing level of 1,25 (OH) D^{4,5}.

Fasting blood glucose (FBG) was used as a parameter to distinguish diabetes mellitus patients from non-mellitus diabetes patients because fasting blood glucose has high specificity. World Health Organization also recommends measurement of HbA1c level to diagnose diabetes, HbA1c tends to be stable in the blood for a longer period and the examination can be done at any time without special preparation. According to Metcalf PA, *et al* (2017), HbA1c has correlation with macrovascular complications (cardiovascular disease and ischemic stroke) and

microvascular (retinopathy, nephropathy and neuropathy)^{6,7}.

The latest guidance from scientific bodies around the world recommends measuring 25 (OH) D serum as the test of choice for assessing vitamin D status.⁸ Serum 25 (OH) D level is the best indicator for assessing vitamin D status because 25 (OH) D level reflects the production of skin vitamin D3 and vitamin D (D2 and D3) from food. 25 (OH) D also has a long half-life in circulation which is 3-4 weeks. Although the active metabolite of vitamin D is 1,25 (OH) 2D, measurement of serum 1,25 (OH) 2D level is not recommended to determine vitamin D status because the half-life in short circulation is 4-6 hours and its level in serum is very low when compared with a level of 25 (OH) D. When vitamin D deficiency occurs, parathyroid hormone secretion will increase as a compensatory response that will stimulate the kidneys to increase production of 1,25 (OH) D so that when vitamin D deficiency occurs, 25 (OH) D level are decreased, while level of 1,25 (OH) 2D is maintained at normal or even elevated level⁹.

Until now then, research on the relationship of fasting blood glucose and HbA1c level with vitamin D (25 (OH) D) levels in diabetes mellitus especially in Indonesia is still limited, and also there were differences in previous studies between these variables. Therefore, it needs advanced investigations.

MATERIALS AND METHODS

An observational study with cross sectional approach was conducted in Rumah Sakit Nasional Diponegoro (RSND) Semarang on April to September 2019. The subjects of the

study were 46 diabetes mellitus patients who met inclusion and exclusion criterias by consecutive sampling. The inclusion criterias were diabetic patients, >18 years old, normal body temperature, normal liver function test, and willing to be respondent, while the exclusion criterias were anemia, pregnant, smoker, consuming vitamin D supplements, patient with history of chronic liver, patient with history of chronic kidney disease, and stroke patient.

Data in this study were identity of the patients, weight, height, sport activity, blood pressure, percentage of body fat, hemoglobin levels, AST, ALT, FBG, HbA1c levels, and 25(OH)D levels. Blood sampling, blood pressure and the percentage of body fat measurement were done at laboratory of RSND. The measurement of hemoglobin levels, AST, ALT, FBG, HbA1c, and 25(OH)D was done at laboratory of RSND, GAKI laboratory faculty of medicine UNDIP, and laboratory of Prodia.

The percentage of body fat was done by omron body composition monitor HBF-375. Fasting blood glucose were measured by GOD-PAP method on automatic clinical chemistry analyzer architech C-4000, normal level of FBG is ≤ 100 mg/dl. HbA1c levels were measured by high performance liquid chromatography (HPLC) on a D-10 system, normal level of HbA1c is less than 5,7%. 25(OH)D levels were measured by *Enzyme-Linked Immuno Sorbent Assay* (ELISA) method, normal level of 25(OH) is 30-100ng/mL.

Data analysis was performed by statistical programme computer. Normality test was *Saphiro-Wilk* and data that were not normally distributed carried out data transformation. *Pearson* correlation test was performed to test correlation of HbA1c levels with vitamin D (25(OH)D) levels and *Spearman* correlation was performed to test relation of fasting blood glucose with vitamin D (25(OH)D). $p > 0,05$ was significant.

Ethical clearance was obtained from the Medical and Health Research Ethics Commission (KEPK) of the Faculty of Medicine, Diponegoro University. Subjects were given a brief explanation of the purpose, benefits, research protocols, and possible side effects, and written informed consent was done.

RESULT

The population of this study were diabetes mellitus patients in RumahSakit Nasional Diponegoro (RSND) Semarang. This study was conducted from April to September 2019 in RSND Semarang and the samples were examined in laboratory of RSND Semarang. There were 79 subjects who were willing to be sampled, but only 46 subjects met the inclusion and exclusion criteria. Descriptive are shown in table 1.

The gender distribution in this study was 17 (36,96%) males and 29 (63,04%) females. The average age of respondents was 57.85 ± 11.58 years. The youngest age in this study was 34 years and the oldest age was 79 years. There were 16 (34,78%) subjects who routinely did sport activities and there were 30 (65,22%) subjects who did not routinely do sport activities.

Table 1. Characteristics of subjects

Parameter	Mean \pm SD
Age (years)	57,85 \pm 11,58
Gender (%)	
Male	36,96
Female	63,04
Sport activity (%)	
Yes	34,78
No	65,22
Blood pressure (mmHg)	
Sistol	131,89 \pm 11,73
Diastol	82,83 \pm 6,08
Percentage of body fat (%)	32,56 \pm 6,48
Male	26,76 \pm 6,77
Female	35,56 \pm 5,03
Body weight (kilograms)	63,19 \pm 11,54
Height (centimeters)	156,29 \pm 7,87
Body Mass Index (kg/m ²)	25,72 \pm 4,15
Hemoglobin levels (mg/dL)	13,18 \pm 0,95
SGOT (mg/dL)	19,37 \pm 5,2
SGPT (mg/dL)	18,65 \pm 5,33
Fasting blood glucose (mg/dl)	168,78 \pm 86,21
HbA1c levels (%)	8,46 \pm 2,20
Vitamin D (25(OH)D) levels (ng/mL)	15,85 \pm 8,75

4
The percentage of body fat in female is higher than male with BMI obtained in normal to overweight range. Fasting blood glucose levels in this study population ranged from normal to high, where the proportion of normal FBG level was less than those of high levels. There were 21 (45,65%) subjects with normal FBG and 25 (54,35%) subjects with increased FBG, while HbA1c levels in all subjects of this study were found to be increasing from normal. 25(OH)D levels in this study were found decreased based on normal values.

Table 2: Correlation test results

Variable	Vitamin D (25(OH)D)	
	r	p
HbA1c	0,214	0,153 ^a
Fasting blood glucose	0,195	0,193 ^b

^a*Pearson* correlation test

^b*Spearman* correlation test

Data was analyzed by parametric *Pearson* correlation test for HbA1c and vitamin D (25(OH)D) and non-parametric *Spearman* correlation test for fasting glucose test and vitamin D (25(OH)D). Based on the *Pearson* correlation test between HbA1c and vitamin D (25(OH)D), the result shows there was no significant correlation between HbA1c and vitamin D (25(OH)D). Based on the *Spearman* correlation test between fasting blood glucose and vitamin D (25(OH)D), the result shows there was no significant correlation between fasting blood glucose and vitamin D (25(OH)D) as shown in table 2.

DISCUSSION

1
The result of *Spearman* correlation test between fasting blood glucose with vitamin D (25(OH)D) showed that there was no significant correlation. The study conducted by Alkhatatbeh in 2018 found that there was no significant relationship between fasting blood glucose with vitamin D (25(OH)D) in diabetes mellitus¹⁰. Tandon in 2014 also stated there was no significant correlation between fasting

ORIGINAL ARTICLE

blood glucose with vitamin D (25(OH)D) in diabetes mellitus¹¹.

The result of this study differs from some previous studies. Study conducted by Pannu in 2017 found there was a negative relationship between fasting blood glucose and vitamin D (25(OH)D) in DM¹². Another study in 2017 by Kanakaraju shows there was a negative relationship between vitamin D (25(OH)D) with blood glucose parameters¹³. The same result was also mentioned by Bhatt in 2018 there was a negative relationship between fasting blood glucose with vitamin D (25(OH)D) in pre-diabetic population^{14,23}.

Insignificant relationship between fasting blood glucose and vitamin D (25(OH)D) can be caused by several factors. One of them is the percentage of body fat. Individuals with a high percentage of body fat can reduce level of vitamin D (25(OH)D) because excess fat tissue can store fat-soluble vitamin D thus it reduces vitamin D is released in the bloodstream and serum 25(OH)D level become low^{15,16}.

The use of drugs such as glucocorticoid drugs, anti-hypertension may affect vitamin D (25(OH)D) level, glucocorticoids may affect function of osteoblasts and osteoclasts thus it reduces vitamin D levels. Some glucocorticoids also increase degradation of 25(OH)D and 1,25(OH)D due activation of *pregnane X receptors*¹⁷. In addition, physical activity also affects plasma vitamin D concentrations, a study conducted by Fernandes in 2017 showed that the concentration plasma vitamin D increases with physical activity¹⁸. Research conducted by Sakungin 2018 showed that physical activity may increase insulin sensitivity and increases glucose uptake by body tissues thus it reduce blood glucose level¹⁹.

The result of *Pearson* correlation test between HbA1c with vitamin D (25(OH)D) showed there was no significant correlation. The study conducted by Kumar in 2017 showed that diabetes mellitus could not be proven to have a significant relationship between HbA1c and vitamin D (25(OH)D) levels in people with diabetes mellitus²⁰. Coskun in 2012 also stated there was no significant relationship between HbA1c with vitamin D (25(OH)D) levels in diabetes mellitus²¹.

The result of this study differs from some previous studies. Study conducted by Al-Timimi in 2013 stated that patients with poor glycemic control (HbA1c) had lower serum 25(OH)D²². In another study in 2017 by Pannu shows that high levels serum 25(OH)D was associated with a low HbA1c levels¹². The same result was also mentioned by Buhary in 2017 that there was a negative relationship between HbA1c and vitamin D (25(OH)D) in diabetes mellitus²³.

Insignificant relationship between HbA1c and vitamin D (25(OH)D) can be caused by several factors. One of them is exposure of sun's ultraviolet B (UV B) rays, UV B has an important role in the synthesis of vitamin D. UV B play a role in breaks B ring at 7-dehydrocholesterol into pre-D3 which is isomerized into D3 by irradiation of UV B. Thus, the lack of exposure from UV B sunlight can interfere with the synthesis of vitamin D²⁴.

Gender can also affect the level of vitamin D (25 (OH) D) because men have outside activities more frequently thus they are more exposed to sunlight, usage of sunblock

more frequent in women can also reduce vitamin D synthesis, and also women have a higher percentage of fat than men thus they have lower concentrations of vitamin D (25 (OH) D) than men. Furthermore, age also affects vitamin D level due to reduced production of vitamin D in the skin, lack of exposure to UV B sunlight, reduced food intake, reduced skin thickness, impaired absorption in the intestine, and reduced hydroxylation in the liver and kidneys. In the study conducted by Al-Horani in 2016 showed that there was a relationship between gender, age, and body mass index with vitamin D (25(OH)D) levels²⁵.

Another factors that can influence HbA1c level in this study is glycation factors which can be influenced by the levels of consumption of vitamin C, the levels of consumption of vitamin E, and alcoholism, as well as Yunika (2017)⁹. In the study conducted by Ardekani in 2007 showed that consumption of vitamin D decreases serum level of fasting blood glucose, LDL, and HbA1c significantly²⁶.

Limitations in this study are the current study subjects, they are not distinguished based on physical activity, gender and age. This study also did not distinguish DM subjects based on BMI or body fat percentage. Medicines were consumed by patients, where the use of several drugs such as anti-hypertensive drugs, glucocorticoids, vitamin C supplements, vitamin E supplements can influence the results of the study.

CONCLUSION

Relationship between FBG and HbA1c levels with vitamin D (25(OH)D) level in diabetes mellitus patients are not significant. Assessment of vitamin D deficiency base on the measurement of the vitamin D in diabetes mellitus. The next study needs to be adjusted with interference factors related to variables detail that mention in limitation.

REFERENCES

1. International Diabetes Federation. IDF Diabetes Atlas. 8 ed. International Diabetes Federation. Brussel: International Diabetes Federation; 2017.
2. Kesehatan BP dan P. Hasil Utama Riset Kesehatan Dasar. Kementerian Kesehatan Republik Indonesia. 2018;1-100.
3. American Diabetes Association. Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care*. 2010;33(1):S62-9.
4. Akash MSH, Rehman K, Chen S. Role of Inflammatory Mechanisms in Pathogenesis of Type 2 Diabetes Mellitus. *J Cell Biochem*. 2013;114(3):525-31.
5. Mangin M, Sinha R, Fincher K. Inflammation and vitamin D: the infection connection. *Inflamm Res*. 2014;63(10):803-19.
6. Rathod SD, Crampin AC, Musicha C, Kayuni N, Banda L, Saul J, et al. Glycated haemoglobin A1c (HbA1c) for detection of diabetes mellitus and impaired fasting glucose in Malawi: A diagnostic accuracy study. *BMJ Open*. 2018;8(5):1-8.
7. Metcalf PA, Kyle C, Kenealy T, Jackson RT. HbA1c in relation to incident diabetes and diabetes-related complications in non-diabetic adults at baseline. *J Diabetes Complications*. 2017;31(5):814-23.
8. Herrmann M, Farrell CJL, Pusceddu I, Fabregat-Cabello N, Cavalier E. Assessment of Vitamin D status - A changing landscape. *Clin Chem Lab Med*. 2017;55(1):3-26.
9. Puspa Dewi, Yunika. (2017). An Overview: Vitamin D. Available at: <https://www.researchgate.net/publication/319997190>.

- Accessed October 5, 2019.
10. Alkhatatbeh MJ, Abdul-Razzak KK. Association between serum 25-hydroxyvitamin D, hemoglobin A1c and fasting blood glucose levels in adults with diabetes mellitus. *Biomed Reports*. 2018;9(6):523–30.
 11. Tandon V, Gillani Z, Khajuria V, Mahajan S, Mahajan A, Raina K, et al. Prevalence of vitamin d deficiency among Indian menopausal women and its correlation with diabetes: A first Indian cross sectional data. *J Midlife Health*. 2014;5(3):121.
 12. Pannu PK, Piers LS, Soares MJ, Zhao Y, Ansari Z. Vitamin D Status is Inversely Associated with markers of risk for type 2 diabetes: A population based study in Victoria, Australia. *PLoS One*. 2017;12(6):e0178825.
 13. Kanakaraju K, Ranganathan RS, Shankar R. Correlation of Vitamin D3 Levels and the Blood Sugar Parameters among the Patients with type 2 Diabetes Mellitus. 2017;4(4):844–7.
 14. Bhatt SP, Misra A, Gulati S, Singh N, Pandey RM. Lower Vitamin D levels are associated with higher blood glucose levels in Asian Indian women with pre-diabetes: A population-based cross-sectional study in North India. *BMJ Open Diabetes Res Care*. 2018;6(1):1–9.
 15. Sundari LPR. Defisiensi Vitamin D Pada Obesitas. *Sport Fit J*. 2018;6(1):1–5.
 16. Zhang M, Li P, Zhu Y, Chang H, Wang X, Liu W, et al. Higher visceral fat area increases the risk of vitamin D insufficiency and deficiency in Chinese adults. *Nutr Metab*. 2015;12(1):50.
 17. Gröber U, Kisters K. Influence of drugs on vitamin D and calcium metabolism. *Dermatoendocrinol*. 2012;4(2):158–66.
 18. Fernandes MR, Dos Reis Barreto W. Association Between Physical Activity and Vitamin D: A Narrative Literature Review. *Rev Assoc Med Bras*. 2017;63(6):550–6.
 19. Sakung JM, Sirajuddin S, Zulkifli A, Rahman SA, Sudargo T. Physical Activity is Associated with Lower Blood Glucose Level in High School Teachers in Palu, Indonesia. *Int J Community Med Public Heal*. 2018;5(8):3176.
 20. Akshay Kumar S V., Nanda SK, Bharathy N, Ravichandran K, Dinakaran A, Ray L. Evaluation of vitamin D status and its correlation with glycated haemoglobin in type 2 diabetes mellitus. *Biomed Res*. 2017;28(1):66–70.
 21. Coskun H, Aydin Y, Kutlucan A, Yildirim H, Kudas O, Kir S, et al. Vitamin D3 deficiency effect on the HbA1c in type 2 diabetic Population. 15th Int 14th Eur Congr Endocrinol. 2012;29:733.
 22. Al-Timimi DJ, Ali AF. Serum 25(OH) D in diabetes mellitus type 2: Relation to glycaemic control. *J Clin Diagnostic Res*. 2013;7(12):2686–8.
 23. Buhary BM, Almohareb O, Aljohani N, Alrajhi S, Elkaisi S, Sherbeeni S, et al. Association of Glycosylated Hemoglobin Levels With Vitamin D Status. *J Clin Med Res*. 2017;9(12):1013–8.
 24. Tsiaras WG, Weinstock MA. Factors influencing vitamin d status. *Acta Derm Venereol*. 2011;91(2):115–24.
 25. Al-Horani H, Abu Dayyih W, Mallah E, Hamad M, Mima M, Awad R, et al. Nationality, gender, age, and body mass index influences on Vitamin D concentration among elderly patients and young Iraqi and Jordanian in Jordan. *Biochem Res Int*. 2016:1-8.
 26. Afkhami-Ardekani M, Shojaoddiny-Ardekani A. Effect of vitamin C on blood glucose, serum lipids and serum insulin in type 2 diabetes patients. *Indian J Med Res*. 2007;126(5):471–4.

Correlation Between Fasting Blood Glucose and Hba1c with Vitamin D in Diabetes Mellitus

ORIGINALITY REPORT

17 %	%	17 %	%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

- 1** Ghadeer K. Al-Shaikh, Gehan H. Ibrahim, Amel A. Fayed, Hazem Al-Mandeel. "Impact of vitamin D deficiency on maternal and birth outcomes in the Saudi population: a cross-sectional study", BMC Pregnancy and Childbirth, 2016
Publication **3%**
 - 2** Mangin, Meg, Rebecca Sinha, and Kelly Fincher. "Inflammation and vitamin D: the infection connection", Inflammation Research, 2014.
Publication **1%**
 - 3** Rabenberg, Martina, Christa Scheidt-Nave, Markus A. Busch, Nina Rieckmann, Birte Hintzpeter, and Gert B.M. Mensink. "Vitamin D status among adults in Germany – results from the German Health Interview and Examination Survey for Adults (DEGS1)", BMC Public Health, 2015.
Publication **1%**
-

4

Sunil J Wimalawansa. "Vitamin D Deficiency is a Surrogate Marker for Visceral Fat Content, Metabolic Syndrome, Type 2 Diabetes, and Future Metabolic Complications", Journal of Diabetes, Metabolic Disorders & Control, 2016

Publication

1%

5

Akash, Muhammad Sajid Hamid, Kanwal Rehman, and Shuqing Chen. "Role of inflammatory mechanisms in pathogenesis of type 2 diabetes mellitus", Journal of Cellular Biochemistry, 2013.

Publication

1%

6

Dallak. "*In vivo*, Acute, Normo-hypoglycemic, Antihyperglycemic, Insulinotropic Actions of Orally Administered Ethanol Extract of *Citrullus colocynthis* (L.) Schrab Pulp", American Journal of Biochemistry and Biotechnology, 2009

Publication

1%

7

Antara Sen, Kallol Bhattacharjee, Giridhari Kar, Chandra Prakash Thakur. "SPECTRUM OF VITAMIN D IN TYPE 2 DIABETES MELLITUS: A HOSPITAL BASED STUDY", Journal of Evidence Based Medicine and Healthcare, 2018

Publication

1%

8

Alireza Khabbazi, Morteza Ghojazadeh, Sakineh Hajebrahimi, Zeinab Nikniaz. "Relationship Between Vitamin D Level and Bechcet's

1%

Disease Activity: A Systematic Review and Meta-Analysis", International Journal for Vitamin and Nutrition Research, 2019

Publication

9

Molly Bond, Natalie Moll, Alicia Rosello, Rod Bond et al. "Vitamin D status in chronic tic-disorder and comorbid obsessive-compulsive disorder and attention-deficit/hyperactivity disorder: A pan-European study", Cold Spring Harbor Laboratory, 2019

Publication

1%

10

"Calcium in Internal Medicine", Springer Science and Business Media LLC, 2002

Publication

1%

11

Hanan Al-Horani, Wael Abu Dayyih, Eyad Mallah, Mohammed Hamad, Mohammad Mima, Riad Awad, Tawfiq Arafat. "Nationality, Gender, Age, and Body Mass Index Influences on Vitamin D Concentration among Elderly Patients and Young Iraqi and Jordanian in Jordan", Biochemistry Research International, 2016

Publication

<1%

12

U. Ozuguz, S. Oruc, M. S. Ulu, H. Demirbas, A. Acay, B. Coker, B. Beyazit, M. Yaman, T. Koken. "Does vitamin D have any role in the improvement of diabetic peripheral neuropathy in type 1 diabetic patients?", Journal of

<1%

13

Mzwandile A. Mabhala, Adetoyosi Babanumi, Anthony Olagunju, Eloho Akata, Asmait Yohannes. "Evaluating the Use of Vitamin D Supplementation to Improve Glycaemic Outcome in Type 2 Diabetes Mellitus Patients: A Systematic Review of Evidence", Journal of Diabetes Mellitus, 2017

Publication

<1%

14

Klenk, J., K. Rapp, M. D. Denking, G. Nagel, T. Nikolaus, R. Peter, W. Koenig, B. O. Boehm, and D. Rothenbacher. "Seasonality of vitamin D status in older people in Southern Germany: implications for assessment", Age and Ageing, 2013.

Publication

<1%

15

Jain, S K, D Micinski, L Huning, G Kahlon, P F Bass, and S N Levine. "Vitamin D and L-cysteine levels correlate positively with GSH and negatively with insulin resistance levels in the blood of type 2 diabetic patients", European Journal of Clinical Nutrition, 2014.

Publication

<1%

16

Mohammed I. Oraby, Mona Abu Srie, Sanaa Abdelshafy, Eman Elfar. "Diabetic peripheral neuropathy: the potential role of vitamin D

<1%

deficiency", The Egyptian Journal of Neurology, Psychiatry and Neurosurgery, 2019

Publication

17

Volmer, Dietrich A., Luana R.B.C. Mendes, and Caroline S. Stokes. "Analysis of vitamin D metabolic markers by mass spectrometry: Current techniques, limitations of the "gold standard" method, and anticipated future directions : MEASURING VITAMIN D METABOLIC MARKERS BY MASS SPECTROMETRY", Mass Spectrometry Reviews, 2013.

Publication

<1%

18

"Book of Abstracts", Climacteric, 2009

Publication

<1%

19

Rene F. Chun, Albert Shieh, Carter Gottlieb, Vahe Yacoubian, Jeffrey Wang, Martin Hewison, John S. Adams. "Vitamin D Binding Protein and the Biological Activity of Vitamin D", Frontiers in Endocrinology, 2019

Publication

<1%

20

Mahdieh Golzarand, Bruce W. Hollis, Parvin Mirmiran, Carol L. Wagner, Sakineh Shab-Bidar. "Vitamin D supplementation and body fat mass: a systematic review and meta-analysis", European Journal of Clinical Nutrition, 2018

Publication

<1%

21

Jing Xue, Sarah A. Schoenrock, William Valdar, Lisa M. Tarantino, Folami Y. Ideraabdullah.

"Maternal vitamin D depletion alters DNA methylation at imprinted loci in multiple generations", *Clinical Epigenetics*, 2016

Publication

<1%

22

Wei Zhu, Daniel P. Heil. "Associations of vitamin D status with markers of metabolic health: A community-based study in Shanghai, China", *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 2018

Publication

<1%

23

Tânia Valladares, Maria Regina Cardoso, Jose Mendes Aldrighi. "Higher serum levels of vitamin D are associated with lower blood glucose levels", *Menopause*, 2019

Publication

<1%

24

Surya Prakash Bhatt, Anoop Misra, Seema Gulati, Naamrata Singh, Ravindra Mohan Pandey. "Lower vitamin D levels are associated with higher blood glucose levels in Asian Indian women with pre-diabetes: a population-based cross-sectional study in North India", *BMJ Open Diabetes Research & Care*, 2018

Publication

<1%

25

Christopher T. Sempos, Annemieke C. Heijboer, Daniel D. Bikle, Jens Bollerslev et al. " Vitamin

<1%

D Assays and the Definition of Hypovitaminosis D: Results from the 1 International Conference on Controversies in Vitamin D ", British Journal of Clinical Pharmacology, 2018

Publication

26

Kostas Archontogeorgis, Nikolaos Papanas, Evangelos C. Rizos, Evangelia Nena et al. "Reduced Serum Vitamin D Levels Are Associated with Insulin Resistance in Patients with Obstructive Sleep Apnea Syndrome", Medicina, 2019

Publication

<1%

27

Maria Creusa Rolim, Bárbara Mendes Santos, Gildasio Conceição, Paulo Novis Rocha. "Relationship between vitamin D status, glycemic control and cardiovascular risk factors in Brazilians with type 2 diabetes mellitus", Diabetology & Metabolic Syndrome, 2016

Publication

<1%

28

Sunali Padhi, Subham Suvankar, Venketesh K. Panda, Abhijit Pati, Aditya K Panda. "Lower levels of vitamin D are associated with SARS-CoV-2 infection and mortality in the Indian population: an observational study", International Immunopharmacology, 2020

Publication

<1%

29

Eyad M Mallah. "Plasma concentrations of 25-

<1%

hydroxyvitamin D among Jordanians: Effect of biological and habitual factors on vitamin D status", BMC Clinical Pathology, 2011

Publication

30

G Karatekin, A Kaya, Ö Salihoğlu, H Balci, A Nuhoğlu. "Association of subclinical vitamin D deficiency in newborns with acute lower respiratory infection and their mothers", European Journal of Clinical Nutrition, 2007

Publication

<1%

31

Rukshana Shroff, Craig Knott, Lesley Rees. "The virtues of vitamin D—but how much is too much?", Pediatric Nephrology, 2010

Publication

<1%

32

Jeong Ki Paek, Ji Hye Won, Hye Rin Shin, Dea Youn Kim, Kyunam Kim, Seon Yeong Lee. "Association between Vitamin D Concentration and Visceral Fat Area in Healthy Korean Adults", Korean Journal of Health Promotion, 2017

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Correlation Between Fasting Blood Glucose and Hba1c with Vitamin D in Diabetes Mellitus

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
