

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : The correlation between galectin-3, creatinine and uric acid on stage v chronic renal failure

Jumlah Penulis : 3 orang

Status Pengusul : **Indranila K S, Guruh A I, Meita H**

Identitas Jurnal Ilmiah : a. Nama Jurnal : **INDONESIAN JOURNAL OF CLINICAL PATHOLOGY AND MEDICAL LABORATORY**

b. Nomor ISSN : p- 0854-4263, e- 2477-4685

c. Vol, Nomor, halaman : Vol 25, No. 1 p:104-114

d. Edisi : November 2018

e. Penerbit : Indonesian Association of Clinical Pathologists

f. Jumlah halaman : 7

g. DOI artikel (jika ada) : <http://dx.doi.org/10.24293/ijcpml.v25i1.1516>

h. Alamat web jurnal : <https://www.indonesianjournalofclinicalpathology.org/index.php/patologi/article/view/1516/1073>

i. Terindeks di : **SINTA 2**

j. Online Turnitin : [https://doc-](https://doc-pak.undip.ac.id/4994/1/Turnitin_THE_CORRELATION_BETWEEN.pdf)

Kategori Publikasi Jurnal Ilmiah : ☐ Jurnal Ilmiah Internasional / Internasional Bereputasi **

(beri ✓ pada kategori yang tepat) ☒ **Jurnal Ilmiah Nasional Terakreditasi**

☐ Jurnal Ilmiah Nasional/Nasional

Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah			Nilai Akhir Yang Diperoleh
	Internasional / Internasional Bereputasi ** <input type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional *** <input type="checkbox"/>	
a. Kelengkapan unsur isi artikel (10%)		2,5		2,5
b. Ruang lingkup dan kedalaman pembahasan (30%)		7,5		6,5
c. Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)		7,5		6,5
d. Kelengkapan unsur dan kualitas terbitan/ jurnal (30%)		7,5		7,5
Total = (100%)		25		23
Nilai Pengusul =	(60% x 23) = 13,8			

Catatan penilaian Artikel oleh Reviewer :

- Kelengkapan unsur isi artikel** : Unsur isi artikel lengkap Abstrak, Pendahuluan, Hasil & Diskusi, Referensi. Abstrak ditulis kurang terstruktur sedangkan unsur lain cukup baik, hasil dan pembahasan tidak dipisah sehingga kurang bisa fokus
- Ruang lingkup dan kedalaman pembahasan** : Ruang lingkup keilmuan sesuai pengusul. Pembahasan cukup dalam disertai referensi yang sesuai
- Kecukupan dan kemutakhiran data/informasi dan metodologi**: Penelitian observasional dengan desain belah lintang. Data dikumpulkan dengan teknologi yang baik dan dianalisis dengan baik pula
- Kelengkapan unsur dan kualitas terbitan/ jurnal**: Jurnal nasional terakreditasi Sinta 2

Semarang,
 Reviewer

Prof. Dr. dr. Tri Nur Kristina, DMM, M.Kes
 NIP. 19590527 198603 2 001
 Unit Kerja : Fakultas Kedokteran Undip
 Bidang Ilmu : Ilmu Kedokteran
 Jabatan : Guru Besar

LEMBAR

HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Judul Jurnal Ilmiah (Artikel) : THE CORRELATION BETWEEN GALECTIN-3, CREATININE AND URIC ACID ON STAGE V CHRONIC RENAL FAILURE

Jumlah Penulis : 3 orang

Status Pengusul : Indranila K S, Guruh A I, Meita H

Identitas Jurnal Ilmiah :

- a. Nama Jurnal : INDONESIA JOURNAL OF CLINICAL PATHOLOGY
- b. Nomor ISSN : p-ISSN: 0854-4263e-ISSN: 2477-4685
- c. Volume nomor bulan tahun : Vol 25, 104-114
- d. Penerbit : Indonesian Journal of Clinical Pathology and Medical Laboratory
- e. DOI artikel (Jika ada) : <http://dx.doi.org/10.24293/ijcpml.v25i1.1516>
- f. Alamat web Jurnal : <https://www.indonesianjournalofclinicalpathology.org/index.php/patologi/article/view/1516/1073>
- h. Terindeks di : SINTA 2
- i. Online turnitin : https://doc-pak.undip.ac.id/4994/1/Turnitin_THE_CORRELATION_BETWEEN.pdf

Kategori Publikasi Jurnal Ilmiah : ☐ Jurnal Ilmiah Internasional / internasional bereputasi *

(beri ✓ pada kategori yang tepat) ☒ Jurnal Ilmiah Nasional Terakreditasi

☐ Jurnal Ilmiah Nasional/ Nasional terindeks di DOAJ CABI COPERNICUS** Tidak Terakreditasi

Hasil Penilaian Peer Review :

NO	KOMPONEN YG DINILAI	Nilai Maksimal Jurnal Ilmiah			
		Internasional/internasional bereputasi	Nasional Terakreditasi	Nasional ***	Nilai Akhir Yang Diperoleh
a	Kelengkapan unsur isi artikel (10%)		25		2,2
b	Ruang lingkup dan kedalaman pembahasan (30%)		7,5		7,3
c	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)		7,5		7,0
d	Kelengkapan unsur dan kualitas terbitan/ jurnal (30%)		7,5		7,0
Nilai Total = (100%)			25		23,5
Nilai Pengusul =				60% x =	14,1

Catatan Penilaian artikel oleh Reviewer

- a. Kelengkapan unsur isi artikel : abstrak lengkap dan Area informatif. Analisis teras, pendahuluan, metode, hasil dan diskusi. Kesimpulan, saran dan rekomendasi. Kesimpulan, saran dan rekomendasi. Kesimpulan, saran dan rekomendasi.
- b. Ruang lingkup dan kedalaman pembahasan : Ruang lingkup dan kedalaman pembahasan. Ruang lingkup dan kedalaman pembahasan. Ruang lingkup dan kedalaman pembahasan.
- c. Kecukupan dan kemutakhiran data/informasi dan metodologi : Kecukupan dan kemutakhiran data/informasi dan metodologi. Kecukupan dan kemutakhiran data/informasi dan metodologi. Kecukupan dan kemutakhiran data/informasi dan metodologi.
- d. Kelengkapan unsur dan kualitas terbitan/ jurnal : Kelengkapan unsur dan kualitas terbitan/ jurnal. Kelengkapan unsur dan kualitas terbitan/ jurnal. Kelengkapan unsur dan kualitas terbitan/ jurnal.

Semarang, 01 September 2020

Reviewer 2

Prof. Dr. drg. Oedijani, M.S.

NIP 194902091979012001

Unit kerja: Fakultas Kedokteran Undip

Bidang ilmu : Ilmu Kedokteran

Jabatan/pangkat: Guru Besar

Vol. 25, No. 1, November 2018

p-ISSN 0854-4263
e-ISSN 2477-4685

INDONESIAN JOURNAL OF
**Clinical Pathology and
Medical Laboratory**

Majalah Patologi Klinik Indonesia dan Laboratorium Medik

IJCP & ML
(Maj. Pat. Klin.
Indonesia & Lab. Med.)

Vol. 25

No. 1

Page 1 - 128

Surabaya
November 2018

ISSN
0854-4263

Published by Indonesian Association of Clinical Pathologists

Accredited No. 36a/E/KPT/2016, Tanggal 23 Mei 2016



INDONESIAN JOURNAL OF Clinical Pathology and Medical Laboratory

Majalah Patologi Klinik Indonesia dan Laboratorium Medik

p-ISSN 0854-4263
e-ISSN 4277-4685

Published by
Indonesian Association of Clinical
Pathologist and Medical Laboratory

[HOME](#) [ABOUT](#) [LOGIN](#) [REGISTER](#) [SEARCH](#) [CURRENT](#) [ARCHIVES](#) [ANNOUNCEMENTS](#) [AUTHOR GUIDELINE](#) [EDITORIAL TEAM](#)

[Home](#) > [About the Journal](#) > [Editorial Team](#)

Editorial Team

Editor-in-chief

Puspa Wardhani, Editor-in-chief Indonesian Journal of Clinical Pathology and Medical Laboratory, Indonesia, Indonesia

Editorial Boards

Ida Parwati, Department of Clinical Pathology, Faculty of Medicine, Padjajaran University, Bandung, Indonesia, Indonesia

Yulia Nadar Indrasari, Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo Hospital, Surabaya, Indonesia, Indonesia

Nuri Diah Indrasari, Department of Clinical Pathology, Faculty of Medicine, Indonesia University/Dr. Cipto Mangunkusumo, Jakarta, Indonesia, Indonesia

Efrida Efrida, Department of Clinical Pathology, Faculty of Medicine, Andalas Univesity/M.Djamil Hospital, Padang, Indonesia, Indonesia

Puspa Wardhani, Editor-in-chief Indonesian Journal of Clinical Pathology and Medical Laboratory, Indonesia, Indonesia

Nyoman Suci Widiastuti, Department of Clinical Pathology, Faculty of Medicine, Diponegoro University/Dr. Kariadi, Semarang, Indonesia, Indonesia

Ulung Bahrin, Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University/Dr. Wahidin Sudirohusodo Hospital, Makassar, Indonesia, Indonesia

Budi Mulyono, Department of Clinical Pathology, Faculty of Medicine, Gajah Mada University/Dr. Sardjito Hospital, Yogyakarta, Indonesia, Indonesia

Editorial Assistant

Dian Wahyu Utami, Indonesian Journal of Clinical Pathology and Medical Laboratory, Indonesia, Indonesia

Peer-Reviewers

Tony Badrick, Royal College of Pathologists of Australia Quality Assurance Programs, Australia

Hans Vrielink, Sanquin Blood Supply, Plesmanlaan 125, 1066 CX Amsterdam, the Netherlands, Netherlands

Wendy Erber, The University of Western Australia, Australia

Teguh Triyono, Department of Clinical Pathology, Faculty of Medicine, Gadjah Mada University/Dr. Sardjito Hospital, Yogyakarta, Indonesia, Indonesia

Che Maraina Bt Che Hussin, Universiti Sains Malaysia, Malaysia

Zilfalil Bin Alwi, Universiti Sains Malaysia, Malaysia

Kadek Mulyantari, Department of Clinical Pathology, Faculty of Medicine, Udayana University/Sanglah General Hospital Bali, Indonesia, Indonesia

Prihatini prihatini, Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo Hospital, Surabaya, Indonesia, Indonesia

Adi Koesoema Aman, Department of Clinical Pathology, Faculty of Medicine, Sumatera Utara University/Adam Malik Hospital, Medan, Indonesia, Indonesia

Lusi Oka Wardhani, Department of Clinical Pathology, Faculty of Medicine, Sebelas Maret University/Dr. Moewardi, Surakarta, Indonesia, Indonesia

Basti Andriyoko, Department of Clinical Pathology, Faculty of Medicine, Padjajaran University/Dr. Hasan Sadikin Hospital, Bandung, Indonesia, Indonesia

Paulus Budiono Notopuro, Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo Hospital, Surabaya, Indonesia, Indonesia

Tonang Dwi Ardyanto, Department of Clinical Pathology, Faculty of Medicine, Sebelas Maret University/Dr. Moewardi Hospital, Surakarta, Indonesia, Indonesia

Agustin Iskandar, Faculty of Medicine, Brawijaya University, Indonesia

Tenri Esa, Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University/Dr. Wahidin Sudirohusodo, Makassar, Indonesia, Indonesia

Rachmawati Adiputri Muhiddin, Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University/Dr. Wahidin Sudirohusodo, Makassar, Indonesia, Indonesia

Liong Boy Kurniawan, Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University/Wahidin Sudirohusodo Hospital, Makassar, Indonesia, Indonesia

SUBMIT



ISSN

p-ISSN: 0854-4263

e-ISSN: 2477-4685

LINK

[Template](#)

[Statement Letter](#)

[Copyright Transfer Form](#)


INDEX



USER

Username
Password
☐ Remember me

CURRENT ISSUE



JOURNAL CONTENT

[Andaru Dahesih Dewi](#), Department of Clinical Pathology, Faculty of Medicine, Gadjah Mada University/Dr. Sardjito Hospital, Yogyakarta, Indonesia, Indonesia

[Umi Solekhah Intansari](#), Department of Clinical Pathology, Faculty of Medicine, Gajah Mada University/Dr. Sardjito Hospital, Yogyakarta, Indonesia, Indonesia

[Banundari Rachmawati](#), Department of Clinical Pathology, Faculty of Medicine, Diponegoro University/Dr. Kariadi Hospital, Semarang, Indonesia, Indonesia

[Agnes Rengga Indrati](#), Department of Clinical Pathology, Faculty of Medicine, Padjajaran University/Hasan Sadikin Hospital, Bandung, Indonesia, Indonesia

[Rikarni Rikarni](#), Department of Clinical Pathology, Faculty of Medicine, Andalas University/M.Djamil Hospital, Padang, Indonesia, Indonesia

[Hani Susianti](#), Department of Clinical Pathology, Faculty of Medicine, Brawijaya University/Saiful Anwar Hospital, Malang, Indonesia, Indonesia

[Adhi Kristianto Sugianli](#), Department of Clinical Pathology, Faculty of Medicine, Padjadjaran University/Dr. Hasan Sadikin Hospital, Bandung, Indonesia, Indonesia

[Anak Agung Wiradewi Lestari](#), Department of Clinical Pathology, Faculty of Medicine, Udayana University/Sanglah Hospital, Bali, Indonesia, Indonesia

[Yetti Hernaningsih](#), Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo Hospital, Surabaya, Indonesia, Indonesia

[Yuyun Widaningsih](#), Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University/Wahidin Sudirohusodo Hospital, Makassar, Indonesia, Indonesia

[Osman Sianipar](#), Department of Clinical Pathology, Faculty of Medicine, Gajah Mada University/Dr. Sardjito Hospital, Yogyakarta, Indonesia, Indonesia

[Rismawati Yaswir](#), Department of Clinical Pathology, Faculty of Medicine, Andalas University/M. Djamil Hospital, Padang, Indonesia, Indonesia

[Maimun Zulhaidah Arthamin](#), Department of Clinical Pathology, Faculty of Medicine, Brawijaya University/Saiful Anwar Hospital, Malang, Indonesia, Indonesia

[Aryati Aryati](#), Department of Clinical Pathology, Faculty of Medicine, Airlangga University/Dr. Soetomo Hospital, Surabaya, Indonesia, Indonesia

[Rahajuningsih Dharma](#), Department of Clinical Pathology, Faculty of Medicine, Indonesia University/Ciptomangunkusumo Hospital, Jakarta, Indonesia

[Mansyur Arif](#), Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University/Wahidin Sudirohusodo Hospital, Makassar, Indonesia

[July Kumalawati](#), Department of Clinical Pathology, Faculty of Medicine, Indonesia University/Ciptomangunkusumo Hospital, Jakarta, Indonesia, Indonesia

[Purwanto AP](#), Department of Clinical Pathology, Faculty of Medicine, Diponegoro University/Dr. Kariadi Hospital, Semarang, Indonesia, Indonesia

[Jusak Nugraha](#), Department of Clinical Pathology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo Hospital, Surabaya, Indonesia, Indonesia

[Edi Widjayanto](#), Department of Clinical Pathology, Faculty of Medicine, Brawijaya University/Saiful Anwar Hospital, Malang, Indonesia, Indonesia

[Ninik Sukartini](#), Department of Clinical Pathology, Faculty of Medicine, Indonesia University/Dr. Ciptomangunkusumo Hospital, Jakarta, Indonesia, Indonesia

[Kusworini Handono](#), Department of Clinical Pathology, Faculty of Medicine, Brawijaya University/Dr. Saiful Anwar Hospital, Malang, Indonesia, Indonesia

[Ranti Permatasari](#), Department of Clinical Pathology, Faculty of Medicine, Sumatera Utara University/Adam Malik Hospital, Medan, Indonesia, Indonesia

Search

Search Scope

All

Search

Browse

[By Issue](#)[By Author](#)[By Title](#)**Contact Us:**

Laboratorium Patologi Klinik RSUD Dr. Soetomo
Jl. Mayjend. Prof. Dr. Moestopo 6-8 Surabaya.



INDONESIAN JOURNAL OF Clinical Pathology and Medical Laboratory

Majalah Patologi Klinik Indonesia dan Laboratorium Medik

p-ISSN 0854-4263
e-ISSN 4277-4685

Published by
Indonesian Association of Clinical
Pathologist and Medical Laboratory

HOME ABOUT LOGIN REGISTER SEARCH CURRENT ARCHIVES ANNOUNCEMENTS AUTHOR GUIDELINE EDITORIAL TEAM

Home > Archives > Vol 25, No 1 (2018)

Vol 25, No 1 (2018)

DOI: <http://dx.doi.org/10.24293/ijcpml.v25i1>

Table of Contents

Articles

- PROPORTION OF ISOMORPHIC ERYTHROCYTE URINE IN DIABETIC KIDNEY DISEASE WITH FLOW CYTOMETRY METHODS

Erica Catarina, Coriejati Rita, Basti Andriyoko, Ida Parwati



Pg. 1-6

- ANALYSIS OF RET-HE IN CHRONIC KIDNEY DISEASE PATIENTS AT DR.WAHIDIN SUDIROHUSODO HOSPITAL, MAKASSAR

Febrina Rovani, Asvin Nurulita, Mansyur Arif



Pg. 7-10

- ANALYSIS OF RED BLOOD CELL DISTRIBUTION WIDTH COEFFICIENT OF VARIATION ON STROKE PATIENT

Kartika Paramita, Agus Alim Abdullah, Mansyur Arif



Pg. 11-15

- IgA ANTI-DENGUE PROFILE IN SAMPLES WITH POSITIVE DENGUE PCR OR NS1

M Thohirin Ramadhani, Aryati Aryati, M Vitanata Arfijanto



Pg. 16-20

- THE ASSOCIATION OF INSULIN RESISTANCE AND LIPID PROFILE RATIO IN METABOLIC SYNDROME

Rini Rahmayani, Adi Koesoema Aman, Santi Safril



Pg. 21-25

- CORRELATION OF FREE HEMOGLOBIN LEVEL AND PLASMA NITRIC OXIDE IN PACKED RED CELL DURING BLOOD BANK STORAGE PERIOD

Ricca Fitria, Zelly Dia Rofinda, Desywar Desywar



Pg. 26-30

- CORRELATION OF LIPID PROFILE WITH INTERLEUKIN-12 IN TYPE 2 DIABETES MELLITUS

Meri Ponda Sari, Hanifah Maani, Ellyza Nasrul, Dia Rofinda



Pg. 31-34

- PLATELET INDICES FOR PREDICTING LIVER FIBROSIS IN PATIENTS WITH



SUBMIT



ISSN

p-ISSN: 0854-4263

e-ISSN: 2477-4685

LINK

Template

Statement Letter

Copyright Transfer Form


INDEX























USER

Username
Password
☐ Remember me

CURRENT ISSUE



JOURNAL CONTENT

	CHRONIC HEPATITIS B INFECTION	
	<i>Shendy Sherly Soeliauwan, Darwati Muhadi, Mutmainnah Mutmainnah</i>	
	THE RELATIONSHIP BETWEEN THE LEVEL OF INTERLEUKIN-6 AND PROCALCITONIN IN SEVERE SEPSIS PATIENTS AT THE ADAM MALIK HOSPITAL	 PDF
	<i>Sesily C Nainggolan, Adi Koesoema Aman, Achsanudin Hanafie</i>	<i>Pg. 38-41</i>
	SPONTANEOUS PLATELET AGGREGATION IN THIRD TRIMESTER PREGNANCY AT ADAM MALIK HOSPITAL, MEDAN	 PDF
	<i>Rezqi Maulani Jusuf, Hotma Partogi Pasaribu, Herman Hariman</i>	<i>Pg. 42-46</i>
	CORRELATION BETWEEN PRESEPSIN AND SEQUENTIAL [SEPSIS-RELATED] ORGAN FAILURE ASSESSMENT (SOFA) SCORE AS AN ORGAN DYSFUNCTION MARKER IN SEPSIS	 PDF
	<i>Stevi Dwiyani, Agnes Rengga Indrati, Leni Lismayanti, Adhi Kristianto S</i>	<i>Pg. 47-52</i>
	CORRELATION OF ATHEROGENIC INDEX OF PLASMA WITH STENOSIS LEVEL OF CORONARY ARTERY IN ACUTE CORONARY SYNDROME	 PDF
	<i>Ilhamifithri Ilhamifithri, Rismawati Yaswir, Eugeny Alia, Efrida Efrida</i>	<i>Pg. 53-57</i>
	THE COMPATIBILITY OF NEUTROPHIL TO LYMPHOCYTE COUNT RATIO WITH SERUM PROCALCITONIN AS BACTERIAL INFECTION MARKERS IN SEPSIS PATIENTS	 PDF
	<i>Elvinawaty Elvinawaty, Hanifah Maani, Zelly Dia Rofinda, Husni Husni</i>	<i>Pg. 58-63</i>
	THE DIAGNOSTIC VALUE OF TROPONIN I TESTING TO CORONARY ANGIOGRAPHY WITH A POINT OF CARE TESTING INSTRUMENT IN PATIENTS WITH ACUTE MYOCARDIAL INFARCTION	 PDF
	<i>Riska Anton, Sheila Febriana, Asvin Nurulita, Uleng Bahrn</i>	<i>Pg. 64-67</i>
	COMPARISONS OF FIBRO Q INDEX AND FIB-4 IN VARIOUS STAGES OF CHRONIC B HEPATITIS	 PDF
	<i>Evy Adrianti, Liong Boy Kurniawan, Ibrahim Abdul Samad</i>	<i>Pg. 68-72</i>
	MICROORGANISM PATTERN ON NASAL CAVITY OF END STAGE RENAL DISEASE PATIENTS WITH REGULAR HEMODIALYSIS AND STAFFS IN HEMODIALYSIS INSTALLATION ADAM MALIK GENERAL HOSPITAL MEDAN	 PDF
	<i>Imelda Damayanti, Ricke Loesnihari, Syafrizal Nasution</i>	<i>Pg. 73-78</i>
	THE CORRELATION BETWEEN THE MEAN PLATELET VOLUME VALUES WITH THROMBOCYTE AGGREGATION IN NEPHROPATHY DIABETIC PATIENTS	 PDF
	<i>Agus Sunardi, Nadjwa Zamalek Dalimoenthe, Coriejati Rita, Adhi Kristianto Sugianli</i>	<i>Pg. 79-85</i>
	THE ROLE OF PLATELET CONCENTRATION TRANSFUSION ON THE CORRELATION BETWEEN PLATELET NUMBER AND MAXIMUM AMPLITUDE WITH BLEEDING VOLUME POST CARDIOPULMONARY BYPASS	 PDF
		<i>Pg. 86-90</i>

Search

Search Scope

All

Search

Browse

By Issue

By Author

By Title

Ryan Bayusantika Ristandi, Nida Suraya, Leni Lismayanti, Sylvia Rachmayati

- THE RELATIONSHIP BETWEEN NITRIC OXIDE AND GLYCEMIC CONTROL IN CONTROLLED AND UNCONTROLLED TYPE 2 DIABETES MELLITUS PATIENTS IN THE ADAM MALIK HOSPITAL MEDAN

Yessy Suziarty, Ratna Akbari Ganie, Santi Syafril



Pg. 91-94

- ANALYSIS OF RED BLOOD CELL DISTRIBUTION WIDTH VALUE TOWARDS FIBROTIC STAGE IN CHRONIC HEPATITIS B

Fatma Idris, Darwati Muhadi, Mutmainah Mutmainah



Pg. 95-98

- CORRELATION OF SERUM HIGH-DENSITY LIPOPROTEIN CHOLESTEROL AND HOMOCYSTEINE LEVEL IN PATIENT WITH ACUTE MYOCARDIAL INFARCTION

Yayie Dwina Putri, Tuty Prihandani, Lillah Lillah, Rismawati Yaswir



Pg. 99-103

- THE CORRELATION BETWEEN GALECTIN-3, CREATININE AND URIC ACID ON STAGE V CHRONIC RENAL FAILURE

Indranila K S, Guruh A I, Meita H



Pg. 104-110

Literature Review

- ROLE OF DELTA CHECK IN CLINICAL LABORATORY SERVICES

Osman Sianipar



Pg. 111-114

Case Report

- PRIMARY MYELOFIBROSIS

Muhammad Irhamsyah, Darwati Muhadi, Mansyur Arif



Pg. 115-120

- LEUKEMIC PHASE OF MALIGNANT LYMPHOMA IN CHILDREN

Sahriany S, Agus Alim Abdulah, Mansyur Arif



Pg. 121-128

Front Matter

- Cover



Pg.

- Contents



Pg.

Back Matter

- Author Guideline and Subscribes Form



Pg.

THE CORRELATION BETWEEN GALECTIN-3, CREATININE AND URIC ACID ON STAGE V CHRONIC RENAL FAILURE

by Indranila Kustarini Samsuria

Submission date: 26-Jan-2021 11:09AM (UTC+0700)

Submission ID: 1494503041

File name: 1516-2917-1-SM_organized.pdf (6.61M)

Word count: 286

Character count: 1113

THE CORRELATION BETWEEN GALECTIN-3, CREATININE AND URIC ACID ON STAGE V CHRONIC RENAL FAILURE

Indranila KS, Guruh AI, Meita H

Department of Clinical Pathology, Faculty of Medicine, Diponegoro University/Dr. Kariadi Hospital, Semarang, Indonesia. E-mail: nila_fkundip@yahoo.com

ABSTRACT

Chronic Renal Failure (CRF) is structural damage and function of the kidneys that cannot excrete toxins and waste products from the blood, characterized by the presence of protein in the urine and decreased glomerulus filtration rate. This study aimed to determine the correlation between Galectin-3 and markers of kidney function which are creatinine and uric acid. This study was being concluded on 33 CRF patients who were doing hemodialysis therapy. This study was conducted in the Dr. Kariadi Semarang Hospital and GAKI Laboratory of Diponegoro Medical Faculty from April to June 2018. The research method was analytical descriptive with cross-sectional approach. Galectin-3 was analyzed using ELISA method with an automatic analyzer, creatinine and uric acid using the colorimetric method with an automatic spectrophotometer. Statistical analysis used Shapiro-Wilk normality test and Spearman correlation test. There is a weak positive correlation test of galectin-3 with creatinine ($r = 0.381$; $p = 0.029$) and galectin-3 with uric acid ($r = 0.374$; $p = 0.048$) in CRF – HD. It is concluded galectin-3 can be used as a marker of kidney function.

Key words: Galectin-3, creatinine, uric acid, chronic renal failure, hemodialysis

INTRODUCTION

Chronic kidney failure is a world health problem with an increased incidence, prevalence, and level of morbidity. Hypertension, diabetes, smoking, the use of analgesic drugs, NSAIDs, and the use of energy drinks affect the occurrence of Chronic Renal Failure (CRF). Chronic renal failure generally develops to the terminal stage or the final stage requiring renal function replacement therapy, known as dialysis.^{1,2} Chronic renal failure lasts for at least three months and can cause persistent kidney disease.^{3,4} Long-term disease, such as hypertension and diabetes, slowly damage the kidneys and reduce kidney function over time. End-Stage Renal Disease (ESRD) requires renal replacement therapy namely Hemodialysis (HD), Continuous Ambulatory Peritoneal dialysis (CAPD) or a kidney transplant to maintain the patients survival.⁵

Galectin-3 is a protein molecule, belonging to the galactoside binding family of lectins that bind β -galactoside to glycoprotein and glycolipid cells also known as IgE binding proteins, laminin binding proteins, and Mac-2 antigens. Galectin-3 is involved in various biological processes such as inter-cell adhesion, proliferation, malignant transformation,

and metastasis. Excessive levels of galectin-3 in the body play a role in promoting the inflammatory and fibrotic response that causes progressive heart failure, liver cirrhosis, and renal failure. Research by Li *et al.* and Drechsler *et al.* stated that galectin-3 concentrations increased with progressive kidney damage.^{6,7}

Creatinine is a residual product of reshaping creatine phosphate that mostly occurs in the muscle (98%) and the rest in the heart, liver, brain, kidneys and body fluids.⁸ Creatinine levels are influenced by muscle mass, diet, physical activity, sex and age, but it is relatively stable over time. Small muscle mass cause little creatinine production.⁹ Creatinine is excreted through the urine, so the kidneys are responsible for creatinine levels in the blood.^{3,4,9} Permanent abnormal high creatinine levels indicate damage or failure of chronic kidney function and can sometimes be detected before the presence of symptoms.^{10,11}

Uric acid is the final product of adenine and guanine catabolism derived from the breakdown of purine nucleotides. Uric acid is the final product of purine metabolism which consists of carbon, nitrogen, oxygen, and hydrogen components with the molecular formula of $C_5H_4N_4O_3$. Johnson *et al.*

stated that the kidneys are responsible for excretion of two-thirds of uric acid levels every day and the other third is excreted through the gastrointestinal tract.¹¹

Chronic renal failure is known to involve an inflammatory process and fibrosis as the pathogenesis of damage to the glomerulus and renal tubules until the kidneys lose their function. Detection of renal dysfunction by urinalysis are albuminuria, hematuria and the presence of urine sediment abnormalities, and need confirmation by other examinations such as serum creatinine and uric acid parameters.¹⁰⁻¹² It is known that an increase in galectin-3 levels influences the development of CRF associated with a decrease in Glomerular Filtration Rate (GFR) with abnormally high creatinine levels as a sign of glomerular filtration function and an increase in uric acid levels or hyperuricemia as a marker of tubular reabsorption and secretion function which indicates damage or failure of kidney function.¹² creased in patients with decreased GFR, but there have not been many studies that state galectin-3 levels in various stages of CRF.⁸ This study wants to prove galactin-3 as a potential novel biomarker in patients with CRF stage V.

METHODS

This study was a descriptive analytic study with a cross-sectional approach to analyzing galectin-3 levels, creatinine levels and uric acid levels as outputs in CRF patients with hemodialysis. Data were coded, entered, cleaned and edited on software programs. Univariate analysis was performed on each variable to determine the characteristics of the sample. Bivariate analysis was conducted to find the relationship between galectin-3 levels and creatinine levels, and galectin-3 levels with uric acid levels in CRF patients with hemodialysis. The inclusion criteria were patients with CRF who received hemodialysis therapy, and willing to sign an informed consent, and had examination of creatinine, uric acid, and galectin-3. Exclusion criteria were as follows: age <18 years, psychiatric illness, and severe comorbidity affecting outcome and prognosis, including cancer, acute hepatocellular insufficiency, acute infectious disease, and exacerbation of chronic diseases. Lysed samples were also excluded. Galectin-3 was analyzed with ELISA method. Creatinine and uric acid using the colorimetric method.

Data normality test was done from each variable using Shapiro-Wilk. The results of the normality test for each data showed that uric acid levels were

normally distributed while galectin-3 and creatinine levels were distributed abnormally even after data transformation, so the Spearman correlation test was carried out. Relationships were considered significant if $p \leq 0.05$. The degree of relationship showed a very weak relationship if $r=0-0.199$, weak relationship if $r=0.2-0.399$, moderate relationship if $r=0.4-0.599$, strong relationship if $r=0.6-0.799$ and the relationship was very strong if $r=0.8-1.00$

This study has obtained ethical clearance No.172 / EC / FK-RSDK / IV / 2018 dated 17 April 2018 from the Health Research Ethics Committee of the FK UNDIP/Dr. Kariadi Hospital. All research subjects/families were asked for their approval by signing written informed consent. The patient's identity was kept confidential, and all costs related to the research were the responsibility of the researcher. All prospective research subjects were given a full explanation of the objectives, benefits and research procedures. Research subjects have the right to refuse to be included in the study.

RESULTS AND DISCUSSION

This study was conducted on CRF patients who did hemodialysis at the Dr. Kariadi Hospital Semarang. Blood samples were examined in the Semarang IDD laboratory in April 2018 - June 2018. A total of 35 samples were collected, two lysis samples were excluded from this study and 33 samples fulfilled the inclusion and exclusion criteria who were willing to sign an informed consent during this period. The study involved 33 patients with the youngest males aged 20 years and the oldest was 62 years. Based on the classification of CRF according to the degree of disease, all patients involved in this study were in the 5th-grade category with a GFR of less than 15 mL/minute. Glomerular Filtration Rate (GFR) used: Chronic Epidemiology Collaboration Equation (CKD-EPI). This study also assessed the condition of hypertension, diabetes mellitus, kidney trauma and consumption of energy drinks/herbs which are considered as risk factors for CRF. It can be seen in Table 1. Patients who had hypertension were 20 cases, diabetes mellitus were three cases, kidney stones were 8 cases and a history of energy drinks or herbal medicine consumption were two cases from all study samples.

The characteristic data of the study sample are shown in Table 2. Data with normal distribution were BMI parameters, systolic blood pressure, while the data with abnormal distribution were parameters of age, weight, height, diastolic blood pressure, galectin-3, creatinine, and uric acid levels. Statistical

Table 1. Descriptive data on risk factors for chronic renal failure



Risk factor	Total	Percentage (%)
Hypertension	20	60
Diabetes	3	9
Kidney stones	8	25
History of energy drinks/ herbal medicine consumption	2	6

tests for data normality use the Shapiro-Wilk normality test. Data were normally distributed if $p > 0.05$.

Data on galectin-3 levels and serum creatinine were abnormally distributed based on data transformation. Spearman correlation test was used to test the relationship in two variables. The relationship test results between galectin-3 levels and creatinine levels obtained were $r = 0.381$ with $p = 0.029$, meaning there was a weak positive relationship. A scatter chart plots the relationship of serum galectin-3 levels with serum creatinine levels in CRF patients with hemodialysis can be seen in Figure 1.

Data on galectin-3 levels and serum uric acid levels were abnormally distributed based on data transformation. Spearman correlation test was used to test the relationship in two variables. The test results of the relationship between galectin-3 levels and serum uric acid levels were obtained $r = 0.347$ with $p = 0.048$, meaning there was a weak positive relationship. A scatter plot graph of the relationship of galectin-3 levels with serum uric acid levels in patients with CRF with hemodialysis can be seen in Figure 2.

This study involved 33 CRF patients with hemodialysis who fulfilled the inclusion and

**Figure 1.** Scatter chart plots the relationship of serum galectin-3 levels with serum creatinine levels in CRF patients with hemodialysis**Figure 2.** Scatter chart plot of the relationship between galectin-3 levels and serum uric acid levels in patients with CRF with hemodialysis**Table 2.** Characteristics of research sample data

Variable	Mean \pm SD	Median (min – max)	Normality	
			p	Transf.
Age (years)	46.33 \pm 10.09	45 (20 – 62)	0.238	0.246
Weight (kg)	63.73 \pm 5.17	63 (52 – 73)	0.033	0.034
Height (cm)	167.65 \pm 4.37	169 (150 – 175)	0.000	0.000
BMI	22.56 \pm 1.51	22.43 (18.44 – 25.00)	0.576	
Systolic (mmHg)	150.30 \pm 19.92	150 (110 – 180)	0.084	
Diastolic (mmHg)	116.36 \pm 157.02	90 (70 – 990)	0.000	0.000
Galectin-3 (ng/dL)	12.43 \pm 1.55	12.95 (5.41 – 13.97)	0.000	0.000
Creatinine (mg/dL)	10.16 \pm 4.43	8.9 (4.4 – 20.4)	0.035	0.123
Uric acid (mg/dL)	9.55 \pm 2.96	9 (5.1 – 16.7)	0.014	0.544*

Note: * normal data distribution

BMI: Body Mass Index

exclusion criteria, all of them were male. The youngest age was 20 years, and the oldest was 62 years. Research by Dharma stated that the incidence of CRF was increasingly prevalent in young adults.¹³ This finding is due to increasing unhealthy lifestyles such as consuming fast food, stress, smoking, lack of body activity, energy drinks, and rare consuming of water.^{13,14} These poor habits are a risk factor for kidney damage.

According to the data from the Indonesian Renal Registry (IRR) in 2012, the risk factors for CRF that occur mostly in young adults were Diabetes Mellitus (DM), hypertension, smoking habits and consumption of supplemental drinks.¹⁵ Risk factors that influence the high incidence of CRF were unmodified factors such as age, sex, race, genetics, and modified factors such as hypertension, diabetes mellitus, kidney stones, and diet can influence the incidence of CRF. Hypertension was the highest risk factor in the subject of this study. It was found that 60% of patients were with hypertension. There were 25% nephrolithiasis patients and 9% of DM patients. About 6% of people had a history of consuming energy supplement and herbal drinks for a long time.^{15,16}

Uncontrolled hypertension can worsen disease progression by causing arteriolar nephrosclerosis due to the hyperfiltration process.¹⁶⁻¹⁷ Hyperfiltration injury is a general course of glomerular damage. It does not depend on the underlying cause of kidney damage. With the loss of nephrons, the remaining nephrons experience structural and functional hypertrophy characterized by increased glomerular blood flow. The driving force for glomerular filtration is the increase in living nephrons. Although the mechanism of hyperfiltration can temporarily maintain kidney function, it will subsequently lead to progressive damage to living glomeruli, due to the direct effect of increased hydrostatic pressure on the integrity of capillary walls and/or the effects of increased protein across the capillary wall. Over time, with an increased sclerosing nephrons population, surviving nephrons will experience an increased burden of excretion, which will cause a vicious cycle of hyperfiltration and increased glomerular blood flow, associated with intravascular volume overload and/or excessive renin production associated with glomerular disease.¹⁸

Nephrolithiasis is the crystallization of minerals and matrixes such as pus, blood, tissue, and tumors. The increase in solution concentration is a result of low intake and also an increase in organic matter due

to infection of the urinary tract or static urine thus making a place for stone formation. Kidney stones cause some nephrons (including glomeruli and tubules) to be damaged. Intact nephrons undergo hypertrophy, furthermore because the number of damaged nephrons increases, oliguria, and retention of residual products arise. The symptoms in the patient become more explicit, and there are typical symptoms of kidney failure if approximately kidney function has disappeared 80% - 90%. The renal function becomes low at this level, and creatinine clearance values drop to 15 mL/minute or lower.¹⁷

The condition of hyperglycemia in patients with uncontrolled DM will force the glomerulus and tubules to work harder infiltration, secretions, and reabsorption which will slowly damage the kidney nephron. Damage to the kidney nephron will be more extensive and continuous in uncontrolled DM conditions until kidney damage occurs due to diabetes is known as diabetic nephropathy. This condition generally continues towards CRF which ends with hemodialysis as a therapy for kidney replacement.¹³

According to Dharma, supplement drinks are related to the wrong eating and drinking habits.¹³ People tend to be lazy to consume nutritious food and then switch to supplements as a substitute for vitamin intake. Supplements contain synthetic vitamins resulting from chemical products that are not free from carcinogenic substances. Excessive consumption of supplemental beverages can aggravate kidney work. The supplements are containing various substances, including taurine and caffeine. Taurine is a detoxifying amino acid that gives an effect like glycine in neutralizing all types of toxins, but excess consumption of taurine in supplements in amounts that exceed the threshold of 50-100 mg makes the kidneys work harder. According to Hidayati, caffeine can narrow the arteries to the kidneys so that the blood leading to the kidneys decreases.¹⁴ As a result, the kidneys will lack food intake and oxygen. The state of kidney cells lacking oxygen and food will cause kidney cells to experience ischemia and spur the emergence of an inflammatory reaction that can end with a decrease in the ability of kidney cells to filter blood.¹⁴ The supplemental drink also contains several dangerous chemicals such as preservatives, food coloring, flavorings, and artificial sweeteners. Research conducted by Nugroho concluded that the more frequent consumption of supplements could lead to faster kidney damage.¹⁹

Galectin-3 levels in the study subjects had a mean value of 12.43 ± 1.55 with the lowest levels of 5.41 ng/mL and the highest of 13.97 ng/mL. Galectin-3 serum levels in this study were within normal limits. Galectin-3 is a profibrotic mediator that is elevated in CRF with progressive fibrosis. Normal galectin-3 serum levels can occur because CRF patients with hemodialysis have achieved stable and permanent fibrosis in the kidneys so that galectin-3 secretion as a profibrotic marker return to normal or decreased. The fibrogenesis of the kidneys that runs for more than four months has stabilized which means that the inflammatory and fibrosis processes in the kidneys have been reduced.^{20,21} The development of CRF is characterized by the development of glomerular and tubular fibrosis where galectin-3 is a profibrotic mediator in the kidneys which increases in progressive fibrosis.²²

Creatinine levels in this study had a mean value of 8.9 mg/dL with the lowest level of 4.4 mg/dL and the highest level of 20.4 mg/dL. The creatinine in this study was used as a parameter for evaluating glomerular function, wherein normal conditions almost all creatinine would be filtrated in the glomerulus and then excreted through urine. Increased serum creatinine levels indicate glomerular filtration failure, where creatinine is used to assess GFR to determine the need for dialysis as a therapy for renal replacement in CRF patients.²³

Uric acid levels in this study had a mean of 9.0 mg/dL with the lowest levels of 5.1 mg/dL and the highest levels of 16.7 mg/dL. Uric acid in this study was used as a parameter for the assessment of tubular function where most uric acid would be absorbed, partly secreted by the tubules and discharged through urine. Tubular damage causes reduced ability of absorption and secretion of uric acid and causes an increase in uric acid levels in the blood.^{24,25}

Creatinine comes from a substance called creatine, which is formed when food turns into energy through a metabolic process. About 2% of body creatine is converted to creatinine every day. Creatinine is transported through the bloodstream to the kidneys. The kidneys filter out most of the creatinine and throw it in the urine. Therefore if the kidney is disrupted, creatinine in the blood serum increases. High creatinine levels indicate kidney damage or failure, even before patients complaints arise.²¹ It is consistent with the study of Suryawan *et al.*, where serum creatinine levels are directly proportional to the degree of glomerular damage.²³

Creatinine is released from the blood by the kidneys. Damage to the glomerular causes reduced glomerular filtration which causes blood creatinine levels to increase. The wider the glomerular damage, the less creatinine is filtered out. The level of glomerular damage affects creatinine levels in the blood so that creatinine levels in the blood and urine can be used to calculate creatinine clearance, as well as GFR.²¹⁻²³ Low creatinine levels in the blood, are used as important indicators of kidney function and can be used to determine whether a person with impaired renal function needs hemodialysis or not.^{23,26-28}

Research by Okamura *et al.* states that in rats with extensive kidney damage with galectin-3 deficiency, total collagen increased, but myofibroblast (Mfb) synthesis and extracellular matrix decreased.²⁸ These results are in accordance with the study of O'Seaghdha *et al.* that found that in high serum galectin-3 levels there was a decrease in GFR in the incidence of CRF.²⁹ Tan *et al.* stated that galectin-3 levels in CRF patients with hemodialysis (GFR < 15 mL/min) higher than the control group.³⁰ It showed that galectin-3 affects the kidneys from chronic injury and fibrosis.

The test results of the relationship between galectin-3 levels and creatinine acid levels were obtained by $r = 0.381$ with $p = 0.029$, meaning there was a weak positive relationship. Rebholz stated that galectin-3 was related to urine albumin-creatinine ratio (UACR), $r = 0.08$, $P < 0.001$.³¹ The result was in accordance with Song *et al.* that was $p < 0.001$.³²

Uric acid is mainly synthesized in the liver which is catalyzed by the xanthine oxidase enzyme. Uric acid is transported to the kidneys by blood to be filtered, partially reabsorbed, and finally excreted through urine. Increased uric acid levels in urine and serum (hyperuricemia) depend on kidney function, the speed of purine metabolism, and dietary intake of foods containing purines. Purines derived from nucleic acid catabolism in the diet are converted into gout directly. The breakdown of purine nucleotides occurs in all cells, but uric acid is only produced by tissues containing xanthine oxidase especially in the liver and small intestine.¹¹

Xanthine will be converted by xanthine oxidase to uric acid. Uric acid in the kidneys will experience four stages, namely uric acid from the capillary plasma into the glomerulus and filtration in the glomerulus. Around 98-100% will be reabsorbed in the proximal tubule, then secreted into the distal lumen of the proximal tubule and reabsorbed in the distal tubule.

Uric acid will be excreted in the urine around 6% - 12% of the amount of filtration. After vein filtration in the glomerulus, almost all reabsorbed in the proximal tubule.^{12,16,30}

Mazzali *et al.* state that gout can be associated with worsening renal function, increased proteinuria, glomerulosclerosis, renal interstitial fibrosis, and preglomerular vasculopathy.³³ High uric acid can stimulate aldosterone secretion, an adrenal gland hormone that functions to increase blood pressure. High blood pressure causes the kidneys to work extra so that they can cause inflammation. Inflammation can increase galectin-3 which will trigger fibrogenesis.

The role of galectin-3 in renal fibrosis is still investigated, but galectin-3 deficiency causes a decrease consistent with MFb activation.³⁴ Galectin-3 deficiency inhibits MFb accumulation/activation and fibrosis, while thinning of specific macrophages reduces fibrosis severity where galectin-3 deficiency does not affect macrophage recruitment or proinflammatory cytokine profile of macrophages in response to interferon- γ /lipopolysaccharide.³³ Galectin-3 secretion by macrophages is very essential in activating renal fibroblasts to the profibrotic phenotype so that galectin-3 deficiency causes a decrease consistent with MFb activation.³⁴

Galectin-3 deficiency increases total collagen, but there is a decrease in myofibroblast synthesis and extracellular matrix.³⁵ Galectin-3 could protect renal tubules from chronic injury by limiting apoptosis and improving matrix and fibrosis remodeling. Serum uric acid levels cannot predict the incidence of CRF but are independently associated with the progression of CRF that has already occurred.^{16,36} CRF is associated with increased vascular risk. A study by Mok *et al.* in 2012, stated that serum uric acid levels were associated with serum creatinine levels and inversely related to GFR.³⁷

The test results of the relationship between galectin-3 levels and serum uric acid levels were obtained by $r = 0.347$ and $p = 0.048$, meaning there was a weak positive relationship. This result was following Opatowsky's research; Galectin-3 increases with increasing uric acid with $r = 0.36$; $p = 0.003$.³⁸

The research limitations were the sample size is relatively small and can affect the strength of the study to detect the results of studies that show a weak positive relationship. Thus, the correlation of galectin-3 with uric acid and creatinine, requires further research, not only to validate the robustness

of our findings but also to show that galectin-3 can provide information as an alternative biomarker and other clinical variables.

CONCLUSIONS AND SUGGESTIONS

Based on the results of research in CRF patients with hemodialysis it can be concluded that: There is a significant weak positive relationship between galectin-3 levels and serum creatinine levels as a marker of kidney function; There is a significant weak positive relationship between galectin-3 levels and serum uric acid levels as a marker of kidney function.

Based on the results of the study and the limitations of the study, it is recommended: Further assessment of galectin-3 levels, creatinine levels, and serum uric acid levels are needed as an additional parameter to assess kidney function; Grouping patients with CRF without hemodialysis needs to be considered in further research.

REFERENCES

1. Pranandari R, Supadmi W. Faktor risiko gagal ginjal kronik di unit hemodialisis RSUD Wates Kulon Progo. *Majalah Farmaseutik, UGM*, 2015; 11(2): 316-20.
2. Robinson BE. Epidemiology of chronic kidney disease and anemia. *JAMDA*, 2006; 7(9): S3-6.
3. Suwitra K. Penyakit ginjal kronik. Dalam Sudoyo AW, Setiyohadi B, Idrus A. Editors: Buku ajar ilmu penyakit dalam. Jilid II, Edisi V., Jakarta, Interna Publishing, 2009; 1035-40.
4. Prodjosudjadi W, Suhardjono, Suwitra K. Detection and prevention of chronic kidney disease in Indonesia: Initial community screening. *Nephrology*, 2009; 14(7): 669-74.
5. Prodjosudjadi W, Suhardjono A. End-stage renal disease in Indonesia: Treatment development. *Ethn Dis*. 2009; 9(1): S1-33-6.
6. Li L-C, Jun L, Jian G. Function of galectin-3 and role in fibrotic disease. *J Pharmacol Exp Ther*, 2014; 351(2): 336-43.
7. Drechsler C, Graciela D, Christoph W, Katja B, Stefan P, Andreas T, *et al.* Galectin-3, renal function, and clinical outcomes: Results from the LURIC and 4D studies. *J Am Soc Nephrol*. 2015; 26(9): 2213-21.
8. Chen A, Hou W, Zhang Y, Chen Y, He B. Prognostic value of serum galectin-3 in patients with heart failure: A meta-analysis *IJC*, 2015; 182: 168-170.
9. Banerjee A. Renal physiology. In: *Clinical physiology an examination primer*. USA, Cambridge University Press, 2005; 61.
10. Ricke L. Peran analisa urine pada penanganan penyakit ginjal dan traktus urinarius. *Majalah Kedokteran Nusantara*, 2012; 45(3): 167-76.
11. Johnson RJ, Kang D-H, Feig D, Kivlighn S, Kanellis J,

- Watanabe S, *et al.* Is there a pathogenetic role for uric acid in hypertension and cardiovascular and renal disease. *Hypertension*, 2003; 41(6): 1183–90.
12. Cain L, Shankar A, Ducatman AM, Steenland K. The relationship between serum uric acid and chronic kidney disease among Appalachian adults. *Nephrol Dial Transplant*, 2010; 25(11): 3593–9.
13. Dharma PS. Penyakit ginjal deteksi dini dan pencegahan. Yogyakarta, CV Solusi Distribusi, 2014; 1–11.
14. Hidayati T, Kushadiwijaya H, Suhardi. Hubungan antara hipertensi, merokok dan minuman suplemen energi dan kejadian penyakit ginjal kronis. *Berita Kedokteran Masyarakat*, 2008; 24(2): 90–102.
15. PERNEEFRI. 5th Report of Indonesian Renal Registry. Jakarta, Perhimpunan Nefrolog Indonesia, 2012; 1–40.
16. Johnson RJ, Segal MS, Srinivas T, Ejaz A, Mu W, Roncal C, *et al.* Essential hypertension, progressive renal disease, and uric acid: A pathogenetic link. *J Am Soc Nephrol*, 2005; 16: 1909–19. doi:10.1681/ASN.2005010063.
17. Gulati S. Chronic kidney disease. *Int Urol Nephrol*, 2010; 42(4): 1055–62.
18. Kliegman RM. Chronic kidney disease. *Nelson textbook of pediatrics*. 18th Ed., Philadelphia, Elsevier/Saunders, 2007; 535:2.
19. Nugroho SHP. Hubungan frekuensi konsumsi suplemen energi dengan stadium chronic kidney disease di ruang hemodialisis RSUD Ibnu Sina Gresik. *J Surya*, 2015; 07(01): 54–9.
20. Gopal DM, Kommineni M, Ayalon N, Koelbl C, Ayalon R, Biolo A, *et al.* Relationship of Plasma Galectin-3 to renal function in patients with heart failure: Effects of clinical status, pathophysiology of heart failure, and presence or absence of heart failure. *JAHA* 2010; 1:e000760 doi:10.1161/JAHA.112.000760.
21. Suthahar N, Meijers WC, Sillje HHW, Ho JE, Liu F-T, de Boer RA. Galectin-3 Activation and Inhibition in Heart Failure and Cardiovascular Disease: An update. *Theranostics*, 2018; 8(3): 593–609.
22. Dickerson VM, Rissi DR, Brown CA, Brown SA, Schmiedt CW. Assessment of acute kidney injury and renal fibrosis after renal ischemia protocols in cats. *The American Association for Laboratory Animal Science*, 2017; 67(1): 56–66.
23. Suryawan DGA, Arjani IAMS, Sudarmanto IG. Gambaran kadar ureum dan kreatinin serum pada pasien gagal ginjal kronis yang menjalani terapi hemodialisis di RSUD Sanjiwani Gianyar. *Meditory*, 2016; 4(2): 145–53.
24. Amin N, Mahmood R, Asad M, Zafar M, Raja A. Evaluating urea and creatinine levels in chronic renal failure pre and post dialysis: A prospective study. *JCVD*, 2014; 2(2): 1–4.
25. Gaedeke J. Renal function test. In *Laboratory and diagnostic test handbook*. New York, Academic Press Inc, 2000; 706–15.
26. Islam N, Jahan SS, Badrudduza S, Hossain Z. Evaluation of primary screening test for platelet homeostasis in patients with chronic kidney disease. *Bangladesh Journal of Medicine*, 2012; 21(2): 55–57.
27. Gbinigie O, Price CP, Heneghan C, Van den Bruel A, Plüddemann A. Creatinine point-of-care testing for detection and monitoring of chronic kidney disease: primary care diagnostic technology update. *Br J Gen Pract*, 2015; 65(640): 608–9.
28. Okamura DM, Pasichnyk K, Lopez-Guisa JM, Collins S, Hsu DK, Liu FT, *et al.* Galectin-3 preserves renal tubules and modulates extracellular matrix remodeling in progressive fibrosis. *Am J Physiol Renal Physiol*, 2011; 300(1): F245–53.
29. O'Seaghdha CM, Hwang S-J, Ho JE, Vasan RS, Levy D, Fox CS. Elevated galectin-3 precedes the development of CKD. *J Am Soc Nephrol*, 2013; 24(9): 1470–7.
30. Tan R, Liu X, Wang J, Lu P, Han Z, Tao J, *et al.* Alternations of galectin levels after renal transplantation. *Clin. Biochem*, 2014; 47(15): 83–8.
31. Rebholz CM, Selvin E, Liang M, Ballantyne CM, Hoogeveen RC, Aguilar D, *et al.* Plasma galectin-3 levels are associated with the risk of incident chronic kidney disease. *Kidney Int*, 2018; 93(1): 252–259.
32. Song G, Sun H, Han P, Ge N, Wang W, Yi T, Li S. Increased serum galectin-3 levels are associated with the development of type 2 diabetic nephropathy: A novel marker for progression. *Int J Clin Exp Med*, 2018; 11(7): 7156–64.
33. Mazzali M, Hughes J, Kim YG, Jefferson JA, Kang DH, Gordon KL, *et al.* Elevated uric acid increases blood pressure in the rat by a novel crystal-independent mechanism. *Hypertension*, 2001; 38(5): 1101–6.
34. Henderson NC, Mackinnon AC, Farnworth SL, Kipari T, Haslett C, Iredale JP, *et al.* Galectin-3 expression and secretion link macrophages to the promotion of renal fibrosis. *AJP*, 2008; 172(2): 288–98.
35. de Boer RA, Voors AA, Muntendam P, Van Gilst WH, Van Veldhuisen DJ. Galectin-3: A novel mediator of heart failure development and progression. *Eur J Heart Fail*, 2009; 11(9): 811–7.
36. Astor BC, Hallan SI, Miller IIIER, Yeung E, Coresh J. Glomerular filtration rate, albuminuria, and risk of cardiovascular and all-cause mortality in the U.S. Population. *Am J Epidemiol*, 2008; 167(10): 1226–34.
37. Mok Y, Lee JS, Kim MS, Wenying C, Young M, Sun HJ. Serum uric acid, and chronic kidney disease: The severance cohort study. *Nephrol Dial Transplant*, 2012; 27(5): 1831–5.
38. Opatowsky AR, Baraona F, Owumi J, Loukas B, Singh MN, Valente AM, *et al.* Galectin-3 is elevated and associated with adverse outcomes in patients with single-ventricle fontanocirculation. *J Am Heart Assoc*, 2016; 5(1): e002706.

THE CORRELATION BETWEEN GALECTIN-3, CREATININE AND URIC ACID ON STAGE V CHRONIC RENAL FAILURE

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7