

Optimal Control Approach For HIV-1 Infection in CD4+T Cells with RTI and PI Treatment

by R. Heru Tjahjana

Submission date: 01-Jul-2020 09:53AM (UTC+0700)

Submission ID: 1352053561

File name: 6416-17571-1-RV-email_200520.docx (78.92K)

Word count: 2965

Character count: 15078

Optimal Control Approach For HIV-1 Infection in CD4⁺T Cells RTI and PI Treatments*

R. Heru Tjahjana and Sutimin

Abstract— The purpose of this paper is to expose the optimal approach of controlling HIV infection in CD4⁺T cells with Reverse Transcriptase Inhibitors (RTI) and Protease Inhibitors (PI) treatments. The scope of the paper includes a proposed model of the dynamic system of HIV-1 infection in CD4 cells with RTI and PI as control, and a proposed objective function model that minimizes infected CD4⁺T Cells, the population of free virus and therapeutic costs. From the dynamics system model and objective function model are designed for optimal control for HIV-1 infection control. In this paper are obtained optimal control for RTI and PI therapies. The results of this paper obtained are as follows, with the optimal control approach obtained infectious control strategy that minimizes actively infected CD4⁺T Cells, the population of free virus and the cost of treatment. In other words, optimal control can be used as a good approach in determining infection control strategies that minimizes the objective function.

I. INTRODUCTION

Acquired Immunodeficiency Syndrome (AIDS) is caused by HIV infection, one of the dangerous diseases that leads to death. In Indonesia, based on the report from the Ministry of Health, until March 31st, 2016, there are more than 13,200 people died from AIDS, 198,219 people are HIV infected and 78,292 people are AIDS. From the data, 53% of the infection are transmitted among heterosexuals and it is the highest group getting HIV infection. Many researchers are interested in studying about HIV infection.

The study about HIV infection was conducted by Srivastava *et al.* They considered complete Reverse Transcription (RT) of HIV life cycle in CD4⁺ T cells and the effect of RTI drug to block the infection of CD4⁺ T cells [1]. They developed a model to capture HIV infection regarding the interaction of virus with both CD4⁺ T cells and macrophage and studied the behavior of global stability of equilibriums for the dynamical model due to the effect of antiretroviral treatment [2]. Tarfulea proposed the mathematical model taking into account the effect of combined RTI and PI drug classes to capture the dynamical behavior of HIV infection. It shows that the adherence level in taking drug determine the progress of HIV infection. The combination of RTI and PI treatments become a better treatment than that of monotherapy of RTI or PI, and drug resistance increases viral load in the body [3]. Perelson and Ribeiro used a simple model to reveal the quantitative

findings of HIV biology for acute HIV infection, drug response and the emergence of viral variants [4]. Chirove *et al.* developed a host model to capture the spread of HIV-1 infection that is transmitted from free virus interaction to target cells and contact of infected cells to healthy cells. Chirove *et al.* also took into account Langerhans target cells in the mathematical model. The infection among target cells take place due to the interaction between virus and target cells, and the interaction between infected cells and susceptible target cells. The results show that the progress of HIV infection is determined by the critical values of virus production by infected CD4⁺ T cells. Other results show that the infection cycles in the infection process are obtained [5]. Meanwhile, Sutimin *et al.* developed a model taking into account the combination effect of RTI and PI drugs to block the HIV-1 infection. The drug response of target cells depends on the characteristic of immune cells, thereby the immune system of each individual shows different drug response. In the model, they divided Langerhans and CD4⁺ T cells into sub classes namely: sub class containing the active drugs and subclass which is not acquiring drugs [6-7]. Sutimin *et al.* developed and analyzed a mathematical model incorporating the infection process through cell to cell contact of CD4⁺T cells, CTL compartment and the combination of RTI and PI treatments [8].

II. THE PROPOSED MODEL

In this section the proposed mathematical model is explained in detail. The mathematical model is developed from [1, 6-7]. In [1], Sivastava *et al.* did not incorporating cell-cell viral transmission. In [6-7], although considering cell-cell transmission but in the studies did not take into account incomplete reverse transcription in latent infection of CD4⁺ T cells. In the proposed model, we incorporate viral transmission both through cell-to-cell and the effect of incomplete reverse transcription in latent infection of CD4⁺T cells. The mathematical model introduced the interaction between the infected CD4⁺ T cells and healthy CD4⁺ T cells, virus clearance by CD4⁺ T cells, and the effect of PI drug in blocking the viral replication in infected CD4⁺ T cells. The model consists of two models. The first model is dynamical model and the second one is model of objective function. The dynamical model describes the spread of HIV-1 infection within host cells and the objective functional model describes minimization of infected CD4⁺T Cells, the population of free virus and the cost of therapy.

The proposed dynamical model can be presented as follows

*Research supported by RPI Diponegoro University.

R. Heru Tjahjana and Sutimin are with the Department of Mathematics, Diponegoro University, Semarang, 50275 Semarang (corresponding author to provide phone and fax: 024-76480922; e-mail: heru.math.undip@gmail.com).

$$\begin{aligned}
\frac{dT}{dt} &= \lambda - \mu_t - \beta_1 VT - \beta_2 VT + (\varepsilon_{RTI}\alpha + \rho)T_1 \\
\frac{dT_1}{dt} &= \beta_1 VT + \beta_2 VT - (\mu_1 + \alpha + \rho)T_1 \\
\frac{dT_i}{dt} &= (1 - \varepsilon_{RTI})\alpha T_1 - (\mu_t + \delta)T_i \\
\frac{dV}{dt} &= N\delta(1 - \varepsilon_{PI})T_i - \mu_v V - \phi VT.
\end{aligned} \tag{1}$$

The proposed objective function model that must be minimized is

$$J = \int_{t_0}^{t_f} (C_1 T_i + C_2 V + w_1 \varepsilon_{RTI}^2 + w_2 \varepsilon_{PI}^2) dt. \tag{2}$$

Consider (1), the populations of susceptible CD4⁺ T cells are symbolized by T with the constant rate μ_t . The recruitment rate of CD4⁺ T cells can be denoted by λ . The population is infected by free virus and CD4⁺ T cells with the constant rate β_1 and β_2 respectively. Sometimes, viruses enter into CD4⁺ T cell and Reverse Transcription (RT) takes place in cytoplasm. The symbol T_1 is incomplete RT CD4⁺ T cells and may return to susceptible with the constant rate ρ . Once the cells acquire active RTI drug, it returns to be healthy with the rate α due to the effectiveness of drug. The population diminish due to natural death with the constant rate μ_t . The symbol T_i represents CD4⁺ T cells that take place complete RT, thereby it is called the actively infected CD4⁺ T cells. These population diminish due to natural death and lysis at the constant rate μ_t, δ respectively, and the infection rate of the population in class T_1 reduce to $(1 - \varepsilon_{RTI})\alpha$ due to the effectiveness of RTI drug. The population of free virus V is produced by the infected CD4⁺ T cells with the average of the virus number decreases $(1 - \varepsilon_{PI})N\delta$ due to the effect of PI treatment, and diminishes due to natural death and clearance at the constant rates μ_v and ϕ respectively. The constant N and μ_t are the average number of virions produced by infected CD4⁺ T cells and the constant natural death rate of CD4⁺ T cells, respectively. In control theory, model (1), can be seen consists 4 state variables and two controls. The state variables are T, T_1, T_i and V . The control variables are ε_{RTI} and ε_{PI} .

Considering equation (2), physically, in objective function model (2) means to minimize T_i and V , also minimize ε_{RTI} and ε_{PI} . In this paper, we denoted ε_{RTI} as u_1 and ε_{PI} as u_2 respectively and follows the paper written by Garira *et al.* in [9], values of u_1 and u_2 satisfy $0 \leq u_1, u_2 \leq 1$. Logically, from objective function model (2) one wants to minimize T_i and V , and also to minimize the cost of treatment both PI treatment cost and RTI treatment cost.

The values of the parameters used in this paper can be seen in Table I.

TABLE I VALUES OF THE PARAMETERS

Parameters	Values	Descriptions	Units	Source(s)
λ	10 or 20	Production rate of CD4 ⁺ T cells	Cells/day	[10]
ϕ	(2,9)	Constant degradation rate by Langerhans cells	l/day	[11-12]
β_1	0.0001	CD4 ⁺ T cells infection rate by free virus	l/day	[13-14]
β_2	0.00015	CD4 ⁺ T cells infected rate by CD4 ⁺ T cells	l/day	[5]
δ	0.2	Constant lysis death rate of infected CD4 ⁺ T cells	l/day	[10]
α	0.26	Transmission rate of T_1 due to incomplete the RT	l/day	[1]
μ_t	0.01	Natural death rate of CD4 ⁺ T cells	l/day	[15]
μ_v	2.40	Natural death rate of free virus	l/day	[16]
μ_1	0.015	Natural death rate of CD4 ⁺ T cells due to inflammation	l/day	[17]
ρ	0.025	Constant rate from incomplete RT class to healthy CD4 ⁺ T	l/day	[1]
N	(100, 1000)	The average number of virions produced by infected CD4 ⁺ T cells	Virions/cells day	[18-20]

The parameter values used in this paper are based on estimation and data of HIV sufferers from the laboratory as reported in the paper referenced. For parameter values according to the reference values are in an interval, estimated by taking one value that is in the interval. Another reason why this paper takes parameter values such as those used in the next simulations is because recent papers on HIV-1 [6-8] also refer to those values. With the explanation regarding the development of the model at the beginning of section II and the reasons for taking the values of the parameters above, it can be said that the proposed model is a valid model.

III. MATHEMATICAL ANALYSIS

In this section, controllability analysis and the existence of optimal control is reported. In control theory, dynamical model (1) is nonlinear control system. The controllability analysis of the model (1) presented in the Theorem 3.1 as follows.

Theorem 3.1

The dynamical model (1) is locally accessible.

Proof:

The proof start from change the form of model (1) to Affine system. In Affine system form, model (1) has a drift. The next step is generate controllability matrix through Lie Bracket. After check the rank of controllability matrix and the result is full rank, so we can conclude that the model (1) is locally accessible. ■

After controllability analysis, the next analysis is existence of optimal control. The existence of optimal control can be exposed in Theorem 3.2 as follows.

Theorem 3.2

Optimal control problem with dynamical model (1) and objective function (2) is exist.

Proof:

The proof base on convexity of each summand in objective function (2). Since each summand is convex and summation of convex function is also convex then the objective functional (2) is convex. If the objective function (2) is convex then the optimal control problem is exist. ■

IV. SIMULATION RESULT

Besides values of the parameters, another important thing in the simulation is determination of initial condition $T(0)$, $T_1(0)$, $T_i(0)$ and $V(0)$. The process of control design for simulation, starts from dynamical model (1). The values of the parameters from table I are substituted in (1) and (2), so we have a system of differential equation. With the system of differential equation and objective functional (2), the Hamiltonian function is obtained. After Hamiltonian function, the Hamiltonian system is obtained. Next, the Pontryagin Maximum Principle is used to get the optimal condition for each control variables. Therefore, the equation for each control variable is obtained. Then, each control variable is substituted to Hamiltonian system and we have the ordinary differential equation system with control variable which has been substituted. The simulation result is obtained from this ordinary differential equation system with initial values for $T(0)$, $T_1(0)$, $T_i(0)$ and $V(0)$ are 850, 0, 41 and 3.76, respectively. The initial and final times, namely 0 and 70, respectively. ■

In this paper, the numerical method used to solve the ordinary differential equation system numerically is Runge-Kutta method. The computation performed using the Matlab software. The Matlab version used to make simulations is Matlab R 2008a.

The main result of simulation can be described in Figure 1 and Figure 2 as follows. The values of objective function J and control variables u_1 and u_2 are plotted in Figure 1 and Figure 2, respectively.




Figure 1. Values of objective function.

Considering objective function (2), as described in section II, the objective of this paper is to minimize the objective cost. Trend of objective function value is decreasing as can be seen in Figure 1. The plot of simulation to minimize the objective function shows this trend. Plot of objective function in Figure 1 indicates that efforts to minimize objective functions have been seen, although the functional plot of J is seen to increase at a time interval of about 350 to 450 increased compared to the value of the function at the previous interval of 150 until 345. This is possible because there are values in functional costs whose value rises at the time in that interval. Next, after 450, it can be seen that the trend of the value of J looks to continue to decreased.




Figure 2. Plot of controls versus time.

Subsequent simulations are u_1 and u_2 . As the objective of this paper, it is to find the optimal strategy through optimum control for 70 days, for each of the minimum u_1 and u_2 , the simulation result is shown in Figure 2. It is noted that the objective function (2), besides minimizing J , the main objective of this paper is to determine how the minimum possible u_1 and u_2 can minimize the objective function value. Since u_1 represents the cost of RTI and u_2 represents the cost of PI, and it is only natural that these costs are the matters that must be minimized. Plot of u_1 and u_2 versus time can be seen in Figure 2.

V. CONCLUSION

With the optimal control approach, optimal control for RTI and PI treatments can be found [11] they simultaneously minimize functional objectives. The mathematical model for the spread of HIV-1 infection is within the host cells of four nonlinear differential equations. From the dynamics model (1) and objective function (2), with the optimal control approach, it can be obtain the simulation results of minimizing objective functional value and simulation of control determination for the minimum RTI and PI. The results are reproduced in plots u_1 and u_2 throughout the initial and final times, namely 0 and 70. For the record, different parameter values can provide plot results that may not be the same as the simulation plot results, however, the values of parameters taken from these references in the source make a convincing result for the reader. From Figure 1 can be seen that the value of objective function can be minimized. From Figure 2, the minimization u_1 and u_2 was successfully shown.

21

ACKNOWLEDGMENT

This work was financially supported by the Diponegoro University under the International Scientific Publication Research 20 nt. The authors thank to the reviewers for the valuable comment to improve the quality of this paper.

REFERENCES

- [1] P. K. Srivastava, M. Banerjee and P.Chandra, "Modeling the drug therapy for HIV infection", *J. Biological System*, vol. 17(2), 2009, pp. 213–223.
- [2] M. Elaiw, "Global properties of a class of HIV models", *Non Linear Analysis Real World Applications*, vol. 11(4) ,2010, pp. 2253–2263.
- [3] N.Tarfulea, "A mathematical model for HIV treatment with time-varying antiretroviral therapy", *Int. J. Comp. Math.*, vol. 88(15) , 2011, pp. 3217–3235.
- [4] A.S. Perelson, and R.M Ribeiro, "Modeling the within-host dynamics of HIV infection" , *BMC Biology*, vol. 11(96), 2013, pp. 1–10
- [5] F. Chirove, Sutimin, E. Soewono and N. Nuraini, "Analysis of combined Langerhans and CD4+ T Cells HIV infection", *SIAM J. Appl. Math.*, vol. 74, 2014, pp. 1174 – 1193.
- [6] Sutimin, F. Chirove, E. Soewono, N. Nuraini, and L.B. Suromo, "A model incorporating combined RTIs and PIs therapy during early HIV-1 infection", *Mathematical Biosciences*, vol. 285, 2017, pp. 102–111.
- [7] Sutimin, N. Nuraini , F. Chirove, and L.B. Suromo, "Modelling Multiple Dosing with Drug Holiday in Antiretroviral Treatment on HIV-1 Infection", *J. Math. Fund. Sci.*, vol. 49(1), 2017, pp. 1–20.
- [8] Sutimin, Sunarsih, and RH. Tjahjana, "Modeling CD4+T cells and CTL response in HIV-1 infection with antiretroviral therapy", *Commun. Biomath Sci.*, vol.1, 2018, pp.100-109.
- [9] Garira, S.D. Musekwa and T. Shiri, "Optimal control of combine therapy in a single strain HIV-1 model", *Electronic Journal Of Differential Equations*, vol. 52, 2005, pp. 1–22.
- [10] R. Culshaw and S. Ruan, "A Delay-differential equation model of HIV infection of CD4+T Cell", *Math. Biosci.*, vol 165(1), 2000, pp. 27–39
- [11] A.S. Perelson and P.W. Nelson, "Mathematical analysis of HIV-1 dynamics in vivo", *Siam Rev. Science*, vol. 41(1) ,1999, pp. 3–44
- [12] A.S. Perelson, J.M. Leonard, and D.D. Ho, "HIV-1 dynamics in vivo: virion clearan rate, infected cell life-span and viral generation time", *Science*, vol. 271(5255) , 1996, pp. 1582– 1586
- [13] D. Kirschner, "Using mathematics to understand HIV Immune Dynamics", *Notices Amer. Math. Soc.*, vol. 43(2) ,1996, pp. 191–202
- [14] Z. Wang and X. Liu, "A chronic viral infection model with immune impairment", *J. Theor. Biol.*, vol. 249(3) , 2007, pp. 532–542
- [15] M.A. Stafford, L.Corey, Y. Cao.,, E.S. Daar,D.D. Ho, and A.S. Perelson, "Modeling plasma virus concentration during primary HIV infection", *J. Theor. Biol.*, vol. 203(3) , 2000, pp. 285–301
- [16] A.S.Perelson, D.E. Kirschner, and R. D. Boer, "Dynamics of HIV infection of CD4+ T cells", *Math Bioschi*, vol. 114(1) . 1993, pp. 81-125
- [17] A.T. Kamath, S. Henri, F. Battye, D.F.Tough and K. Shortman, "Developmental kinetics and lifespan of denritics cells in mouse lymmploid organs", *Blood* , vol. 100(5) , 2002, pp. 1734-1741
- [18] B.M. Adams, , H.T. Banks, M.Davidian, H.D. Kwon, H.T. Tran, S.N. Wynne, and E.S. Rosenberg, "HIV dynamics: modeling, data analysis, and optimal treatment protocols", *J. Comput. Appl. Math.* vol. 184(1), 2005, pp. 10– 49
- [19] D.S. Dimitrov, R.L. Willey, H. Sato, L.J. Chang, R. Blumenthal, and M.A. Martin, "Quantitation of human immunodeficiencyvirus type 1 infection kinetic", *J Virol.*, vol. 67(4), 1993, pp. 2182–2190
- [20] A.T. Haase, K. Henry, M. Zupancic, G. Sedgewick, R.A. Faust, H. Melroe, W. Cavert, K. Gebhard, K. Staskus, Z.Q. Zhang, P.J. Dailey, H.H Balfour, A. Erice and A.S. Perelson, "Quantitative image analysis of HIV-1 infection in lymphoid tissue", *Science*, vol. 274. (5289), 1996, pp. 985–989.

Optimal Control Approach For HIV-1 Infection in CD4+ T Cells with RTI and PI Treatments

ORIGINALITY REPORT

14%

SIMILARITY INDEX

6%

INTERNET SOURCES

11%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Submitted to School of Business and Management ITB
Student Paper 1%
- 2** journals.itb.ac.id
Internet Source 1%
- 3** Submitted to University of the Philippines Baguio
Student Paper 1%
- 4** www.hindawi.com
Internet Source 1%
- 5** Sutimin, F. Chirove, E. Soewono, N. Nuraini, L.B. Suromo. "A model incorporating combined RTIs and PIs therapy during early HIV-1 infection", Mathematical Biosciences, 2017
Publication 1%
- 6** Submitted to Higher Education Commission Pakistan
Student Paper 1%
- 7** Ding, Y.. "A fractional-order differential equation 1%

model of HIV infection of CD4⁺ T-cells",
Mathematical and Computer Modelling, 200908

Publication

-
- | | | |
|-----------------|---|-----|
| 8 | Pehuén Pereyra Gerber, Mercedes Cabrini, Carolina Jancic, Luciana Paoletti et al. "Rab27a controls HIV-1 assembly by regulating plasma membrane levels of phosphatidylinositol 4,5-bisphosphate", The Journal of Cell Biology, 2015 | 1% |
| Publication | | |
| 9 | archpedi.ama-assn.org | <1% |
| Internet Source | | |
| 10 | systemsmedicine.de | <1% |
| Internet Source | | |
| 11 | www.srcf.ucam.org | <1% |
| Internet Source | | |
| 12 | T. Dumrongpokaphan, Y. Lenbury, R. Ouncharoen, Y. Xu. "An Intracellular Delay-Differential Equation Model of the HIV Infection and Immune Control", Mathematical Modelling of Natural Phenomena, 2008 | <1% |
| Publication | | |
| 13 | Culshaw, R.V.. "A delay-differential equation model of HIV infection of CD4 ⁺ T-cells", Mathematical Biosciences, 200005 | <1% |
| Publication | | |
-

14

Yan Cheng, Fumin Zhang, Min Zhao. "A stochastic model of HIV infection incorporating combined therapy of HAART driven by Lévy jumps", *Advances in Difference Equations*, 2019

Publication

<1%

15

Elaiw, A.M.. "Global properties of a class of HIV models", *Nonlinear Analysis: Real World Applications*, 201008

Publication

<1%

16

International Journal of Numerical Methods for Heat & Fluid Flow, Volume 22, Issue 1 (2011-12-24)

Publication

<1%

17

Gesham Magombedze. "Optimal control for HIV-1 multi-drug therapy", *International Journal of Computer Mathematics*, 2010

Publication

<1%

18

fixedpointtheoryandapplications.springeropen.com

Internet Source

<1%

19

Wai-Yuan Tan, Zhengzheng Ye. "Assessing Effects of HIV Heterogeneity and Macrophage on the HIV Pathogenesis in HIV-Infected Individuals", *Journal of Theoretical Medicine*, 2000

Publication

<1%

20

Submitted to National Chung Hsing University

Student Paper

<1%

21

eprints.itb.ac.id

Internet Source

<1%

22

www.tandfonline.com

Internet Source

<1%

23

Ferrante Neri, Jari Toivanen, Giuseppe Leonardo Cascella, Yew-Soon Ong. "An Adaptive Multimeme Algorithm for Designing HIV Multidrug Therapies", IEEE/ACM Transactions on Computational Biology and Bioinformatics, 2007

Publication

<1%

24

waset.org

Internet Source

<1%

25

Wang, X.. "Global properties of a delayed HIV infection model with CTL immune response", Applied Mathematics and Computation, 20120515

Publication

<1%

26

Salvatore T. Butera. "Therapeutic targeting of human immunodeficiency virus type-1 latency: current clinical realities and future scientific possibilities", Antiviral Research, 2000

Publication

<1%

R. R. Regoes, R. Antia, D. A. Garber, G.

27

Silvestri, M. B. Feinberg, S. I. Staprans. "Roles of Target Cells and Virus-Specific Cellular Immunity in Primary Simian Immunodeficiency Virus Infection", Journal of Virology, 2004

Publication

<1%

28

Submitted to Middle East Technical University

Student Paper

<1%

29

Submitted to Universiti Teknologi Malaysia

Student Paper

<1%

30

Submitted to Medizinischen Universität Wien

Student Paper

<1%

31

Mevlüde Yakıt Ongun, İlkem Turhan. " A Numerical Comparison for a Discrete HIV Infection of CD4 T-Cell Model Derived from Nonstandard Numerical Scheme ", Journal of Applied Mathematics, 2013

Publication

<1%

32

Submitted to University of Hong Kong

Student Paper

<1%

33

Faraimunashe Chirove, Sutimin, Edy Soewono, Nuning Nuraini. "Analysis of Combined Langerhans and CD⁴⁺ T Cells HIV Infection", SIAM Journal on Applied Mathematics, 2014

Publication

<1%

34

Xia, X.. "Modelling of HIV infection: Vaccine

readiness, drug effectiveness and therapeutical failures", Journal of Process Control, 200703

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off