

Blood serum biochemistry responses and digestive enzyme activities of tilapia (*Oreochromis niloticus*) according to different dietary protein level consumption

by Subandiyono Subandiyono


Submission date: 02-May-2021 05:31PM (UTC+0700)

Submission ID: 1575733952

File name: 2020.3566-3573.pdf (285.96K)

Word count: 4516

Character count: 23543



Blood serum biochemistry responses and digestive enzyme activities of tilapia (*Oreochromis niloticus*) according to different dietary protein level consumption

Sri Hastuti, Subandiyono Subandiyono

22 Department of Aquaculture, Faculty of Fisheries and Marine Science, Diponegoro University, Semarang, Indonesia. Corresponding author: Sri Hastuti, hastuti_hastuti@yahoo.com.

15 **Abstract.** The present experiment was carried out for 40 days to determine the effects of dietary protein on haematological parameters and digestive enzyme activities of tilapia (*Oreochromis niloticus*). The four isolipid test diets contained 28, 32, 36, and 40% of protein. The diet was formulated to be tested randomly on 600 tilapia (5.0 ± 0.5 g). The experiment used three replications. The fish was reared in 60x40x50 cm aquaria; each unit was equipped with aeration and thermostat. Results indicated that haematological parameters including triacylglycerol, aspartate aminotransferase, and alkaline phosphatase decreased as the dietary protein levels increased. The lowest pepsin activity was observed in 28% of dietary protein, which indicated the lower availability of protein as a substrate for protease activities. Furthermore, the highest amylase activity was found in the fish fed on 28% of protein, as was indicated and related to higher carbohydrate level of the diet. Based on the polynomial regression analysis of pepsin and amylase activities, the dietary protein requirements for the trial tilapia were estimated at 35.73% and 35.82%, respectively, with the dietary lipid level of 10.05%.

Key Words: amylase, dietary protein requirements, pepsin, protease, tilapia haematological profile.

Introduction. Digestibility of feed ingredients is crucial in learning fish energy needs and assessment from various diets containing different protein levels. Digestive enzymes are produced in the fish body to break down nutrients in the food consumed, so that it can be absorbed easily. The role of digestive enzymes is very important for the digestive process in the body of the fish. The ability of fish to digest food is very dependent on the availability of digestive enzymes (Fitriyanti 2011). Digestive enzyme activity is an indicator to determine digestive capacity. Whenever digestive enzyme activity is high, it can be physiologically indicated by the abilities of the fish to process the diet consumed (Infante & Cahu 2007). Therefore, the activity of digestive enzymes are influenced by dietary nutrients consumed by the fish (Nurhayati et al 2014; Yamin & Palinggi 2007; Suzer et al 2007).

Feed ingredient digestibility in the fish is affected by several factors, including the feed type, the feed nutritional content (protein), and the digestion enzyme content and types in the fish digestive tract (NRC 1993; Tillman et al 1991; Hepher 1990). Fish ability in digesting feeds is very dependent on the availability of digestive enzymes. It seems that the nutrient content of feed (protein) also affects the enzyme activity digestion. Therefore, digestive enzyme activity indicators to measure dietary protein requirements are used.

Biochemistry of blood serum such as total cholesterol, triglycerides, or aminotransferase is usually used as indicators for assessing the fish nutritional and health status (Shi et al 2016). Variations in dietary nutritional profiles can cause changes in the physiological functions of fish (Kamalam et al 2017). The relationship between enzymatic and intermediate activity metabolic steps may be responsible for controlling nutrient uptake as well as specific metabolic pathways for each nutrition (Kamalam et al 2017).

Trends in increasing tilapia (*Oreochromis niloticus*) production from the aquaculture industries on 2030 reached 128 million tons (FAO 2017). As a result, the need for pellet feeds to support tilapia production will also increase. Currently, pellet feeds is commonly applied in tilapia farming. Furthermore, natural food resources in the form of trash fish and earthworms were declining (Tacon & Metian 2008) as were being used for fish feeds. Therefore, the development of pellet diets for tilapia farming at certain growth stages is important to formulate. Protein play an important role in the feed formula and it is required for better growth of *O. niloticus*. Reducing dietary protein level could affect the fish growth performances. On the other hand, the increasing of protein content above the optimal level will increase costs, as well as waste production and implicit ambient water pollution (Xu et al 2011). Yet, despite the fact that the potential production of pellet feeds for tilapia is very high, there is no accurate information available on protein requirements of the tilapia in relation to the blood serum biochemistry responses and digestive enzyme activities. This study was designed to compare the biochemistry of blood serum and digestive enzymes in different levels of dietary protein.

Material and Method

Experimental design. The experiment was arranged in a completely randomized design with total of 4 treatments and performed in triplicates. Four levels of dietary protein, namely 28, 32, 36, and 40%, each for diet 1, 2, 3, and 4, respectively, were tested on tilapia (weight 5.0 ± 0.5 g). The experimental fish were reared in $60 \times 40 \times 50$ cm aquaria for 40 days. Each aquarium was equipped with an aerator and thermostat. This research was conducted at the Laboratory of the Fish Seed Center (FSC), Siwarak, Semarang, Central Java, Indonesia.

Experimental diets. Four experimental diets were formulated to contain different levels of crude protein, namely 28, 32, 36, and 40% (Table 1).

Formulation and composition of test diets

Feed ingredients (g 100 g ⁻¹)	Diets (Crude protein, %)			
	Diet 1 (28)	Diet 2 (32)	Diet 3 (36)	Diet 4 (40)
Fish meal	40.50	40.50	40.50	40.50
Casein	0.00	4.50	9.15	11.55
Fish oil	7.15	7.15	7.15	7.15
Wheat powder	11.15	11.15	11.15	11.15
Starch	32.70	27.20	21.65	16.05
Vitamin premix*	1.00	1.00	1.00	1.00
Mineral premix**	2.00	2.00	2.00	2.00
Choline chloride	0.50	0.50	0.50	0.50
Ca(H ₂ PO ₄) ₂	1.00	1.00	1.00	1.00
Cellulose	1.00	2.00	2.90	6.10
Sodium carboxymethyl cellulose	2.00	2.00	2.00	2.00
Phagostimulant	1.00	1.00	1.00	1.00
Total	100.00	100.00	100.00	100.00
Proximate composition (%)				
Ash	7.96	8.05	7.96	7.78
Lipid	10.05	10.05	10.05	10.05
Crude protein (CP)	27.90	32.03	36.05	40.06
Gross energy(kJg ⁻¹)	15.58	15.55	15.61	15.65

*Vitamin (mg kg⁻¹ of diet): VB₁ 50 mg, VB₂ 200 mg, VB₆ 50 mg, VB₁₂ 20 mg, folic acid 15 mg, VC (30%) 325 mg, pantothenate 400 mg, inositol 1,500 mg, D-biotin (2%) 5 mg, niacin 750 mg, VA 2.5 mg, VE (50%) 100 mg, VD₃ 2 mg, VK 20 mg. **Mineral (mg kg⁻¹ of diet): Ca(H₂PO₄)₂ 1,800 mg, KH₂PO₄ 1,350 mg, NaCl 500 mg, MgSO₄·7H₂O 750 mg, NaH₂PO₄·2H₂O 650 mg, KI 1.5 mg, COSO₄·6H₂O 2.5 mg, CuSO₄·5H₂O 15 mg, ZnSO₄·7H₂O 350 mg, FeSO₄·7H₂O 1,250 mg, MnSO₄·4H₂O 80 mg, Na₂SeO₃ 6.00 mg.

Fish meal and casein was used as the main source of protein, while fish oil was used as the main lipid source. All macro-ingredients were sieved through a 160 µm-mesh size and then mixed using a blender. The micro-ingredients (vitamin premix, mineral premix, choline chloride and calcium carbonate) were mixed in a plastic container before being added to the macro-ingredients. Fish oil was homogenized before added to the mixture. The final mixture was then passed through a meat grinder to form a 3 mm-diameter pellets. Furthermore, the pellet feed was dried in an oven at temperature of 60°C. Dry pellets were packaged in plastic bags and stored at 0°C until usage.

Experimental fish and culture management. Six hundred tilapia with the body weight of 5.0 ± 0.5 g were obtained from the Fish Seed Center (Siwarak, Semarang) and used as experimental subjects. After 2 weeks-acclimatization period, the fish were randomly assigned to twelve aquaria of 60×40×50 cm with a density of 50 fish in each aquarium. Acclimatization of the experimental diet was carried out gradually, where the fish was used to be fed on commercial feed, then the test diet was proportionally added up to 20, 40, 60, 80, and finally 100%. During the 40 days trial, the fish were fed three times a day by applying the ad satiation method. During the experiment, feed intake was recorded and manure was daily removed from the aquarium. The temperature was maintained around $29.92 \pm 2.55^\circ\text{C}$, with dissolved oxygen level of $>3 \text{ mg L}^{-1}$ and ammonia nitrogen $<0.05 \text{ mg L}^{-1}$.

Sample collection and analysis. After fasting for 24 hours, all fish in the aquarium were anesthetized, counted, and weighed at the end of the experiment. The results of these measurements were used to calculate growth and survival variables. The gastrointestinal tract including the stomach and intestines of six fish per treatment ($n=24$) were weighed and then homogenized as described by Li et al (2012) for determination of digestive enzyme activities. The extract was centrifuged at 1,150 rpm at 4°C for 10 minutes, then the supernatant was collected as an enzyme source. Pepsin, amylase, and lipase activities were analyzed using a commercial kit.

Blood samples of three fish from each treatment ($n=12$) were collected via the caudal vein. Blood serum was prepared by centrifugation (1,150 rpm, 10 minutes, 4°C) for measurement of haematological parameters, including total cholesterol, triacylglycerol, alanine aminotransferase (ALT), aspartate aminotransferase (AST), and alkaline phosphatase using the colorimetric method.

Statistical analysis. Results on the measurement of the enzyme activity, blood serum hematology and growth and SR are presented as mean \pm SD. Statistical analysis was performed using one-way analysis of variance (ANOVA) followed by Duncan's multiple range test with SPSS 22.0 software. The significance test was carried out with a probability level of $p < 0.05$. Furthermore, the polynomial regression model was calculated to estimate the optimal dietary protein requirements.

Results

Biological parameters. Biological parameters, including body weight, growth performances, and survival of the experimental fish fed on four test diets (diets 1-4) for 40 days are shown in Table 2. Fish survival rates ranged from 91.69 to 95.18%, which was similar among groups ($p < 0.05$). Furthermore, dietary protein levels significantly affected feed consumption and *O. niloticus* growth ($p > 0.05$).

Table 2

Biological parameters of *Oreochromis niloticus* fed on different dietary protein levels

Biological parameters	Diets (Crude protein, %)			
	Diet 1 (28)	Diet 2 (32)	Diet 3 (36)	Diet 4 (40)
Initial weight	5.02±0.17	5.25±0.14	5.42±0.62	5.48±0.59
Final weight (g)	18.90±1.37 ^c	23.40±1.71 ^b	25.52±2.16 ^a	26.10±1.57 ^a
Feed consumption (g)	19.29±0.22 ^b	22.69±0.31 ^a	23.12±0.87 ^a	24.74±0.58 ^a
SGR (% day ⁻¹)	3.31±1.26 ^c	3.74±0.27 ^b	3.87±0.32 ^a	3.90±0.21 ^a
Survival rate (%)	91.69±3.40 ^a	95.18±3.30 ^a	94.07±4.46 ^a	94.96±5.58 ^a

Data are expressed as mean±SD of three replications. Values on the same row with different superscripts show statistically significant differences ($p < 0.05$).

35

Digestive enzyme activities. The activities of digestive enzymes, including lipase, amylase, and pepsin are presented in Table 3. Furthermore, the response of digestive enzyme activities to dietary protein levels is presented in Figure 1. Second order of polynomial regression analysis of the pepsin (Figure 1a) and amylase (Figure 1b) in relation to the dietary protein levels, followed the equations of $Y = -8.2894x^2 + 592.4x - 9889.5$ ($R^2 = 0.9558$) and $Y = 0.0084x^2 - 0.6018x + 10.972$ ($R^2 = 0.9994$), respectively. Therefore, the optimal dietary protein level calculated from the equation were 35.73 and 35.82%, with the feed lipidic level of 10.05%.

3

Table 3

Digestive enzyme activities of *Oreochromis niloticus* fed on different dietary protein levels

Enzymatic parameters	Diets (Crude protein, %)			
	Diet 1 (28)	Diet 2 (32)	Diet 3 (36)	Diet 4 (40)
Lipase (U g ⁻¹ protein)	0.99±0.24 ^a	1.01±0.05 ^a	1.30±0.12 ^a	1.12±0.12 ^a
Amylase (U mg ⁻¹ protein)	0.74±0.16 ^a	0.35±0.07 ^b	0.25±0.11 ^b	0.40±0.07 ^b
Pepsin (U mg ⁻¹ protein)	216.47±13.38 ^c	525.61±25.26 ^b	747.99±23.10 ^a	525.61±32.86 ^b

Data are expressed as mean±SD of three replications. Values on the same row with different superscripts show statistically significant differences ($p < 0.05$).




Figure 1. The response of digestive enzyme activities of *Oreochromis niloticus* fed on different dietary protein levels.

Blood serum biochemistry. The haematological parameters of *O. niloticus* are presented in Table 4. The total cholesterol value and triacylglycerol activities decreased as the levels of dietary protein increased, but there was no significant difference for the triacylglycerol content between the 3-4 feed treatments ($p>0.05$). Concerning the alanine aminotransferase (ALT) activities, only in the case of treatment with 28% protein level showed significant higher value, while in the other treatments (i.e. 32, 36, and 40% of protein) the ALT activity was equal. On the other hand, a continuous decreasing trend was observed for the aspartate aminotransferase (AST) activities. Fish that were fed on a diet containing 28% of protein showed the highest alkaline phosphatase level.

Table 4
Haematological parameters of *Oreochromis niloticus* fed on diets containing different protein levels

Haematological parameters	Diets (Crude protein, %)			
	Diet 1 (28)	Diet 2 (32)	Diet 3 (36)	Diet 4 (40)
Total cholesterol (mg dL ⁻¹)	318.60±26.68 ^a	210.73±3.48 ^b	194.10±5.03 ^c	156.98±6.96 ^d
Triacylglycerol (mM)	5.38±1.22 ^a	2.75±0.32 ^b	0.88±0.08 ^c	0.5±0.06 ^c
Alanine aminotransferase (U L ⁻¹)	3.43±1.46 ^a	2.43±0.32 ^b	2.7±0.88 ^b	2.56±0.25 ^b
Aspartate aminotransferase (UL ⁻¹)	55.2±19.25 ^a	26.0±6.33 ^b	12.6±0.75 ^b	9.76±3.28 ^c
Alkaline phosphatase (UL ⁻¹)	4.33±1.27 ^a	1.66±0.45 ^b	1.30±0.26 ^b	2.46±0.7 ^b

Data are expressed as mean±SD of three replications. Values in the same row with different superscripts are significantly different ($p<0.05$).

Discussion. The diet that is consumed by tilapia and then enters the digestive tract will undergo to the digestion processes. As long as in the fish intestine, the diet is digested and hydrolyzed by various enzymes into simpler forms or nutrients it can be more easily absorbed by the intestinal walls, and finally entering the circulatory systems (Tu et al 2015; Infante & Cahu 2007). The substrate availability is a factor that regulates enzyme activities in fish (Kuzmina 1996). High protein content in a diet is associated with low starch levels (Table 1), thereby the increasing of pepsin activities (Table 2). Similar results were found in rainbow trout (*Oncorhynchus mykiss*) (Hepher 1990). In the present study, dietary protein levels significantly affected the digestive enzyme activities of *O. niloticus* (Table 1). Furthermore, optimum dietary protein requirements for the maximum enzyme activity are among 35.73 to 35.82% (Figure 1).

Pepsin is one of the key enzymes responsible for proteolysis. In the present study, pepsin activity was relatively low in *O. niloticus* consumed diet 1. This was due to the lower availability of dietary protein as a substrate for pepsin activity; similar phenomenon was reported in hybrid catfish (*Clarias batrachus* × *Clarias gariepinus*) (Giri et al 2003). In addition, lipase activities remained constant among the four diet groups (Table 3), in accordance with the equal lipid concentrations in the related four diets (Table 1); similar results were reported for *Labeo rohita* seeds (Debnath et al 2007) as well as for goldfish (*Cyprinus carpio* var. Jian) juveniles (Liu et al 2009). Amylase activities in the fish group of diet 1 was markedly higher than that found in the other of three test diets. This phenomena is related to the highest carbohydrate and the lowest protein level of diet 1, as well as the previous studies (Giri et al 2003; Liu et al 2009; Mohanta et al 2008).

Digestible enzymes also affect the utilization of dietary protein for fish growth (Tu et al 2015). Results of the present study showed that the growth of *O. niloticus* increased as increased the dietary protein levels (Table 2) as well as on the pepsin activities (Table 3). Dietary protein is the most important component affecting the growth performances of fish (Lee & Kim 2005). Furthermore, the higher activities of digestive enzymes (i.e. pepsin in diet 3) indicates that *O. niloticus* are physiologically able to utilize the protein consumed, being in accordance with the findings of Infante & Cahu (2007). Therefore, *O. niloticus* that consumed diets containing higher protein level (Table 3) tends to undergo better growth (SGR) (Table 2). Our SGR values obtained are higher than that of the study conducted by Kushayadi et al (2020), 3.5±0.02% for *O. niloticus* (6.7±1.2 g) fed on a diet containing 30% of protein.

¹ Blood serum biochemistry of *O. niloticus* including total cholesterol, triacylglycerol, alanine aminotransferase (ALT), aspartate aminotransferase (AST), and alkaline phosphatase were influenced by dietary protein levels ($p < 0.05$) (Table 4). The increase in dietary protein content tends to decrease the blood serum biochemical values. Otherwise, the decrease in starch content (Table 1) is thought to be a factor that reduces the blood serum biochemical values. *O. niloticus* fed on diet 1 (i.e. containing 32.70% of starch) produced the highest values of total cholesterol, triacylglycerol, ALT, AST, and alkaline phosphatase (Table 4). The increasing of cholesterol content might be caused by the excessive consumption of fat (Sun et al 2015), and carbohydrates (Anand et al 2015). Diets containing high carbohydrates will increase the levels of fructose 2,6 biphosphate, so that the phosphorfructokinase-1 being more active and stimulating glycolysis reaction. The increment on glycolysis will increase the converted glucose to become fat. Furthermore, these free fatty acids are combined with glycerol to form triglycerides (Tsalissavrina et al 2006).

Triglycerides as the result of esterification process between glycerol and three fatty acid also called triacylglycerol. Triacylglycerol levels related to cholesterol levels in the blood. Therefore, the increase in triacylglycerol could cause the total cholesterol level in the blood increase, too (Tsalissavrina et al 2006). The total cholesterol value of *O. niloticus* in diet 1 was $318.60 \pm 26.68 \text{ mg dL}^{-1}$ (Table 3), which was above the normal values. The normal value of blood cholesterol levels of *O. niloticus* is $64\text{--}299 \text{ mg dL}^{-1}$ (Hrubec et al 2008).

Quantitatively, hepatic ALT and AST activities were the most important degrading enzymes in amino acid catabolism (Metón et al 1999). Results of the present study showed that the high level of AST in *O. niloticus* fed on diet 1 (Table 4) was as an indicator of malnutrition and responsible for the poor growth performance of this treatment. Additionally, AST was more active than ALT in *O. niloticus* fingerlings, and was consistent with that in tiger puffer (*Takifugu rubripes*) juveniles (Kim & Lee 2009). The activities of alkaline phosphatase could be used as markers for well-differentiated intestinal brush border membranes (Kvåle et al 2009). Higher alkaline phosphatase activities in diet 1 indicated abnormal intestinal digestive function in fish fed on this diet. Furthermore, triacylglycerol content was gradually decreasing from diet 1 to diet 4 (Table 4), which suggested the deduction of lipid transportation as dietary protein levels increased.

Conclusions. It was concluded that the optimum protein levels required by *O. niloticus* fingerlings were ranged between 35.73 to 35.82%, based on the polynomial regression analysis of pepsin and amylase activities. Dietary protein levels significantly influenced on the growth performances, blood serum biochemistry, and digestive enzyme activities of the experimental fish.

¹⁶ **Acknowledgements.** Thanks to the Dean of Faculty of Fisheries and Marine Science, Diponegoro University and the Head of the Fish Seeds Center (FSC) Siwarak, Semarang, who provided the research facilities.

References

- Anand S. S., Howkes C., de Souza R. J., Mente A., Dehghan M., Nugent R., Popkin B. M., 2015 Food consumption and its impact on cardiovascular disease: Importance of solutions focused on the globalized food system: A report from the workshop convened by the world heart federation. *Journal of the American College of Cardiology* 66(14):1590-1614.
- Debnath D., Pal A., Sahu N., Yengkokpam S., Baruah K., Choudhury D., Venkateshwarlu, G., 2007 Digestive enzymes and metabolic profile of *Labeo rohita* fingerlings fed diets with different crude protein levels. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* 146:107–114.
- Fitriliyani I., 2011 [The activity of digestive tract enzymes of tilapia (*Oreochromis niloticus*) with feed containing lamtoro leaf flour (*Leucaena leucophala*) hydrolyzed

- and without hydrolysis with extracts of sheep rumen fluid enzymes]. *Bioscientiae* 8(2):16-31. [In Indonesian].
- Giri S. S., Sahoo S. K., Sahu A. K., Meher P. K., 2003 Effect of dietary protein level on growth, survival, feed utilisation and body composition of hybrid *Clarias* catfish (*Clarias batrachus* × *Clarias gariepinus*). *Animal Feed Science and Technology* 104(1-4):169-178.
- Hepher B., 1990 Nutrition of pond fishes. Cambridge University Press, New York.
- Hrubec T. C., Cardinale J. L., Smith S. A., 2008 Hematology and plasma chemistry reference intervals for cultured tilapia (*Oreochromis* Hybrid). *Veterinary Clinical Pathology* 29:7-12.
- Infante Z. J. L., Cahu C. L., 2007 Dietary modulation of some digestive enzymes and metabolic processes in developing marine fish: Applications to diet formulation. *Aquaculture* 268(1-4):98-105.
- Kamalam B. S., Medale F., Panerat S., 2017 Utilisation of dietary carbohydrates in farmed fishes: New insights on influencing factors, biological limitations and future strategies. *Aquaculture* 467:3-27.
- Kim S. S., Lee K. J. 2009 Dietary protein requirement of juvenile tiger puffer (*Takifugu rubripes*). *Aquaculture* 287:219-222.
- Kushayadi A. G., Suprayudi M. A., Jusadi D., Fauzi I. A., 2020 Evaluation of rubber seed oil as lipid source in red tilapia (*Oreochromis* sp.) diet. *Aquaculture Research* 51:114-123.
- Kuzmina V. V., 1996 Influence of age on digestive enzymes activity in some freshwater teleosts. *Aquaculture* 148:25-37.
- Kvåle A., Harboe T., Mangor-Jensen A., Hamre K., 2009 Effects of protein hydrolysate in weaning diets for Atlantic cod (*Gadus morhua* L.) and Atlantic halibut (*Hippoglossus hippoglossus* L.). *Aquaculture Nutrition* 15:218-227.
- Lee S. M., Kim K. D., 2005 Effect of various levels of lipid exchanged with dextrin at different protein level in diet on growth and body composition of juvenile flounder *Paralichthys olivaceus*. *Aquaculture Nutrition* 11:435-442.
- Liu Y., Feng L., Jiang J., Liu Y., Zhou X. Q., 2009 Effects of dietary protein levels on the growth performance, digestive capacity and amino acid metabolism of juvenile Jian carp (*Cyprinus carpio* var. Jian). *Aquaculture Research* 40:1073-1082.
- Li X., Jiang Y., Liu W., Ge X., 2012 Protein-sparing effect of dietary lipid in practical diets for blunt snout bream (*Megalobrama amblycephala*) fingerlings: Effects on digestive and metabolic responses. *Fish Physiology Biochemistry* 38(2):529-541.
- Metón I., Mediavilla D., Caseras A., Cantó E., Fernández F., Baanante I., 1999 Effect of diet composition and ration size on key enzyme activities of glycolysis-gluconeogenesis, the pentose phosphate pathway and amino acid metabolism in liver of gilthead sea bream (*Sparus aurata*). *British Journal of Nutrition* 82:223-232.
- Mohanta K., Mohanty S., Jena J., Sahu N., 2008 Protein requirement of silver barb, *Puntius gonionotus* fingerlings. *Aquaculture Nutrition* 14:143-152.
- Nurhayati, Utomo N. B. P., Setiawati M., 2014 [Development of digestive enzymes and growth of African catfish, *Clarias gariepinus* Buchell 1822, larvae, fed silk worms and artificial feed combination]. *Jurnal Iktiologi Indonesia* 14(3):167-178. [In Indonesian].
- Tacon A. G. J., Metian M., 2008 Global overview on the use of fish meal and fish oil in industrially compounded aquafeeds: Trends and future prospects. *Aquaculture* 285:146-158.
- Tillman A. D., Hartadi H., Reksohadiprodjo S., Prawirokusumo S., Lebdoesoekojo S., 1991 [Basic animal feed science]. Gadjah Mada University Press, Yogyakarta, Indonesia. [In Indonesian].
- Tsalissavrina I., Djoko W., Dian H., 2006 Effect of high carbohydrate diets compared to high fat diets on blood triglycerides and HDL levels on *Rattusno vergicuswistar* strain. *Medical Journal Brawijaya* 22(2):86-88.
- Tu Y. Q., Xie S. Q., Han D., Yang Y. X., Jin J. Y., Liu H. Q., Zhu X. M., 2015 Growth performance, digestive enzyme, transaminase and GH-IGF-I axis gene

- responsiveness to different dietary protein levels in broodstock allogynogenetic gibel carp (*Carassius auratus gibelio*) CAS III. *Aquaculture* 446:290–297.
- Shi Z., Li X., Chowdhury M. A. K., Chen J., Leng X., 2016 Effects of protease supplementation in low fish meal pelleted and extruded diets on growth, nutrient retention and digestibility of gibel carp, *Carassius auratus gibelio*. *Aquaculture* 460:37–44.
- Sun Y., Neelakantan N., Wu Y., Lote- Oke R., Pan A., van Dam R. M., 2015 Palm oil consumption compared with vegetable oils low in saturated fat in a meta-analysis of clinical trials. *The Journal of Nutrition* 145(7):1549–1558.
- Suzer C., Kamaci H. O., Coban D., Saka S., Firat K., Ozkara B., Ozkara A., 2007 Digestive enzyme activity of the red porgy (*Pagrus pagrus*, L.) during larval development under culture conditions. *Aquaculture Research* 38(16):1778–1785.
- Xu J., Chen W., Ai C. X., 2011 Reviews on nutrient requirements and formulated feed quality evaluation system for ricefield eel (*Monopteras albus*). *Feed Industry* 32(4): 33–38.
- Yamin M., Palinggi N., 2007 [Activity of protease enzymes and condition of digestion in the gut of rapid fish (*Epinephelus fuscoguttatus*) after giving feed]. *Jurnal Riset Akuakultur* 2(2):281–288. [In Indonesian].
- *** FAO, 2017 The state of world fisheries and aquaculture 2017. FAO, Rome, Italy, 230 p.
- *** National Research Council (NRC), 1993 Nutrient requirements of warm water fishes. National Academy of Sciences, Washington DC.

Received: 02 November 2020. Accepted: 04 December 2020. Published online: 11 December 2020.

Authors:

Sri Hastuti, Department of Aquaculture, Faculty of Fisheries and Marine Science, Diponegoro University, Semarang 50275, Central Java, Indonesia, e-mail:hastuti-hastuti@yahoo.com
 S.Subandiyono, Department of Aquaculture, Faculty of Fisheries and Marine Science, Diponegoro University, Semarang 50275, Central Java, Indonesia, e-mail:s_subndiyono@yahoo.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Hastuti S., Subandiyono S., 2020 Blood serum biochemistry responses and digestive enzyme activities of tilapia (*Oreochromis niloticus*) according to different dietary protein level consumption. *AACL Bioflux* 13(6):3566–3573.

Blood serum biochemistry responses and digestive enzyme activities of tilapia (Oreochromis niloticus) according to different dietary protein level consumption

ORIGINALITY REPORT

11 %
SIMILARITY INDEX

5 %
INTERNET SOURCES

10 %
PUBLICATIONS

1 %
STUDENT PAPERS

PRIMARY SOURCES

1 ClinicalTrials.gov
Internet Source <1 %

2 Fernandez, F.. "Effects of diets with distinct protein-to-carbohydrate ratios on nutrient digestibility, growth performance, body composition and liver intermediary enzyme activities in gilthead sea bream (Sparus aurata, L.) fingerlings", Journal of Experimental Marine Biology and Ecology, 20070430
Publication <1 %

3 Daniela Ferraz Bacconi Campeche, Douglas Henrique de Holanda Andrade, Anderson Miranda de Souza, José Fernando Bibiano Melo et al. " Dietary protein:lipid ratio changes growth, digestive enzyme activity, metabolic profile and haematological parameters in hybrid surubim (×)", Aquaculture Research, 2018
Publication <1 %

4	www.karger.com Internet Source	<1 %
5	tel.archives-ouvertes.fr Internet Source	<1 %
6	ijcmas.com Internet Source	<1 %
7	Kennedy, C.J.. "Energy intake affects the biotransformation rate, scope for induction, and metabolite profile of benzo[a]pyrene in rainbow trout", Aquatic Toxicology, 20081121 Publication	<1 %
8	Yong Liu, Lin Feng, Jun Jiang, Yang Liu, Xiao-Qiu Zhou. " Effects of dietary protein levels on the growth performance, digestive capacity and amino acid metabolism of juvenile Jian carp (var. Jian) ", Aquaculture Research, 2009 Publication	<1 %
9	Gupta, S.K., A.K. Pal, N.P. Sahu, A.K. Jha, and S. Kumar. "Effects of dietary microbial levan on growth performance, RNA/DNA ratio and some physio-biochemical responses of Labeo rohita (Hamilton) juveniles", Aquaculture Nutrition, 2015. Publication	<1 %
10	S.-M. LEE. "Effect of various levels of lipid exchanged with dextrin at different protein level in diet on growth and body composition	<1 %

of juvenile flounder *Paralichthys olivaceus*",
Aquaculture Nutrition, 12/2005

Publication

11

Giri, S.S.. "Nutrient digestibility and intestinal enzyme activity of *Clarias batrachus* (Linn.) juveniles fed on dried fish and chicken viscera incorporated diets", Bioresource Technology, 200001

Publication

<1 %

12

J. Gaye-Siessegger, U. Focken, K. Becker. "Effect of dietary protein/carbohydrate ratio on activities of hepatic enzymes involved in the amino acid metabolism of Nile tilapia, *Oreochromis niloticus* (L.)", Fish Physiology and Biochemistry, 2006

Publication

<1 %

13

Jener Alexandre Sampaio Zuanon, Antonio Policarpo Souza Carneiro, Lidiane da Silva Nascimento, Donizete Aparecido da Silva et al. "Protein requirement for *Trichogaster lalius*, blue variety", Revista Brasileira de Zootecnia, 2013

Publication

<1 %

14

K.N. MOHANTA. "Optimal dietary lipid level of silver barb, *Puntius gonionotus* fingerlings in relation to growth, nutrient retention and digestibility, muscle nucleic acid content and

<1 %

digestive enzyme activity", Aquaculture Nutrition, 2/25/2008

Publication

15

Karakatsouli, Nafsika. "An Overview of the Use of Fatty Acids in Fish Farming Research during the Last Decade, with Particular Emphasis on Fish Quality", Journal of the World Aquaculture Society, 2012.

Publication

<1 %

16

T ATSUYA ODA. "Detoxifying ability of marine bacterium Flavobacterium sp. against a toxic contaminant in the reagent sodium chloride to Nannochloropsis oculata", Fisheries Science, 4/2000

Publication

<1 %

17

Yurong Zou, Qinghui Ai, Kangsen Mai. "Ontogenic development of digestive enzyme activities in juvenile soft-shelled turtle (Pelodiscus sinensis) under cultured conditions", Frontiers of Agriculture in China, 2012

Publication

<1 %

18

es.scribd.com

Internet Source

<1 %

19

eujournal.org

Internet Source

<1 %

20

projetos.unioeste.br

Internet Source

<1 %

21

www.academicjournals.org

Internet Source

<1 %

22

www.iaeme.com

Internet Source

<1 %

23

www.innspub.net

Internet Source

<1 %

24

www.pesca.sp.gov.br

Internet Source

<1 %

25

Ahlem Hamza, Kais Fdhila, Dora Zouiten, Ahmed Sleheddine Masmoudi. "Virgibacillus proomii and Bacillus mojavensis as probiotics in sea bass (*Dicentrarchus labrax*) larvae: effects on growth performance and digestive enzyme activities", Fish Physiology and Biochemistry, 2015

Publication

<1 %

26

Bamidele O. Omitoyin, Emmanuel K. Ajani, Olugbenga Orisasona, Happiness E. Bassey, Kazeem O. Kareem, Friday E. Osho. " Effect of guava (L.) aqueous extract diet on growth performance, intestinal morphology, immune response and survival of challenged with ", Aquaculture Research, 2019

Publication

<1 %

27

García-Meilán, I., B. Ordóñez-Grande, and M.A. Gallardo. "Meal timing affects protein-sparing effect by carbohydrates in sea bream: Effects on digestive and absorptive processes", *Aquaculture*, 2014.

Publication

<1 %

28

Henrique Gomes Cornélio, Fernando, Douglas Amaral da Cunha, Jennifer Silveira, Daniel Alexandre, Carlos Peres Silva, and Débora Machado Fracalossi. "Dietary Protein Requirement of Juvenile Cachara Catfish, *Pseudoplatystoma reticulatum* : PROTEIN REQUIREMENT OF CACHARA CATFISH", *Journal of the World Aquaculture Society*, 2014.

Publication

<1 %

29

Michael D. Johnston, Tanya E. Whiteside, Michelle E. Williamson, David M. Kurtz. "Toxigenic Profile of Strains Isolated from Natural Ingredient Laboratory Animal Diets ", *Cold Spring Harbor Laboratory*, 2021

Publication

<1 %

30

Pradhan, Pravata Kumar, Joykrushna Jena, Gopa Mitra, Neeraj Sood, and Enric Gisbert. "Ontogeny of the digestive enzymes in butter catfish *Ompok bimaculatus* (Bloch) larvae", *Aquaculture*, 2013.

Publication

<1 %

31 Rania F. Ismail, Norhan E. Saleh, Alaa El-Din H. Sayed. "Microplastics induced endocrine disruption, alteration in testicular tissue in tilapia (*Oreochromis niloticus*) pre-fed on *Amphora coffeaeformis*", Research Square, 2021
Publication

32 Sedigheh Babaei, Abdolmohammad Abedian Kenari, Mehdi Hedayati, Mohammad Ali Yazdani Sadati, Isidoro Metón. "Effect of diet composition on growth performance, hepatic metabolism and antioxidant activities in Siberian sturgeon (*Acipenser baerii*, Brandt, 1869) submitted to starvation and refeeding", Fish Physiology and Biochemistry, 2016
Publication

33 Xavier, B., K.K. Jain, A.K. Pal, N.P. Sahu, G. Maheswarudu, D. Gal, and S. Kumar. "Mixed feeding schedule of low and high protein in the diet of *Labeo rohita* (Hamilton) fingerlings: effect on growth performance, haemato-immunological and stress responses", Aquaculture Nutrition, 2015.
Publication

34 www.animalsciencepublications.org
Internet Source

35 www.ojs.jmolekul.com
Internet Source

36

Mengxue Xu, Tianli Sun, Xuexi Tang, Keyu Lu, Yongshun Jiang, Sai Cao, You Wang. "Title: CO2 and HCl-induced seawater acidification impair the ingestion and digestion of blue mussel *Mytilus edulis*", Chemosphere, 2020

Publication

<1 %

37

Jean - Paul Lallès. "Intestinal alkaline phosphatase in the gastrointestinal tract of fish: biology, ontogeny, and environmental and nutritional modulation", Reviews in Aquaculture, 2019

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Blood serum biochemistry responses and digestive enzyme activities of tilapia (Oreochromis niloticus) according to different dietary protein level consumption

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8