

cek eva 1

by Widyandani Sasikirana

Submission date: 26-May-2021 07:15AM (UTC+0700)

Submission ID: 1481837432

File name: ECT_OF_LONGAN_PERICARPEXTRACT_IN_PARACETAMOL-INDUCED_IN_RATS.pdf (367.31K)

Word count: 2282

Character count: 12024



Nabilah Saniyya Wahyudi, Lusiana Batubara,
Endang Sri Sunarsih, Eva Annisaa*

ANTIOXIDANT EFFECT OF LONGAN PERICARP EXTRACT IN PARACETAMOL-INDUCED IN RATS

Nabilah Saniyya Wahyudi¹, Lusiana Batubara², Endang Sri Sunarsih³, Eva Annisaa^{3*}

¹Under Graduate Student, Faculty of Medicine, Diponegoro University, Semarang, Indonesia

²Department of Biochemistry, Faculty of Medicine, Diponegoro University, Semarang, Indonesia

³Department of Pharmacology, Faculty of Medicine, Diponegoro University, Semarang, Indonesia

Jl. Prof. Soedharto SH Tembalang Semarang, Central Java, Indonesia, 50275.

Corresponding author : evaannisaa@lecturer.undip.ac.id

ABSTRACT

Background: Oxidative stress is a condition which were free radicals harm biological molecules, like lipid, protein, and DNA. Consume paracetamol can increase oxidative stress, which can cause increased level of AST and ALT serum. Longan pericarp (*Dimocarpus longan* L.) is one of natural ingredients that contain phenolic and flavonoids as an antioxidant to prevent oxidative stress. This study aims to prove the effect of longan pericarp (*Dimocarpus longan* L.) on AST and ALT serum levels of Wistar rats induced paracetamol. **Methods:** Experimental research with post-test only control group design. Thirty males wistar aged 2-3 months weighing 140-250 grams randomly divided into 5 groups. Group K1 was normal control group. Group K2 was induced by paracetamol 2000 mg/kgBW. Group P1, P2, P3 were induced by paracetamol 2000 mg/kgBW administered by longan shell extract (*Dimocarpus longan* L.) in multilevel dose 5 mg/200 gBW, 10 mg/200 gBW, and 15 mg/200 gBW. Blood samples in each group were collected to measure AST and ALT serum levels of wistar rats. **Results:** There were significant differences groups of AST serum. There were significant difference groups of K1-K2 ($p=0,000$), K2-P2 ($p=0,041$), and K2-P3 ($p=0,010$). There were significant differences groups of ALT serum. There were significant differences groups of K1-K2 ($p=0,001$), K2-P2 ($p=0,032$), and K2-P3 ($p=0,006$). **Conclusions:** The longan pericarp extract can lower the AST and ALT serum levels of Wistar rats induced by paracetamol.

Keywords: ALT; AST; longan pericarp extract; paracetamol.

INTRODUCTION

Oxidative stress is a condition that free radicals attack biological molecules, like lipid, protein, and DNA. Oxidative stress occur when there was free radicals and antioxidant imbalance in the body.^{1,2} Drugs consumption, like paracetamol can cause oxidative stress increased and can cause hepatocyte cell membrane damage. So, the hepatocyte cell membrane permeability increased, and then enzymes in the liver, including AST and ALT will be released to the blood supply. So that the levels of AST and ALT serum rise.^{3,4}

Oxidative stress can reduced by consuming antioxidant. Antioxidant molecules can protect cell from damage, because antioxidant netralize free radicals by donating hydrogen to atom that does not

pairing to another electron. There is high antioxidant levels on plants, especially on pericarp. Antioxidant activity related to phenolic and flavonoid agent on longan pericarp.⁵

Many natural resources used in society to improve health, protect the body from diseases. Longan pericarp is one of natural resource that can be used to increase antioxidant levels in the body to reduce free radicals levels and prevent liver damage caused by stress oxidative. However, longan pericarp is underused and often become waste.⁶

Therefore, the writer wants to know the effect of longan pericarp extract (*Dimocarpus longan* L.) to AST and ALT serum levels on wistar rats that induced by paracetamol.



Nabilah Saniyya Wahyudi, Lusiana Batubara,
Endang Sri Sunarsih, Eva Annisaa*

METHOD

Experimental animals and study design

This research is experimental research with Post Test Control Group Design approach. Thirty male Wistar rats (*Rattus norvegicus*) aged minimal 8 weeks, weighing 140-250 grams, healthy and active randomly divided into 5 groups. Each group consists of 6 rats. Twenty four rats carried out the induction with paracetamol. The rats had not received treatment before. Exclusion criteria : the rats have some anatomical abnormalities. Dropout criteria : rats that died during the experiment. Rats were divided into 5 groups randomly, where each group consisted of 6 rats to be treated as follows;

- K1: Normal control group, were administered standard diet for 14 days.
- K2: Negative control group, were administered standard diet and induced by paracetamol 2000 mg/kgBW on day-14.
- P1: First treatment group, were administered longan pericarp extract with 5 mg/200 grBW dose for 14 days and induced by paracetamol 2000 mg/kgBW on day-14.
- P2: Second treatment group, were administered longan pericarp extract with 10 mg/200 grBW dose for 14 days and induced by paracetamol 2000 mg/kgBW on day-14.
- P3: Third treatment group, were administered longan pericarp extract with 15 mg/200 grBW dose for 14 days and induced by paracetamol 2000 mg/kgBW on day-14.

Longan Pericarp Extract (*Dimocarpus longan* L.) Preparation

Longan pericarp (*Dimocarpus longan* L.) washed and separated from the pericarp. Then, the longan pericarp thin cut and dried in chamber temperature. Longan pericarp extracted by soxhlet method. We used alcohol 95% for this method. Then, put the sample that wrapped in the filter paper to soxhlet tube. Next, put the sample in the porcelain cup and evaporate in water bath with 80°C until form some paste.

Statistical Analysis

One-Way Analysis of Variance (ANOVA) and Kruskal-Wallis were used for statistical analysis. Statistical analyses were done using SPSS version 22.0 for windows.

Ethical Clearance

This research has been accepted and given an ethical clearance from Komisi Etik Penelitian Kesehatan (KEPK) Faculty of Medicine, Diponegoro University with ethical clearance number is 64/EC/H/KEPK/FK-UNDIP/V/2019 on May, 21st 2019.

RESULT

Longan Pericarp Extraction

Yield is a ratio between quantity of extract resulting from extraction process. Yield obtained by compare the extract weight with dried and mashed up material multiplied 100%.⁷ Yield of longan pericarp extraction can be seen in Table 1

Table 1. Total Yield of Longan Pericarp Extraction


Solvent	Material	Dried weight (gram)	Extract weight (gram)	Yield (%)
Etanol 96%	Longan pericarp	215,3386	16,43	7,62



Nabilah Saniyya Wahyudi, Lusiana Batubara,
Endang Sri Sunarsih, Eva Annisaa'

AST Serum Levels

Data analysis of AST serum levels can be seen in Graph 1.



Graph 1. AST Serum Levels Mean

One Way ANOVA parametric test shows significant difference between groups ($p=0,009$). Then further

data analysis will be continued using the Post Hoc Test to assess meaningful differences between the 2 groups.

Table 2. Post Hoc Result for AST Serum Levels

Groups (n=6)	AST Serum Levels (U/L)	K1	K2	P1	P2
K1	113,91 \pm 15,51	-	-	-	-
K2	155,60 \pm 15,81	0,000*	-	-	-
P1	134,74 \pm 22,74	0,054	0,054	-	-
P2	133,37 \pm 19,55	0,071	0,041*	0,895	-
P3	126,78 \pm 14,28	0,223	0,010*	0,447	0,529

Information: $p<0.05$; Significant

Based on the Post Hoc Test, the results showed that there were significant differences between the K2 group with K1 ($p<0,001$), K2 with P2 ($p = 0.041$), K2 with P3 ($p = 0,010$) which meant there were significant differences in the group induced paracetamol but not given longan pericarp


extract (K2) with the group induced paracetamol and given longan pericarp extract with 10 mg/200 grBW dose (P2) and 15 mg/200 grBW dose (P3)

ALT Serum Levels

Data analysis of ALT serum levels can be seen in Graph 2.



Nabilah Saniyya Wahyudi, Lusiana Batubara,
Endang Sri Sunarsih, Eva Annisaa*



Graph 2. ALT Serum Levels Mean

One Way ANOVA parametric test shows significant difference between groups ($p=0,012$). Then further data analysis will be

continued using the Post Hoc Test to assess meaningful differences between the 2 groups.

Table 3. Post Hoc Result for ALT Serum Levels

Kelompok (n=6)	Kadar ALT Serum Tikus (U/L)	K1	K2	P1	P2
K1	63,92 \pm 12,65	-	-	-	-
K2	86,61 \pm 8,40	0,001*	-	-	-
P1	74,45 \pm 7,63	0,090	0,053	-	-
P2	73,01 \pm 12,75	0,141	0,032*	0,812	-
P3	68,73 \pm 9,15	0,429	0,006*	0,348	0,480

Information: $p<0.05$; Significant

Based on the Post Hoc Test, the results showed that there were significant differences between the K2 group with K1 ($p=0,001$), K2 with P2 ($p = 0,032$), K2 with P3 ($p = 0,006$) which meant there were significant differences in the group induced paracetamol but not given longan pericarp extract (K2) with the group induced paracetamol and given longan pericarp extract with 10 mg/200 grBW dose (P2) and 15 mg/200 grBW dose (P3).

DISCUSSION

This study aimed to determine the AST and ALT serum levels of paracetamol-induced wistar rats in the given of longan pericarp extract. The results showed that the paracetamol induced group had higher AST and ALT serum levels than the normal control group. This was in accordance with previous studies where toxic dose of paracetamol 2000mg / kgBW in wistar rats can AST and ALT serum levels.⁸ K1 were



Nabilah Saniyya Wahyudi, Lusiana Batubara,
Endang Sri Sunarsih, Eva Annisaa*

induced by paracetamol continued to increase of AST and ALT serum levels, it can be concluded that paracetamol can work more effectively to increase AST and ALT serum levels.

Longan pericarp extract contains phenolic and flavonoid which act as an antioxidant. Phenolic compounds can donate hydrogen atom to free radicals that does not have electron pair, so the new molecules will be more stable. Phenolic compounds as an antioxidant can also interacted to protein in the body, so can prevent the enzyme that plays a role in free radicals form, like *lipooxygenase*, *cyclooxygenase*, and *xanthine oxidase*.⁹ This result match to the previous research that phenolic and flavonoid compounds can decrease AST and ALT serum levels in wistar rats.^{10,11}

The results showed that longan pericarp extract had a therapeutic effect in reducing AST and ALT serum levels. Negative control groups who were not given therapy experienced an increase in AST and ALT serum levels, while the treatment group was given a therapy of longan pericarp extract with 5 mg / 200grBW, 10mg /200grBW, 15 mg/200grBW doses can reduce AST and ALT serum levels in paracetamol-induced wistar rats. Based on data analysis all the doses above can significantly reduce uric acid levels but did not find significant differences between groups P1, P2, and P3. However, when we observed from the means of AST and ALT serum level, the AST and ALT levels serum difference between the groups, and from the average p value obtained, the P3 group given a longan pericarp extract with a dose of 15 mg /200grBW was the best result.

CONCLUSIONS

From this study we can conclude that longan pericarp extract can reduce AST and ALT serum levels in paracetamol-induced wistar rats.

REFERENCES

1. Betteridge JD. What Is Oxidative Stress? *Metab - Clin Exp*. 2000;49(2):3-8.
2. Kehrer JP, Klotz LO. Free radicals and related reactive species as mediators of tissue injury and disease: Implications for Health. Vol. 45, *Critical Reviews in Toxicology*. 2015. 765-798 p.
3. Kusuma IW. Media Medika Muda. Pengaruh Parasetamol Dosis Analgesik Terhadap Kadar Serum Glutamat Piruvat Transaminase Tikus Wistar Jantan [Internet]. 2013. Available from: <https://ejournal3.undip.ac.id/index.php/medico/article/view/4937/4763>
4. Moyer AM, Fridley BL, Jenkins GD, Batzler AJ, Pelleymounter LL, Kalari KR, et al. Acetaminophen-NAPQI hepatotoxicity: A cell line model system genome-wide association study. *Toxicol Sci*. 2011;120(1):33-41.
5. Muhtadi, Hidayati AL, Suhendi A, Sudjono TA, Haryoto. Kulit Buah Asli Indonesia Dengan Metode Ftc K-50 K-51. *Simp Nas RAPI XIII - 2014 FT UMS*. 2014;50-8.
6. Yang X, Yan F, Huang S, Fu C. Antioxidant activities of fractions from longan pericarps. 2014;2014(006290):1-5.
7. Fitokimia S, Sansevieria D, Dewatisari WF, Rumiyaniti L, Rakhmawati I. Rendemen dan Skrining Fitokimia pada Ekstrak Daun Sansevieria sp. Rendemen and Phytochemical Screening using Leaf extract of Sansevieria Sp. *Res Gate*. 2018;17(Jurnal Penelitian Pertanian Terapan):197-202.
8. Wang X, Wu Q, Liu A, Anadón A, Rodríguez JL, Martínez-Larrañaga MR, et al. Paracetamol: overdose-induced oxidative stress toxicity, metabolism, and protective effects of various compounds in vivo and in vitro. *Drug*



Nabilah Saniyya Wahyudi, Lusiana Batubara,
Endang Sri Sunarsih, Eva Annisaa'

-
- Metab Rev [Internet]. 2017;49(4):395–437. Available from: <http://dx.doi.org/10.1080/03602532.2017.1354014>
9. Pereira DM, Valentão P, Pereira JA, Andrade PB. Phenolics: From Chemistry to Biology. *Molecules*. 2009;2202–11.
 10. Rahayu L, Yantih N, Supomo Y. Analisis SGPT dan SGOT pada Tikus yang Diinduksi Isoniazid untuk Penentuan Dosis dan Karakteristik Hepatoprotektif Air Buah Nanas (*Ananas comosus* L . Merr) Mentah (Analysis of SGPT and SGOT in Isoniazid-induced Rats for Dose Determination and Hepatoprotective Characteristics of Unripe Pineapple Fruit Water (*Ananas Comosus* L . Merr). 2018;16(1):100–6.
 11. Hardiningtyas SD, Purwaningsih S, Handharyani E, Agatis J, Hewan FK. Antioxidant Activity and Hepatoprotective Effect of Green Mangrove Leaves. *JPHPI*. 2014;17:80–91.

ORIGINALITY REPORT

26%

SIMILARITY INDEX

21%

INTERNET SOURCES

18%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

1

ejournal3.undip.ac.id

Internet Source

7%

2

jifi.farmasi.univpancasila.ac.id

Internet Source

2%

3

Submitted to Universitas Diponegoro

Student Paper

2%

4

Xueyi Chen, Linglu Qi, Xiaorong Fan, Hehua Tao et al. "Prenatal hypoxia affected endothelium-dependent vasodilation in mesenteric arteries of aged offspring via increased oxidative stress", Hypertension Research, 2019

Publication

1%

5

www.apjtb.com

Internet Source

1%

6

Ratih Budinastiti, Henna Rya Sunoko, Nyoman Suci Widiastiti. "The Effect of Cloud Ear Fungus (*Auricularia polytricha*) on Serum Total Cholesterol, LDL And HDL Levels on Wistar

1%

Rats Induced by Reused Cooking Oil", E3S Web of Conferences, 2018

Publication

7	www.ncbi.nlm.nih.gov Internet Source	1 %
8	digilib.uns.ac.id Internet Source	1 %
9	repository.lppm.unila.ac.id Internet Source	1 %
10	"Poster Presentations", FEBS Journal, 6/2006 Publication	1 %
11	Nirmala Chongtham. "Nutritional Properties of Bamboo Shoots: Potential and Prospects for Utilization as a Health Food", Comprehensive Reviews in Food Science and Food Safety, 05/2011 Publication	1 %
12	Yibin LI, Pufu LAI, Junchen CHEN, Hengsheng SHEN, Baosha TANG, Li WU, Minjie WENG. "Extraction optimization of polyphenols, antioxidant and xanthine oxidase inhibitory activities from Prunus salicina Lindl.", Food Science and Technology, 2016 Publication	1 %
13	sipora.polije.ac.id Internet Source	1 %

14	Submitted to University of Surrey Student Paper	1 %
15	www.spandidos-publications.com Internet Source	1 %
16	cwww.intechopen.com Internet Source	1 %
17	Z. Chen, Y.-Y. Xu, R. Wu, Y.-X. Han, Y. Yu, J.-F. Ge, F.-H. Chen. "Impaired learning and memory in rats induced by a high-fat diet: Involvement with the imbalance of nesfatin-1 abundance and copine 6 expression", Journal of Neuroendocrinology, 2017 Publication	1 %
18	www.jcdr.net Internet Source	1 %
19	www.researchgate.net Internet Source	<1 %
20	opendentistryjournal.com Internet Source	<1 %
21	www.irdrjournal.com Internet Source	<1 %
22	Antonio Giorgio, Luciano Tarantino, Giorgio de Stefano, Nunzia Farella et al. "Interstitial laser photocoagulation under ultrasound guidance of liver tumors: results in 104 treated	<1 %

patients", European Journal of Ultrasound,
2000

Publication

23

Cornelius Coli, Gadis Meinar Sari, Purwo Sri
Rejeki. "Acute Moderate Intensity Exercise
Decreases Oxygen Saturation In Obese
Women", STRADA Jurnal Ilmiah Kesehatan,
2020

Publication

<1 %

24

Maria Celeste Martino. "Intravenous infection
of virulent shigellae causes fulminant hepatitis
in mice", Cellular Microbiology, 1/2005

Publication

<1 %

25

docobook.com

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off