

The association between adiponectin level and non alcoholic fatty liver disease (NAFLD) in obese adolescents

by Mexitalia Maria

Submission date: 12-Aug-2021 03:44PM (UTC+0700)

Submission ID: 1630561019

File name: on_alcoholic_fatty_liver_disease_NAFLD_in_obese_adolescents.docx (54.02K)

Word count: 2676

Character count: 14705



Original Article

The association between adiponectin level and non alcoholic fatty liver disease (NAFLD) in obese adolescents

Maria Mexitalia¹, Suci Romadhona^{1,2}

¹Bagian / KSM Ilmu Kesehatan Anak Fakultas Kedokteran Universitas Diponegoro / RSUP Dr. Kariadi Semarang

²Direktorat Mutu Akreditasi Rumah Sakit Kementerian Kesehatan Republik Indonesia Jakarta

p-ISSN: 2301-4369 e-ISSN: 2685-7898
<https://doi.org/10.36408/mhjcm.v7i2.501>

Diajukan: 27 Agustus 2019
Diterima: 27 Mei 2020

Afiliasi Penulis:
Bagian / KSM Ilmu Kesehatan Anak
Fakultas Kedokteran Universitas Diponegoro
/ RSUP Dr. Kariadi Semarang

Korespondensi Penulis:
Maria Mexitalia
Jl. Dr. Sutomo No. 16, Semarang,
Jawa Tengah 50244, Indonesia

E-mail:
dr.mexitalia@gmail.com

Abstrak

Latar belakang : Non alcoholic fatty liver disease (NAFLD) has been associated with cytokines and inflammatory mediators. Adiponectin has insulin sensitizing effects and has correlation with severity of NAFLD disease. However, the study about the relationship between adiponectin level and NAFLD is lacking. The objective of the study was to determine the association between adiponectin level and NAFLD in obese adolescents through the role of insulin resistance.

Methods : This was a cross-sectional study, that was done in August 2007. The inclusion criteria were obese adolescents aged 11–14 years, and normal weight adolescent as control group. Adiponectin was assessed by using ELISA, insulin resistance was obtained by Homeostasis Model Assessment Insulin Resistance (HOMA-IR). NAFLD was confirmed by abdominal ultrasound, which represented by fatty liver imaging. The comparison of adiponectin level and HOMA-IR among 3 groups were analyzed by Kruskal Wallis test, meanwhile the correlation between adiponectin level and some variables were analyzed by Spearman correlation.

Results : There were 73 subjects, consisted of 37 obese and 36 non obese. Among obese subjects, 54.1% got NAFLD. All of our obese subjects were insulin resistance, the HOMA-IR level of obese non NAFLD was 6.1 and obese with NAFLD was 6.8. The adiponectin levels in normal children was (5.1g / ml), obese non NAFLD (4.1g / ml) and obese with NAFLD (4.0g / ml) ($p < 0.001$). There were no association between adiponectin level and other variables.

Conclusions : There were significant differences of adiponectin levels and insulin resistance measured by HOMA-IR between normal and obese subjects, but no significant differences between the obese groups with or without NAFLD.

Keywords: NAFLD, adiponectin, HOMA-IR, obesity, adolescent

INTRODUCTION

Obesity has emerged as a global epidemic in children with a spectrum of psychosocial and medical consequences manifesting a lifespan. The serious health problem of obese children includes metabolic syndrome, insulin resistance and non alcoholic fatty liver disease (NAFLD).^{1,2} Our previous study in Semarang Indonesia showed that the prevalence of metabolic syndrome among obese children was 31.6%.³

The amount of NAFLD increases with increasing cases of obesity, and the prevalence is around one-third of the obese children.⁴ NAFLD characterized by the accumulation of fat in the liver without a history of alcoholism or known liver pathology.^{1,5} There was increasing evidence that NAFLD often represents a component of the metabolic syndrome characterized by obesity, hyperinsulinemia, insulin resistance, diabetes, hypertriglyceridemia, and hypertension.^{2,6,7} It was reported that 53% of obese children have NAFLD.^{8,9} NAFLD was confirmed by abdominal ultrasonography which was represented by bright liver / fatty liver imaging. Abdominal ultrasonography has 94% sensitivity and 84% specificity to detect NAFLD that described increase echogenicity as a bright liver.^{10,11}

Adiponectin also known as Acrp30 is a 30k Dalton protein that is almost exclusively secreted from white adipose tissue. It is a potent modulator of glucose and lipid metabolism and an indicator of metabolic disorders. The gene expression and plasma levels of this hormone correlated with the insulin sensitive state. Dysregulation in the synthesis and/or secretion from the adipose tissue may play a role in the pathogenesis of insulin resistance in obesity. Adiponectin level was lower in the obese children and had associated with insulin resistance, which was obtained by homeostasis model assessment (HOMA-IR) equation.¹²⁻¹⁴ Low adiponectin level was an indicator of NAFLD in adults, but it seems do not correlate with the degree of hepatic steatosis in children.¹⁵

Therefore, the purpose of the study was to determine the association between adiponectin level and NAFLD in obese adolescent and the correlation between adiponectin with insulin resistance measured by HOMA-IR.

METHODS

The design was a cross-sectional study. The inclusion criteria were obese adolescents aged 11-14 years grade VIII from private high school in Semarang, and non obese adolescents for control. The study was conducted at August 2007. Anthropometric measurements, weight (to the nearest 100 g) and fat percentage were measured by Bioelectrical impedance analysis (BIA) Tanita BC 545 and height (to the nearest 0,1 cm) were measured by microtoise. All obese adolescents had a body mass index (BMI) above 95th percentile CDC 2000 chart, whereas

nonobese adolescents had a BMI between the 50th and 75th percentiles CDC 2000 growth chart. Venous blood sample was taken after 8 hours of fasting for measuring insulin, fasting blood glucose and adiponectin. Adiponectin level (g/ml) was assessed by using ELISA ELX 800 Universal Microplate Reader. Fasting blood glucose level (mg/dl) and insulin level (IU/L) was assessed using Spectrofotometer COBAS MIRA. Homeostasis model assessment (HOMA-IR) (mg/dl) was calculated from fasting glucose level multiplied by fasting insulin level divided by 22.5. Insulin resistance was considered if HOMA-IR > 3.16 mg/dl.¹⁶ Abdominal ultrasound only conducted for obese subjects, and was assessed by one radiologist medical doctor in Dr. Kariadi Hospital. The same ultrasound printout was reassessed by the same radiologist 2 month after ultrasound examination (Kappa 0.8).

The comparison of adiponectin level and HOMA-IR among 3 groups were assessed by Kruskal Wallis test, meanwhile the correlation between adiponectin level and some variables including HOMA-IR was analyzed by spearman correlation. P value were considered significant if $p < 0.05$ with 95% confidence interval. Data analysis were done using *Statistics Program for Social Science* v.15.0. (SPSS Inc, USA). Informed consents were signed by subject's parents, and the study had been approved by the Faculty of Medicine Diponegoro University / Dr. Kariadi Hospital Ethic Committee (IRB) with number 45/EC/FK/RSDK/2007.

RESULTS

There were 73 subjects, consisted of 37 obese and 36 non obese adolescents as control. Among obese subjects, 26 (70.3%) boys and 11 (29.%) girls; and 20 (54.1%) got NAFLD and 17 (45.9%) without NAFLD. Table 1 showed the characteristic of the obese and non obese (normal) subjects and table 2 showed the correlation between adiponectin and some variables.

Based on the HOMA-IR >3.16 was insulin resistance, our data showed that all obese subjects was insulin resistance.

DISCUSSION

Our study revealed that 54.1% of obese children had NAFLD. This was lower compared to New Delhi, where the prevalence of NAFLD among adolescent obese is 62.5%.¹⁷ Zou in China found that the prevalence of NAFLD was 55.75% of children with obesity, while Aksoy in Turkey revealed that 21.6% patients with NAFLD had already got non alcoholic steatohepatitis (NASH).^{5,18} NAFLD often occurs in obese children. The risk of NAFLD was risen with increasing of waist circumference, elevated total cholesterol, triglycerides,

TABEL 1
The comparison among normal, obese non NAFLD, and NAFLD subjects on various variables

Variables	Normal	Obese Non NAFLD	Obese NAFLD	p
n	36	17	19	
BMI (kg/m ²)	18.4 ± 1.13	27.97 ± 2.23	29.9 ± 2.79	0.025§ *
Fasting Insulin (IU/L)	13.9 ± 9.60	26.9 ± 17.90	27.4 ± 19.84	0.949§
Fasting glucose (mg/dl)	91.2 ± 7.97	90.9 ± 10.17	98.8 ± 10.03	0.023§ *
HOMA-IR	3.1 ± 2.25	6.1 ± 4.22	6.8 ± 5.05	0.716§
Adiponectin (µg/ml)	5.1 ± 1.63	4.1 ± 0.96	4.0 ± 0.98	0.739§

§ Mann-Whitney (the difference between Obese Non NAFLD and Obese NAFLD subjects)

¶ Kruskal Wallis (the difference among normal, obese non NAFLD and obese NAFLD subjects)

*significant p < 0.05

TABEL 2
Correlation between adiponectin level and some variables

Variables	Adiponectin level (µg/ml)	
	r	p
Fasting blood glucose level (mg/dl)	-0.56	0.634
Insulin level (IU/L)	-2.02	0.083
HOMA-IR (mg/dl)	-2.02	0.082

*Spearman correlation test n=75

blood glucose, fasting insulin and the HOMA-IR index. According to Propokowiz's study, the best independent predictive factor for diagnosing NAFLD in obese children was fasting insulin.⁴ Whereas Keskin *et al.* showed that the cut off value of HOMA-IR for diagnosing insulin resistance was 3.16.¹⁶

Insulin resistance is a key mechanism in the pathogenesis of fatty liver disease. It causes fat accumulation in hepatocytes through two main mechanisms namely lipolysis and hyperinsulinemia. Insulin resistance is a pathophysiological abnormality that underlies the development of significant disease in the clinic, including the deterioration of insulin potential in regulation of energy metabolism, control of transmembrane ion transport for protein synthesis, control of gene transcripts, and cell proliferation. Practically insulin resistance is used to explain the deterioration of insulin potential both endogenous and exogenous to increase the uptake and use of glucose by the body's cells. While the main role of insulin itself in hepatocytes is to control liver glucose production by suppressing glycogenesis. Various component, especially those released by adipocytes have

the potential to cause insulin resistance. These include proinflammatory cytokines such as Interleukin-6 (IL-6), Tumor Necrosis Factor-TN (TNF-α), angiotensinogen, leptin, resistin, and adiponectin. Adiponectin is a polypeptide that is useful as an antidiabetic, antiarterogenic, anti-inflammatory which is closely related to systemic insulin sensitivity. Adiponectin increases the oxidation of fatty acids in muscles, increases the action of insulin in the liver and decreases lipid accumulation in macrophages.^{7,19,20}

Our study revealed that insulin fasting level was 27.4 IU/L in the NAFLD obese group, not significantly different from the non NAFLD obese group, 26.9 IU/L, but these results were significantly different from the control group with normal nutritional status i.e 13.9 IU/L. One study found that the best independent predictive risk factor for diagnosing NAFLD in obese children was fasting insulin > 18.9 IU / L.⁴ However we did not try to find a predictive risk factor for NAFLD but in both our subject groups with obesity, the fasting insulin greater than 18.9 IU / L were obtained.

All of our obese subjects were insulin resistance, with the HOMA-IR level of obese non NAFLD was 6.1 and obese with NAFLD was 6.8. There was no difference of HOMA-IR level between obese NAFLD and obese without NAFLD. Our results are almost the same as Kim's study in South Korea where the HOMA-IR of obese adolescents with NAFLD was 6.5 but the obese adolescents without NAFLD was 4.2 that is lower than our results.² Kim also found that HOMA-IR (p=0.030) was a significant indicator of NAFLD. The HOMA-IR of overweight adolescents with NAFLD in New Delhi was 4.2 and overweight without NAFLD 3.0, it was lower than our subjects.¹⁷

Adiponectin level was lower in subjects with obesity and NAFLD or without NAFLD than to normal subjects. Our study found a significant differences

between adiponectin levels in normal children (5.1µg / ml) and obese without NAFLD (4.1µg / ml) and obese with NAFLD (4.0µg / ml), but not significant between groups with or without NAFLD. One study in China which compared three obese children groups, namely the group without fatty liver, the group with fatty liver and the group with fatty liver and liver dysfunction, found that the serum adiponectin in the group without fatty liver was significantly higher (4.24µg/ml) compared to subjects with fatty liver (3.37µg/ml) and subjects with fatty liver and liver dysfunction (3.12µg/ml), but did not differ significantly between the fatty liver groups with or without liver dysfunction.¹⁸

In conclusion, we found significant differences of adiponectin levels and insulin resistance measured by HOMA-IR between normal and obese subjects, but no significant differences between the obese groups with or without NAFLD. Likewise, there is no relationship between adiponectin levels and insulin resistance. In suggestion, we need to do a further study regarding the correlation between adiponectin levels and NAFLD in obese adolescents by using liver biopsy.

REFERENCES

1. Neuschwander-Tetri BA. Non-alcoholic fatty liver disease *BMC Medicine* 2017;15:45. Available at <https://bmcmedicine.biomedcentral.com/articles/10.1186/s12916-017-0806-8>
2. Kim JY, Cho J, Yang HR. Biochemical predictors of early onset non-alcoholic fatty liver disease in young children with obesity. *J Korean Med Sci.* 2018;33(16):e122. Available from <https://pubmed.ncbi.nlm.nih.gov/29651819/>
3. Mexitalia M, Utari A, Sakundarno M, Yamauchi T, Subagio HW, Soemantri A. Sindromametabolik pada remajaobesitas (Metabolic syndrome at obese adolescences). *M Med Indones.*2009;43(6):300-5
4. Prokopowicz Z, Malecka-Tendera E, Matusik P. Predictive value of adiposity level, metabolic syndrome, and insulin resistance for the risk of nonalcoholic fatty liver disease diagnosis in obese children. *Can J Gastroenterol Hepatol* 2018 Apr 26; 2018:9465784. Available at <https://www.hindawi.com/journals/cjgh/2018/9465784/>
5. Zou CC, Liang L, Hong F, Fu JF, Zhao ZY. Serum adiponectin resistin levels and non-alcoholic fatty liver disease in obese children. *Endocr J* 2005;52: 519-24.
6. Mathur P, Das MK, Arora NK. Non-alcoholic fatty liver disease and childhood obesity. *Indian J Pediatr.* 2007; 74: 401-7.
7. Junior WS, dos Santos JS, Sankarankutty AK, de Castro e Silva O. Nonalcoholic fatty liver disease and obesity. *Acta Cir Bras* 2006;21: 72-8.
8. Anderson EL, Howel LD, Jones HE, Higgins JPT, Lawlor DA, Fraser A. The prevalence of non-alcoholic fatty liver disease in children and adolescents: a systematic review and meta-analysis. *PLoS One* 2015 Oct 29;10(10):e0140908. Available at <https://pubmed.ncbi.nlm.nih.gov/26512983/>
9. Temple JL, Cordero P, Li J, Nguyen V, Oben JA. A guide to non-alcoholic fatty liver disease in childhood and adolescence. *Int. J. Mol. Sci.* 2016, 17, 947; doi:10.3390/ijms17060947. Available at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4926480/>
10. Saverymuttu SH, Joseph AE, Maxwell JD. Ultrasound scanning in the detection of hepatic fibrosis and steatosis. *Br Med J* 1986;292:13-5 *Br Med J (Clin Res Ed)* 1986 ;292(6512):13-5.
11. Staufer K, Halilbasic E, Spindelboeck W, Eilenberg M, Prager G, Stadlbauer V *et al.* Evaluation and comparison of six non invasive tests for prediction of significant or advanced fibrosis in nonalcoholic fatty liver disease. *United European Gastroenterol J.* 2019; 7(8):1113-23.
12. Asayama K, Hayashibe H, Dobashi K, Uchida N, Nakane T, Kodera K, *et al.* Decrease in serum adiponectin level due to obesity and visceral fat accumulation in children. *Obes Res* 2003;11:1072-7.
13. Winer JC, Zern TL, Taksali SE, Dziura J, Cali AMG, Wollschlaeger M, *et al.* Adiponectin in childhood and adolescent obesity and its association with inflammatory markers and components of the metabolic syndrome. *J Clin Endocrinol Metab* 2006;91:4415-23.
14. Ogawa Y, Kikuchi T, Nagasaki K, Hiura M, Tanaka Y, Uchiyama M. The usefulness of serum adiponectin level as a diagnostic marker of metabolic syndrome in obese Japanese children. *Hypertens Res* 2005;28: 51-7.
15. Jimenez-Rivera C, Hadjiyannakis S, Davila J, Hurteau J, Aglipay M, Barrowman N, *et al.* Prevalence and risk factors for nonalcoholic fatty liver in children and youth with obesity. *BMC Pediatrics.* 2017;17:113. Available at <https://pubmed.ncbi.nlm.nih.gov/28446162/>
16. Keskin M, Kurtoglu S, Kendirci M, Atabek ME, Yazici C. Homeostasis model assessment is more reliable than the fasting glucose/insulin ratio and quantitative insulin sensitivity check index for assessing insulin resistance among obese children and adolescents. *Pediatrics* 2005; 115(4): e500-3. Available at <https://pediatrics.aappublications.org/content/115/4/e500.1> ong
17. Jain V, Jana M, Upadhyay B, Ahmad N, Jain O, Upadhyay AD, *et al.* Prevalence, clinical and biochemical correlates of non-alcoholic fatty liver disease in overweight adolescents. *Indian J Med Res* 2018;148:291-301
18. Aksoy GK, Artan R, Aksoy C, Özdem S, Atalay A, Yilmaz A. Role of soluble adiponectin receptor 2 in non-alcoholic fatty liver disease in children. *Pediatr Gastroenterol Hepatol Nutr.* 2019; 22(5):470-8.
19. Adams LA, Angulo P, Lindor KD. Non alcoholic fatty liver disease. *CMAJ* 2005; 172(7): 899-905.
20. Sanyal AJ. Mechanism of disease: pathogenesis of nonalcoholic fatty liver disease. *Nat Clin Pract Gastroenterol Hepatol* 2005;2(1): 46-53.

The association between adiponectin level and non alcoholic fatty liver disease (NAFLD) in obese adolescents

ORIGINALITY REPORT

10%

SIMILARITY INDEX

5%

INTERNET SOURCES

8%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | www.eje-online.org
Internet Source | 1% |
| 2 | "COPD", Springer Science and Business Media LLC, 2011
Publication | 1% |
| 3 | Branka Filipović, Alastair Forbes, Bojan Tepeš, Dan L. Dumitrașcu. "Nonalcoholic Fatty Liver Disease", Canadian Journal of Gastroenterology and Hepatology, 2018
Publication | 1% |
| 4 | Suna Özdemir, Hüseyin Görkemli, Kazim Gezginç, Mustafa Özdemir, Aysel Kiyici. "Clinical and metabolic effects of medroxyprogesterone acetate and ethinyl estradiol plus drospirenone in women with polycystic ovary syndrome", International Journal of Gynecology & Obstetrics, 2008
Publication | 1% |
| 5 | www.jrmds.in
Internet Source | 1% |

6	journals.sagepub.com Internet Source	1 %
7	www.scilit.net Internet Source	<1 %
8	Yohei OGAWA. "Usefulness of Serum Adiponectin Level as a Diagnostic Marker of Metabolic Syndrome in Obese Japanese Children", Hypertension Research, 2005 Publication	<1 %
9	Masahiro Sogabe, Toshiya Okahisa, Shingo Hibino, Akira Yamanoi. "Usefulness of differentiating metabolic syndrome into visceral fat type and subcutaneous fat type using ultrasonography in Japanese males", Journal of Gastroenterology, 2011 Publication	<1 %
10	dokumen.pub Internet Source	<1 %
11	estudogeral.sib.uc.pt Internet Source	<1 %
12	www.clinicaltrials.gov Internet Source	<1 %
13	Kohtaro Asayama, Hidemasa Hayashibe, Kazushige Dobashi, Norihiko Uchida et al. "Decrease in Serum Adiponectin Level Due to	<1 %

Obesity and Visceral Fat Accumulation in Children", Obesity Research, 2003

Publication

14

M. Brouwers, M.A.L. Bilderbeek-Beckers, A.M. Georgieval, C.J.H. van der Kallen, M.M.J. van Greevenbroek, T.W.A. de Bruin. "W08-P-003 Fatty liver in hypertriglyceridemic subjects with familial combined hyperlipidemia: Importance of visceral fat", Atherosclerosis Supplements, 2005

Publication

<1 %

15

Michel-Henry Mendler, Philippe Bouillet, Anne Le Sidaner, Eric Lavoine, François Labrousse, Denis Sautereau, Bernard Pillegand. "Dual-energy CT in the diagnosis and quantification of fatty liver: limited clinical value in comparison to ultrasound scan and single-energy CT, with special reference to iron overload", Journal of Hepatology, 1998

Publication

<1 %

16

Sanyal, A.J.. "AGA technical review on nonalcoholic fatty liver disease", Gastroenterology, 200211

Publication

<1 %

17

Tsai, P.Y.. "Association between Abnormal Liver Function and Risk Factors for Metabolic Syndrome among Freshmen", Journal of Adolescent Health, 200708

Publication

<1 %

18

doku.pub
Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

The association between adiponectin level and non alcoholic fatty liver disease (NAFLD) in obese adolescents

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
