

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU *PEER REVIEW*
KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Qualitative Tests of Secondary Metabolite Compounds in Ethanol Extract of *Spirulina platensis* from Karimun Jawa Sea, Indonesia

Jumlah Penulis : 3 Orang

Status Pengusul : Renni Yuniati, Muhammad Zainuri, Hermin Kusumaningrum

Identitas Jurnal Ilmiah : a. Nama Jurnal : Biosaintifika: Journal of Biology & Biology Education

b. Nomor ISSN : 2085-191X, 2338-7610

c. Vol, Nomor, halaman : 12 (3), p:343-349

d. Edisi : Desember 2020

e. Penerbit : Universitas Negeri Semarang dan Perhimpunan Biologi Indonesia

f. Jumlah halaman : 9

g. DOI artikel (jika ada) : <http://dx.doi.org/10.15294/biosaintifika.v12i3.23153>

h. Alamat web jurnal : <https://journal.unnes.ac.id/nju/index.php/biosaintifika/article/view/23153/10939>

i. Terindeks di : Sinta 2

j. On line turnitin : <https://doc-pak.undip.ac.id/7661/1/Turnitin-Qualitative-Tests.pdf>

Kategori Publikasi Jurnal Ilmiah : ☐ Jurnal Ilmiah Internasional/ Internasional Bereputasi**
(beri ✓ pada kategori yang tepat) ☒ Jurnal Ilmiah Nasional Terakreditasi
☐ Jurnal Ilmiah Nasional/Nasional

Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah			Nilai Akhir Yang Diperoleh
	Internasional / Internasional Bereputasi **	Nasional Terakreditasi	Nasional *** 	
a. Kelengkapan unsur isi artikel (10%)		25		2,3
b. Ruang lingkup dan kedalaman pembahasan (30%)		7.5		7,0
c. Kecukupan dan kemutahiran data/informasi dan metodologi (30%)		7.5		7,0
d. Kelengkapan unsur dan kualitas terbitan/ jurnal (30%)		7.5		7,2
Total = (100%)		25		23,5
Nilai Pengusul =			60% x 23,5 = 14,1	

Catatan penilaian Artikel oleh Reviewer :

- a. Kelengkapan unsur isi artikel : abstrak singkat dan jelas. Pendahuluan, metode, hasil, pembahasan ditulis sesuai kaidah penulisan ilmiah yg didukung pustaka yang baru dan relevan
- b. Ruang lingkup dan kedalaman pembahasan : lingkup penelitian sesuai dengan bidang ilmu pengusul, dengan melakukan pemeriksaan fito kimia ekstrak *Spirulina platensis* dari Karimunjawa, pembahasan luas dibandingkan dengan penelitian terdahulu
- c. Kecukupan dan kemutahiran data/informasi dan metodologi: studi kuantitatif menganalisis kandungan metabolit antara lain saponin, alkaloid, flavonoid, quinone dan steroid dari ekstrak *S. platensis* Karimunjawa yang didapat melalui maserasi, dengan cara terstandar di lab. Kimia dan Biokimia FK Undip
- d. Kelengkapan unsur dan kualitas terbitan/ jurnal: Biosaintifika: Journal of Biology & Biology Education, Penerbit :Universitas Negeri Semarang dan Perhimpunan Biologi Indonesia adalah Jurnal Ilmiah Nasional terakreditasi S2

Semarang, 13 Desember 2021
Reviewer 1

Prof. Dr. drg. Oedijani, M.S.
NIP. 194902091979012001
Unit kerja : Fakultas Kedokteran Undip
Bidang ilmu : Kedokteran Gigi
Jabatan pangkat : Guru Besar

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU *PEER REVIEW*
KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Qualitative Tests of Secondary Metabolite Compounds in Ethanol Extract of *Spirulina platensis* from Karimun Jawa Sea, Indonesia

Jumlah Penulis : 3 Orang

Status Pengusul : **Renni Yuniati**, Muhammad Zainuri, Hermin Kusumaningrum

Identitas Jurnal Ilmiah : a. Nama Jurnal : Biosaintifika: Journal of Biology & Biology Education
b. Nomor ISSN : 2085-191X, 2338-7610
c. Vol, Nomor, halaman : 12 (3), p:343-349
d. Edisi : Desember 2020
e. Penerbit : Universitas Negeri Semarang dan Perhimpunan Biologi Indonesia
f. Jumlah halaman : 9
g. DOI artikel (jika ada) : <http://dx.doi.org/10.15294/biosaintifika.v12i3.23153>
h. Alamat web jurnal : <https://journal.unnes.ac.id/nju/index.php/biosaintifika/article/view/23153/10939>
i. Terindeks di : Sinta 2
j. On line turnitin : <https://doc-pak.undip.ac.id/7661/1/Turnitin-Qualitative-Tests.pdf>

Kategori Publikasi Jurnal Ilmiah : ☐ Jurnal Ilmiah Internasional/ Internasional Bereputasi**
(beri ✓ pada kategori yang tepat) ☒ **Jurnal Ilmiah Nasional Terakreditasi**
☐ Jurnal Ilmiah Nasional/Nasional


Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah			Nilai Akhir Yang Diperoleh
	Internasional / Internasional Bereputasi **	Nasional Terakreditasi	Nasional ***	
		25		
a. Kelengkapan unsur isi artikel (10%)		2.5		2,5
b. Ruang lingkup dan kedalaman pembahasan (30%)		7.5		7,4
c. Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)		7.5		7,4
d. Kelengkapan unsur dan kualitas terbitan/ jurnal (30%)		7.5		7,5
Total = (100%)		25		24,8
Nilai Pengusul =			60% x 24,8 = 14,88	

Catatan penilaian Artikel oleh Reviewer :

- Kelengkapan unsur isi artikel : Jurnal nasional terakreditasi berbahasa inggris, Sinta 2, unsur lengkap dari abstrak, pendahuluan, metode, hasil pembahasan hingga daftar pustaka. DOI error.
- Ruang lingkup dan kedalaman pembahasan : Penelitian ini bertujuan untuk mengevaluasi senyawa metabolit sekunder yang terkandung dalam ekstrak etanolik *S. platensis* dari Laut Karimun Jawa, Indonesia.
- Kecukupan dan kemutakhiran data/informasi dan metodologi: *S. platensis* yang kering dimaserasi dan diekstraksi menggunakan etanol, kemudian disaring menggunakan kertas saring Whatman. Serangkaian tes dilakukan untuk menentukan metabolit sekunder dalam minyak ter-ozonasi menggunakan beberapa uji kimia dan reagen dalam ekstrak etanol *S. platensis*.
- Kelengkapan unsur dan kualitas terbitan/ jurnal: Terbitan lengkap memuat vol, no, tahun dan daftar isi dan terindeks di Sinta 2.

Semarang,
Reviewer 2



Dr. Diana Nur Afifah, S.TP., M.Si.
NIP. 198007312008012011
Unit kerja : Fakultas Kedokteran Undip
Bidang ilmu : Ilmu Gizi
Jabatan pangkat : Lektor Kepala


[Home](#) > **Vol 13, No 2 (2021)**

Biosaintifika: Journal of Biology & Biology Education

Nationally Accredited SINTA 2 by Ministry of Research and Technology/National Agency for Research and Innovation 2019-2024, Number 36/E/KPT/2019

Biosaintifika: Journal of Biology & Biology Education [p-ISSN 2085-191X] [e-ISSN 2338-7610] published scientific papers on the results of biology and biology education research (see Focus and Scope). Editor accepts the article has not been published in other media with the writing format as listed on page manuscript writing guidelines (see Author Guidelines). The journal has been indexed in Google Scholar, Microsoft Academic, DOAJ, Dimension, DOI Crossref, EBSCO, CABI, SINTA 2 and published three times a year, on April, August & December and published by Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Semarang Perhimpunan Biologi Indonesia in collaboration with Perhimpunan Biologi Indonesia Tingkat Pusat.

Vol 13, No 2 (2021): August 2021 Article in press

Articles in press are peer reviewed, accepted articles to be published in this publication. When the final article is assigned to volumes/issues of the publication, the article in press version will be removed and the final version will appear in the associated published volumes/issues of the publication. The date an article was first made available online will be carried over. Please be aware that, although articles in press do not have all bibliographic details available yet, they can already be cited using the year of online publication and the DOI, as follows: author(s), article title, Publication (year), DOI.

Table of Contents

Articles

The Mitotic Index of *Cajanus cajan* from Kisar Island, in the Southwest of Maluku

Kristin Sangur, Alwi Smith, Meike Tomaso

10.15294/biosaintifika.v13i2.29496 Views of Abstract: 76

The Histomorphometry of Liver and Kidney of Hyperglycemic Albino Rats after Treatment with *Tithonia diversifolia* Leaf Extract

Rizki Fitrawan Yuneldi, Tyas Rini Saraswati, Enny Yusuf Wachidah Yuniwati

10.15294/biosaintifika.v13i2.29459 Views of Abstract: 68

Blood-Glucose Levels of Rats Given High-Fat Diets after Administration of Neem Leaf Ethanol Extract

Sri Isdadiyanto, Siti Muflichatun Mardiaty, Agung Janika Sitasivi

10.15294/biosaintifika.v13i2.29516 Views of Abstract: 25

Conservation Status of Agarwood-Producing Species (*Gyrinops versteegii*) in Indonesia

Sutomo Sutomo, Rajif Iryadi, I Made Sumerta

10.15294/biosaintifika.v13i2.27809 Views of Abstract: 41

Diversity of Pathogenic Fungi and Disease on Vegetable Crops at Polyculture Systems

Eddy Tri Sucianto, Muachiroh Abbas

10.15294/biosaintifika.v13i2.26987 Views of Abstract: 43

Effects of Light Environments on Leaf Traits and Phenotypic Plasticity of *Canna indica*

Yorianta Hidayat Sasaerila, Sakinah Sakinah, Nita Noriko, Risa Swandari Wijihastuti

10.15294/biosaintifika.v13i2.30175 Views of Abstract: 77

Pancreatic Lipase Inhibitory Activity of Endophytic Actinobacteria from *Rhododendron* spp.

Witri Winanda, Irmanida Batubara, Yulin Lestari

10.15294/biosaintifika.v13i2.29981 Views of Abstract: 29

Antidiabetic Potential of Methanol Extract of Flamboyant (*Delonix regia*) Flowers


Kartini Eriani, Uswatun Hasanah, Rizky Fitriana, Widya Sari, Yunita Yunita, Al Azhar



ABOUT THE JOURNAL

[Aims and Scope](#)
[Publication Ethics](#)
[Indexing & Abstracting](#)
[Editorial Team](#)
[Reviewer Team](#)
[Contact](#)

COLLABORATE WITH



perhimpunan biologi indonesia

FOR AUTHORS

[Guidelines for Author](#)
[Manuscript Template](#)
[Mendeley Guidelines](#)
[Online Submission](#)
[Google Citation Analysis](#)
[Scopus Citation Analysis](#)

Readers

	ID 192,637		GB 611
	US 16,747		IR 539
	IN 2,897		AU 498
	MY 2,500		PK 494
	PH 2,004		VN 464
	TH 1,179		MX 421
	CN 1,106		FR 415
	BR 921		EG 386
	JP 874		NL 384
	SG 698		CA 380
	TR 665		DE 377

Pageviews: 647,608

Flags Collected: 180



00583972 [View My Stats](#)

USER

Username

Password

☐ Remember me


[Home](#) > [About the Journal](#) > **Editorial Team**

Editorial Team

Editor-in-Chief

1. Dyah Rini Indriyanti, [SCOPUS ID: 57194044439] Universitas Negeri Semarang, Indonesia

International Editorial Board

1. Fahrul Huyop, [SCOPUS ID:10240240100] Universiti Teknologi Malaysia, Malaysia
2. Mulyoto Pangestu, [SCOPUS ID: 35488897500] Monash University, Australia
3. Mukh Syaifudin, [SCOPUS ID: 6507831675] Badan Tenaga Nuklir Nasional, Indonesia
4. Yohanes Budi Widianarko, [SCOPUS ID: 6507055401] Soegijapranata Catholic University, Semarang, Indonesia
5. Amir Hamidy, [SCOPUS ID: 24341129500] Research Center For Biology-Cibinong Science Center, Indonesia
6. S. Sudarsono, [SCOPUS ID: 16450098400] Institut Pertanian Bogor, Indonesia
7. Iin Kurnia, [SCOPUS ID 57193537585] Badan Tenaga Nuklir Nasional, Indonesia
8. Mohammad Irham, [SCOPUS ID: 55312221800] Research Center for Biology-Indonesian Institute of Sciences, Bogor, Indonesia

Associate Editors

1. Talitha Widiatningrum, [SCOPUS ID: 56798544200] Universitas Negeri Semarang, Indonesia
2. Margareta Rahayuningsih, (SCOPUS ID: 57094066700) Universitas Negeri Semarang, Indonesia
3. Amin Retnoningsih, [SCOPUS ID: 51864647000] Universitas Negeri Semarang, Indonesia
4. Enni Suwarsi Rahayu, [SCOPUS ID: 56644635000] Universitas Negeri Semarang, Indonesia
5. Niken Subekti, [SCOPUS ID: 57195315756] Universitas Negeri Semarang, Indonesia



This work is licensed under a Creative Commons Attribution 4.0 International License.



ABOUT THE JOURNAL

[Aims and Scope](#)
[Publication Ethics](#)
[Indexing & Abstracting](#)
[Editorial Team](#)
[Reviewer Team](#)
[Contact](#)

COLLABORATE WITH



FOR AUTHORS

[Guidelines for Author](#)
[Manuscript Template](#)
[Mendeley Guidelines](#)
[Online Submission](#)
[Google Citation Analysis](#)
[Scopus Citation Analysis](#)

Readers

ID 192,637	GB 611
US 16,747	IR 539
IN 2,897	AU 498
MY 2,500	PK 494
PH 2,004	VN 464
TH 1,179	MX 421
CN 1,106	FR 415
BR 921	EG 386
JP 874	NL 384
SG 698	CA 380
TR 665	DE 377

Pageviews: 647,608

Flags Collected: 180



00583971 [View My Stats](#)

USER

Username

Password

☐ Remember me


[Home](#) > [Archives](#) > **Vol 12, No 3 (2020)**

Vol 12, No 3 (2020)

December 2020

DOI: <https://doi.org/10.15294/biosaintifika.v12i3>

Table of Contents

Editorial

Frontmatter (Cover and Table of Contents)

PDF

Articles

Stability and Antibacterial Property of Polyherbal Mouthwash Formulated Using Local Ingredients

Juman Nafea, Mohamed Edbeib, Kin Israel R. Notarte, Fahrul Huyop, Harisun Yaakub

[DOI: 10.15294/biosaintifika.v12i3.25243](#) Views of Abstract: 256 | PDF: 34

PDF

288-296

The Antioxidant Content and Activity of Various Plant Organs of Kitolod (*Isotoma longiflora*)

Grahita Rezhi Egarani, Sri Kasmiyati, Elizabeth Betty Elok Kristiani

[DOI: 10.15294/biosaintifika.v12i3.23888](#) Views of Abstract: 172 | PDF: 23

PDF

297-303

Identification of Gamma-Amino Butyric Acid Isolates Lactic Acid Bacteria Results from The Isolation of Rusip

S Feta Avila Prayitno, Agung Supriyadi, Endang Kusdiyantini

[DOI: 10.15294/biosaintifika.v12i3.24995](#) Views of Abstract: 79 | PDF: 22

PDF

304-310

Selection of Soybean (*Glycine max*) Germplasm Against Biotrophic Fungi Disease Based on Anatomical Resistance

Siti Samiyarsih, Ade Yuanita Putri Pratiwi, Juni Safitri Muljowati, Nur Fitrianto

[DOI: 10.15294/biosaintifika.v12i3.25109](#) Views of Abstract: 123 | PDF: 18

PDF

311-318

Carotenoid Production by *Rhodospiridium paludigenum* Using Orange Peel Extract as Substrate

Renna Eliana Warjoto, Jennifer Jennifer, Bibiana Widyawati Lay

[DOI: 10.15294/biosaintifika.v12i3.24867](#) Views of Abstract: 130 | PDF: 25

PDF

319-328

Nutritional Content of Bone Flour Made from Plectos Fish *Pterygoplichthys pardalis* from the Ciliwung River, Indonesia

Handhini Dwi Putri, Dewi Elfidasari, Haninah Haninah, Irawan Sugoro

[DOI: 10.15294/biosaintifika.v12i3.23881](#) Views of Abstract: 110 | PDF: 25

PDF

329-334

Potential of Ecology based Agrotourism for Agricultural Education in Besur Village, Lamongan Regency

Slamet Ifandi, Yustika Aulia Rahma

[DOI: 10.15294/biosaintifika.v12i3.25886](#) Views of Abstract: 72 | PDF: 13

PDF

335-342

Qualitative Tests of Secondary Metabolite Compounds in Ethanol Extract of *Spirulina platensis* from Karimun Jawa Sea, Indonesia

Renni Yuniati, Muhammad Zainuri, Hermin Kusumaningrum

[DOI: 10.15294/biosaintifika.v12i3.23153](#) Views of Abstract: 81 | PDF: 24

PDF

343-349

Antioxidant Activity of Jojoba (*Simmondsia chinensis*) Seed Residue Extract

Arnold Patogi Siahaan, Eti Rohaeti, Ali Mahmoud Muddathir, Irmanida Batubara

[DOI: 10.15294/biosaintifika.v12i3.23323](#) Views of Abstract: 157 | PDF: 19

PDF

350-355



ABOUT THE JOURNAL

[Aims and Scope](#)

[Publication Ethics](#)

[Indexing & Abstracting](#)

[Editorial Team](#)

[Reviewer Team](#)

[Contact](#)

COLLABORATE WITH



perhimpunan biologi indonesia

FOR AUTHORS

[Guidelines for Author](#)

[Manuscript Template](#)

[Mendeley Guidelines](#)

[Online Submission](#)

[Google Citation Analysis](#)

[Scopus Citation Analysis](#)

Readers

ID 192,637	GB 611
US 16,747	IR 539
IN 2,897	AU 498
MY 2,500	PK 494
PH 2,004	VN 464
TH 1,179	MX 421
CN 1,106	FR 415
BR 921	EG 386
JP 874	NL 384
SG 698	CA 380
TR 665	DE 377

Pageviews: 647,608

Flags Collected: 180



00583970 [View My Stats](#)

USER

Username

Password

☐ Remember me

 10.15294/biosaintifika.v12i3.23957  Views of Abstract: 50 | PDF: 14

Larvicidal Activity of *Brugmansia candida* against *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae)

Setiawan Khoirul Himmi, Didi Tarmadi, Dita Meisyara, Anugerah Fajar, Titik Kartika, Ikhsan Guswenrivo, Sulaeman Yusuf

PDF

363-369

 10.15294/biosaintifika.v12i3.24634  Views of Abstract: 83 | PDF: 9

Histopathological of White Rats Aorta Induced by High-Fat Feed After Administered by Neem Leaf Ethanolic Extract

Baiq Miftahul Jannah, Sri Isdadiyanto, Agung Janika Sitaswi

PDF

370-377

 10.15294/biosaintifika.v12i3.25494  Views of Abstract: 65 | PDF: 20

***Bifidobacterium longum*, a Predominant *Bifidobacterium* in Early-life Infant Potentially Used as Probiotic**

Dyah Fitri Kusharyati, Anwar Rovik, Dini Ryandini, Hendro Pramono

PDF

378-383

 10.15294/biosaintifika.v12i3.25895  Views of Abstract: 102 | PDF: 19

Alkaline Phosphatase Expression From Mice Mesenchymal Stem Cells Induced By Flamboyant Flower (*Delonix regia*) Extract

Kartini Eriani, Deby Anggraini, Yudha Bintoro, Ichsan Ichsan, Al Azhar, Silmi Mariya

PDF

384-391

 10.15294/biosaintifika.v12i3.25433  Views of Abstract: 33 | PDF: 8

Science Web-Module Integrated with Tannery Waste as Local Potential to Improve Students's Problem Solving

Atznain Iqma Miswami, Heru Nurcahyo

PDF

392-398

 10.15294/biosaintifika.v12i3.24356  Views of Abstract: 49 | PDF: 6

Morphological Variation of Blue Panchax (*Aplocheilichthys panchax*) Lives in Different Habitat Assessed Using Truss Morphometric

Diah Mustikasari, Suhestri Suryaningsih, Agus Nuryanto

PDF

399-407

 10.15294/biosaintifika.v12i3.26593  Views of Abstract: 28 | PDF: 28

Comparison of Nutritional Content of *Mangifera indica* L. var. Wirasangka with Three Other Varieties in Tegal Regency

Enni Suwarsi Rahayu, Amin Retnoningsih, Muhammad Abdullah, Priyanti Susanti

PDF

408-413

 10.15294/biosaintifika.v12i3.26728  Views of Abstract: 10 | PDF: 14

Ethnobotanical Study of Early Childhood Medicinal Plants Used by the Local People in South Bangka Regency, Indonesia

Henri Henri, Vitryany Nababan, Luchman Hakim

PDF

414-421

 10.15294/biosaintifika.v12i3.22221  Views of Abstract: 22 | PDF: 25

Anti-tumor Potential Ethnomedicinal Plants from Dayak Tribes in Central Kalimantan

Liswara Neneng, Asri Pujirahadju, Ferayani Ferayani, Tri Widayat

PDF

422-430

 10.15294/biosaintifika.v12i3.23671  Views of Abstract: 14 | PDF: 16

Blood Hematological and Biochemical Parameters of *Osteochilus vittatus* with *Spirulina platensis* Supplementation in Biofloc System

Muhammad Azharul Rijal, Susanto Susanto, Sorta Basar Ida Simanjuntak, Hernayanti Hernayanti, Purnama Sukardi

PDF

431-437

 10.15294/biosaintifika.v12i3.20936  Views of Abstract: 23 | PDF: 20

Biochemical and Histopathology Analysis of Liver Damage in Hypercholesterolemic Rats Induced by Tomato Extract

Retno Sri Iswari, Muchamad Dafip, Muhammad Rifa'i

PDF

438-445

 10.15294/biosaintifika.v12i3.23337  Views of Abstract: 19 | PDF: 17

Characteristics of DNA Barcodes from Three *Thrixspermum* Orchids Based on ITS2 Regions

Siti Rohimah, Tri Ratnasari, Mukhammad Su'udi

PDF

446-452

 10.15294/biosaintifika.v12i3.23737  Views of Abstract: 19 | PDF: 20

Plantlet Formation and Acclimatization of Sugarcane cv. PS 881 with Different Types and Concentration of Auxin

Rendie Prasetyo, Sugiyono Sugiyono, Elly Proklamasiningsih, Prita Sari Dewi

PDF

453-458

 10.15294/biosaintifika.v12i3.23482  Views of Abstract: 11 | PDF: 11

Isolation of Potential Plant Growth Promoting Rhizobacteria (PGPR) from Cassava (*Manihot esculenta*) Rhizosphere Soil

Suri Raihan Safriani, Lenni Fitri, Yulia Sari Ismail

PDF

459-468

 10.15294/biosaintifika.v12i3.25905  Views of Abstract: 49 | PDF: 36

In Situ Bioremediation Strategies for the Recovery of Mercury-contaminated Land in Abandoned Traditional Gold Mining in Tadaraja

PDF

469-477

Qualitative Tests of Secondary Metabolite Compounds in Ethanol

by Renni Yuniati

Submission date: 20-Aug-2021 11:03AM (UTC+0700)

Submission ID: 1633496025

File name: 23153-69098-1-PB.pdf (300.33K)

Word count: 4048

Character count: 23425

Qualitative Tests of Secondary Metabolite Compounds in Ethanol Extract of *Spirulina platensis* from Karimun Jawa Sea, Indonesia

Renni Yuniati^{1*}, Muhammad Zainuri², Hermin Kusumaningrum³

¹Department of Dermatology and Venereology, Faculty of Medicine, Universitas Diponegoro, Indonesia

²Department of Oceanography, Universitas Diponegoro, Indonesia

³Department of Biology, Faculty of Science and Mathematics, Universitas Diponegoro, Indonesia

*Email: renniyuniati@yahoo.com

Submitted: 22 January 2020, Revised: 27 July 2020, Accepted: 24 September 2020

Abstract. *Spirulina platensis* is a spiral green-blue algae that is abundant in the subtropical and tropical water bodies. *S. platensis* has several potentials to be used in the medical fields due to its wound healing and antitumor potentials. This research aimed to evaluate the secondary metabolite compounds contained in the ethanolic extract of *S. platensis* from Karimun Jawa Sea, Indonesia. The dried *S. platensis* was macerated and extracted using ethanol, then was filtered using Whatman filter paper. A series of tests has been done to determine the secondary metabolites in the ozonated oil using several chemistry tests and reagents in the ethanol extract of *S. platensis*. The tests found that *S. platensis* microalgae harvested from Karimun Jawa sea contains alkaloid, saponin, flavonoid, and quinone. As *S. platensis* from different region contains different substances, this research shows the substances contained in *S. platensis* from Karimun Jawa sea. With this research, we knew that *S. platensis* from Karimun Jawa islands contains alkaloid, saponin, flavonoid, and quinone which can be used as a medicinal ingredient.

Key words: Extraction; Secondary Metabolite; *S. platensis*

How to Cite: Yuniati, R., Zainuri, M., & Kusumaningrum, H. (2020). Qualitative Tests of Secondary Metabolite Compounds in Ethanol Extract of *Spirulina platensis* from Karimun Jawa Sea, Indonesia. *Biosaintifika: Journal of Biology & Biology Education*, 12(3), 343-349.

DOI: <http://dx.doi.org/10.15294/biosaintifika.v12i3.23153>

INTRODUCTION

Spirulina platensis is a species of filamentous photoautotrophic green-blue algae that belongs to cyanobacteria class (Saranraj & Sivasakthi, 2014; Suratno, 2010; Wollina et al., 2018a). Cyanobacteria are one of the oldest forms of living things found in the Earth. They share similarities to the primitive bacteria due to their lack of cell walls and with plants because of their photosynthesis ability. *Spirulina* is one of the edible cyanobacteria along with some other species such as *Nostoc* and *Aphanizomenon* (Saranraj & Sivasakthi, 2014).

The filaments of *Spirulina* is small, measuring about 0.1 mm, coiled in variable tightness and number of spirals (Saranraj & Sivasakthi, 2014). Under the light microscope, *Spirulina* can be seen as small (6-12 micrometer wide) trichomes which are coated in a thin layer of wall. The trichome has greenish-blue color (hence its name), and has the ability to move freely (Vonshak, n.d.). Several environmental factors, including temperature and physicochemical conditions can affect its growth. *S. platensis* can be found in tropical and subtropical areas with warm temperature, as it grows optimally at the 30-37 degrees Celcius temperature area with alkaline pH and high bicarbonate content (S. K. Ali & Saleh, 2012; Saranraj & Sivasakthi, 2014). Salinity is

also one of the factors affecting the growth of *Spirulina*. *Spirulina* is mostly found in areas with high concentration of sodium chloride (NaCl) and bicarbonate, which resulted in higher content of its phycocyanin pigment (Christwardana et al., 2013; Saranraj & Sivasakthi, 2014).

Recently, *S. platensis* is being studied because of its high nutritional content which can be used in the medical field (Saranraj & Sivasakthi, 2014). Its interest is getting higher since a lot of people are trying to use "back to the nature" concept of using natural ingredients as medical therapeutic agents (Ergina & Pursitasari, 2014). However, further development of drugs and medicinal products are engineered towards the active contents contained in the medicinal product ingredient; in this case *S. platensis*. Therefore, it is important to determine what are the substances contained in the *S. platensis* which can be used as a basis for further drug development. *S. platensis* is mostly composed of proteins (around 50-70% of the dry weight). The essential amino acids contained in *Spirulina* include leucine, valine, and isoleucine (Sotiroidis & Sotiroidis, 2013). *Spirulina*'s green color comes from phycobiliproteins contained within *Spirulina* cells which serves as a light-collecting pigment for its photosynthesis activity. Phycobiliproteins consist of three pigments i.e. allophycocyanin, phycocyanin, and

phycoerythrin. From these three pigments, phycocyanin is the pigment that is mostly studied due to its potential benefits in the medicine field (Saranraj & Sivasakthi, 2014).

Several studies have found health benefits of *S. platensis*. *S. platensis* are found to have antifungal activity against several kinds of fungi, including *Aspergillus fumigatus*, *Mucor vulgaris*, *Penicillium expansum*, *Fusarium solani*, and *Fusarium oxysporum*; all of which contributes to the invasive mycoses infection (Otlu & Rudic, 2016). Phycocyanin pigment of *S. platensis* has been found to be able to reduce edema, histamine release, myeloperoxidase activity of macrophages, and reduces prostaglandin E2 and leukotriene LTB4 in inflammatory lesions (Wollina et al., 2018b). *Spirulina* is also found to be able to improve wound healing by increasing fibroblast proliferation and migration, enhancing wound closure rate, and inhibiting colonization of *Staphylococcus aureus* in wounds (Wollina et al., 2018a). Anticancer activities can also be found in *S. platensis*. One research found that *S. platensis* has anti skin tumor effects in mice that is exposed to ultraviolet B (UVB) rays (Yogianti et al., 2014).

As with many other living organisms, *S. platensis* also produces secondary organic metabolites. Secondary metabolites are substances that are produced as a result of metabolism through secondary reactions of primary organic substances (carbohydrates, fats, and proteins). These precursor substances that are used in the biosynthesis of secondary metabolites came from primary metabolism process (Mardani, 2013). Secondary metabolites are small, specific and unique metabolites. Not all organism will have the similar configuration and amount of secondary metabolites. Secondary metabolites has the potential to be used as a lead compounds that can be used in the drug development (Ergina & Pursitasari, 2014).

Secondary metabolites are classified into three groups: phenolic, alkaloid, and terpenoids. Most common secondary metabolites which can be found in plants are alkaloid, flavonoid, steroid, saponin, terpenoid, and tannin (Ergina & Pursitasari, 2014). Terpenes are substances formed by a chain of isoprenes, which carbon structure is formed by a link of two or more C5 carbons. Terpenes consist of several substances such as monoterpenes and sesquiterpenes which are volatile substances, diterpenes which are semi-volatile, and triterpenes and sterols which are non-volatile (Khotimah & Minarno, 2016). Terpenoids are a group of secondary metabolites that is effective to be used as plants-originated insecticides (Mardani, 2013).

Alkaloids are basic substances that contains at least one nitrogen atom which form heterocyclic chain. Most alkaloids are colorless and has active optic properties. Flavonoids are secondary metabolites which are synthesized from pyruvate acids through the metabolism of amino acids. Flavonoids are mostly found in the form of glycosides on one or more groups of phenolic hydroxyls (Khotimah & Minarno, 2016). There are several kinds of flavonoids which can be identified, including anthocyanin, proanthocyanidins, flavanols, flavone, glycoflavone, biflavonyle, chalcone, aurone, flavanone, and isoflavone (Khotimah & Minarno, 2016). This experiment aimed to determine the secondary metabolites contained in *S. platensis* extract, such as alkaloid, saponin, phenol, flavonoids, quinone, triterpenoid, and steroid. We hoped that this research helps scientists and pharmacists to discover the contents of *S. platensis* from Karimun Jawa and enables the usage of *S. platensis* from Karimun Jawa islands to be used as a pharmaceutical product.

METHODS

This research was done in Chemistry and Biochemistry Laboratory, Faculty of Medicine Diponegoro University, Indonesia. The *S. platensis* algae were obtained from Karimun Jawa Sea, Central Jawa, Indonesia. All of the tests done to analyze the presence of the secondary metabolites were done with *S. platensis* extract as the test sample using several tests. In this research, we analyzed the presence alkaloid, saponin, phenol, flavonoids, quinone, triterpenoid, and steroid in the *S. platensis* extract.

This experiment began with the maceration of *S. platensis* sample. The maceration process was done to extract the active ingredients contained within the *Spirulina* cells, so that they can be analyzed once they are no longer trapped inside the cells. The liquid then was filtered using Whatman filter paper to separate the solute from the solids, before further evaporation to obtain a thick extract of *S. platensis*.



Figure 1. Microscopic image of *Spirulina platensis* (Whitton, 2012)

Extraction of *S. platensis*

The raw *S. platensis* algae gathered from the Karimun Jawa Sea were dried. The dried *S. platensis* were crushed into fine powder. The powdered *S. platensis* microalgae was macerated in 95% ethanol solution with 1:10 concentration (one part of *S. platensis* powder macerated in 10 parts of 95% ethanol solution). The maceration process was done for five days in a glass container. The glass container was stirred every day to make sure the uniformity of the maceration process. After five days, the solution was filtered through Whatman filter paper, and was evaporated using rotary evaporator machine at the ethanol boiling point temperature until a thick extract was obtained. This extract of *S. platensis* was used as the material in the subsequent tests.

Identification of alkaloid

The identification of alkaloid in this study were done by diluting a few drops (around 1 milligram in weight) of *S. platensis* extract into five milliliters of 2 N sulfuric acid. The mixture of *S. platensis* extract and 2 N sulfuric acid were subsequently tested using 5 drops of both Mayer's and Liebermann-Burchard reagents on separate test tubes (Khotimah & Minarno, 2016; Suratno, 2010). The formation of white precipitation at the bottom of the test tube after the reagent was added shows the presence of alkaloid substances (Akerina & Sangaji, 2019; Ergina & Pursitasari, 2014; Rasyid, 2012).

Identification of saponin

Identification of saponin were done using foam test. *Spirulina* extract was diluted into 9 ml of water and then heated for 5 minutes. The solution then was agitated vertically for 10 seconds. The agitation process will result in the formation of foam layer on top of the solution. Afterwards, one drop of 2 N hydrochloric acid was added into the solution. The presence of saponin can be confirmed by the formation of foams (at least 1 cm high) that remains on the top of the solution for at least 10 minutes and does not subside with the addition of 2 N HCL (Rasyid, 2012; Suratno, 2010).

Identification of phenolic substances

The identification of phenolic substances in this study were done using FeCl₃ solution as the reagent. One milliliter of *Spirulina* ethanol extract was mixed with 2 drops of 5% FeCl₃ in a test tube. The presence of phenols can be identified by the formation of green

or greenish-blue color (Akerina & Sangaji, 2019; Suratno, 2010).

Identification of flavonoids

The presence of flavonoids in the ethanol extract of *S. platensis* were identified using Willstatter test. This test uses magnesium strips and hydrochloric acid-ethanol mixture as the test reagent. One milliliter of *S. platensis* ethanol extract was put into a test tube. Subsequently, 0.1 mg of magnesium and 0.4 ml of amyl alcohol (a mixture of 37% hydrochloric acid and 95% ethanol with similar volume) was added into the test tube. Positive result can be seen by the color change of the solution, which will turn dark yellow or orange (Khotimah & Minarno, 2016; Suratno, 2010).

Identification of quinone

The identification of quinone in this study was evaluated using sodium hydroxide (NaOH). One milliliter of sample was mixed with 1N NaOH and then was evaluated for color changes. Positive result can be seen by the formation of yellow or yellowish-green color resulted from the reaction (Suratno, 2010).

Identification of triterpenoids and steroids

The presence of triterpenoid and steroid in this study were evaluated using chloroform, sulfuric acid, and anhydrous acetate. A few drops of sample were added into a test tube containing 2 mL chloroform. Subsequently, 10 drops of anhydrous acetate and 3 drops of sulfuric acid were added into the test tube. The positive results can be seen from the color change of the sample: initially, the color will turn into red; afterwards it will change into blue or green (Suratno, 2010). The red color indicates the presence of terpenoids, while the green color indicates the presence of steroid (Ergina & Pursitasari, 2014).

RESULTS AND DISCUSSION

In this research, we conducted the tests explained in the above paragraphs to our *S. platensis* sample from Karimun Jawa sea. The positive result from the Mayer's reagent showed that *S. platensis* extract contains alkaloid substance. However, alkaloid identification using Liebermann-Beurchard reagent did not give positive result. This result was similar to the previous identification result of two different researches by Mane & Chakraborty (2018) and Suratno (2010).



Figure 2. Positive alkaloid test using Mayer's reagent on *S. platensis*

After agitation of the mixture, a layer of foam was present on the top part of the mixture, which was present until 10 minutes were elapsed. The foam did not subside upon the addition of one drop of hydrochloric acid. Therefore, this result indicated that *S. platensis* extract contains saponin. This result were in line with previous researches (Fithriani et al., 2015; Mane & Chakraborty, 2018; Saputra, 2018; Suratno, 2010) which found that *Spirulina* sp. contains saponin.



Figure 3. Positive saponin test using foam test on *S. platensis*

Phenolic substance identification test results in color change from green to yellow after the addition of FeCl_3 was observed. Therefore, this result showed that the sample of *S. platensis* does not contain phenolic substances.



Figure 4. Negative phenolic identification test using FeCl_3 reagent on *S. platensis*

We noticed a color change from dark green to yellow was observed in our flavonoid test. This result showed that the *S. platensis* sample positively contains chalcone, aurone, and flavone content. Several previous researches also found similar results regarding the identification of *S. platensis* extract from another region, including in India (Fithriani et al., 2015; Mane & Chakraborty, 2018).



Figure 5. Positive flavonoid test result using magnesium and hydrochloric acid-alcohol reagent on *S. platensis*

We observed a greenish-yellow color in our quinone test, which confirmed that the *S. platensis* sample in this research contains quinone. This result was similar to the previous research conducted by Suratno, (2010), but not with the other researches.



Figure 6. Positive quinone test using sodium hydroxide reagent on *S. platensis*

The triterpenoids and steroids identification test, the color was initially brown, which at the end of the observation period turned into yellowish-green color. This result showed that our sample of *S. platensis* does not contain triterpenoids and steroids.



Figure 7. Negative triterpenoid and steroid identification test using chloroform, anhydrous acetate and H_2SO_4 reagent on *S. platensis* sample from Karimun Jawa Sea, Indonesia.

In this research, the sample of *S. platensis* microalgae that we harvested from Karimun Jawa sea contains alkaloid, saponin, flavonoid and quinone. This result is mostly similar to the previous research conducted by (Suratno, 2010), where he found that the *S. platensis* sample from Lamongan, East Jawa contains alkaloid, saponin, phenol, and quinone.. Another result conducted by (Saputra, 2018) found that ethanolic extract of *S. platensis* contains flavonoids, saponin, phenolic substances, and steroids. A study conducted using laboratory-grown *Spirulina* sp. sample which was planted in sterile water medium found that *Spirulina* sp. contains tannin, flavonoids, steroids, glycosides, alkaloids, and saponin (Fithriani et al., 2015).

The active metabolites of *Spirulina platensis* are tightly affected by the environment they are grown with (Notonegoro et al., 2018). Results of phytochemical analysis of *S. platensis* extract from

Rankala Lake, Kolhapur, India was slightly different from our study. They found that their *S. platensis* ethanolic extract contains alkaloids, terpenoids, steroids, saponins, flavonoids, phenols, and coumarins (Mane & Chakraborty, 2018). The phytochemical contents were differing from one solvent to another; they found that methanol can extract every single metabolite from *S. platensis* powder, and the other solvents (ethanol, petroleum ether, acetone, and aqueous extract) does not have the same yield (Mane & Chakraborty, 2018). A quantitative study on *S. platensis* extract from Egypt found that *S. platensis* contains relatively high phycobiliproteins, phenolic substance, alkaloids, and terpenoids (Ali et al., 2014).

Saponin is a glycoside that can be found in many organisms. The saponin contents of an organism is highly affected by the variety and stage of growth. The function of saponin are most likely as a storage form of carbohydrate or might be related to defending mechanism against insects. The test for determining the presence of saponin glycosides involves the ability of the shaken extract in the water to form a stable foam that does not subside after a brief period; afterall, saponins itself are characterized by their ability to form strong foams in an aqueous solution (Man et al., 2010; Nurwidayati, 2012). Saponins are known to exhibit several medicinal properties, for example as an antidermatophytic agent and antimicrobial activity. Saponins also has the properties to reduce *Malassezia furfur* growth, which can be used as an anti-dandruff formula in shampoo (Tamura et al., 2012). Saponins also exhibited anti-cancer properties in terms of inhibiting tumor angiogenesis. It works by suppressing cancer angiogenesis inducer in blood vessel endothelial cells and also prevents the adherence, invasion, and metastasis of tumor cells (Man et al., 2010).

The principle that was used in the alkaloid tests are precipitation reaction caused by ligand exchange. The nitrogen atoms that has free electrons in the alkaloid can replace the iodo ion in the reagent (Khotimah & Minarno, 2016). Most alkaloids are not soluble in water, except if they reacted with acid which causes them to form water-soluble alkaloid salt complex (Khotimah & Minarno, 2016). The white precipitation resulted from the reaction is potassium-alkaloid complex. The nitrogen in the alkaloid will react with the potassium ion from the potassium tetraiodomercurate(II) which forms potassium-alkaloid complex that precipitates in the bottom of the reaction tube (Ergina & Pursitasari, 2014; Suratno, 2010).



Figure 8. Reaction of alkaloid identification (Ergina & Pursitasari, 2014)

The flavonoid test in this experiment was done using magnesium and hydrochloric acid. The addition of magnesium and hydrochloric acid was used to reduce the benzopyrone which is contained in the flavonoid structure. The result of this reaction is the formation of flavylium salts, which has the red or orange color. Flavonoid itself is a substance that contains two aromatic rings with more than one hydroxyl groups (Ergina & Pursitasari, 2014).




Figure 9. Reaction of flavonoid with magnesium and hydrochloric acid (Ergina & Pursitasari, 2014)

The *S. platensis* sample in our study contains quinone. The principle of quinone identification test were based on the colorimetric reaction. The reaction of quinone identification test involves reciprocal reduction which converts quinone into a colored substance which can be identified with naked eye. This reduction reaction will subsequently be reverted if the solution were exposed to the air (Khotimah & Minarno, 2016).

This research enables *S. platensis* from Karimun Jawa Islands, Indonesia to be used as an ingredient for herbal preparation which needs alkaloids, saponin, flavonoids, and quinone. Alkaloids has several medicinal properties, including analgesic, anti-cancer, anti-arrhythmic, anti-hyperglycemic, and anti-psychotropic effects (Hussain et al., 2018). *S. platensis* saponins are known to be cytotoxic to cancer cells and has the potential to decrease cancer cellular viability; thus enables its use as an anticancer candidate (Akbarizare et al., 2019). Alkaloids and quinones are also can be utilized as medicinal ingredient. Different region and different growth conditions results in different substances contained in the *S. platensis* algae. Therefore, this result shows the contents of *S. platensis* from Karimun Jawa islands. We hope that this research increases the use of *S.*

platensis as an ingredient of healthy foods and medicines with potent efficacy.

CONCLUSION

S. platensis microalgae that we harvested from Karimun Jawa sea contains alkaloid, saponin, flavonoid, and quinone.

REFERENCES

- Akbarizare, M., Ofoghi, H., & Hadizadeh, M. (2019). In vitro Anticancer Evaluation of Saponins Obtained From *Spirulina platensis* on MDA, HepG2, and MCF7 Cell Lines. *Multidisciplinary Cancer Investigation*, 3(4), 25–32.
- Akerina, F. O., & Sangaji, J. (2019). Analisis Fitokimia dan Toksisitas serta Aktivitas Antioksidan Beberapa Jenis Teripang di Desa Kakara, Halmahera Utara. *Agrikan: Jurnal Agribisnis Perikanan*, 12(2), 188.
- Ali, H. E. A., Shanab, S. M. M., Abo-State, M. A. M., Shalaby, E. A. A., El Demerdash, U. M. N., & Abdullah, M. A. (2014). Evaluation of Antioxidants, Pigments and Secondary Metabolites Contents in *Spirulina platensis*. *Applied Mechanics and Materials*, 625, 160–163.
- Ali, S. K., & Saleh, A. M. (2012). *Spirulina* - An overview. *International Journal of Pharmacy and Pharmaceutical Sciences*, 4(SUPPL.3), 9–15.
- Christwardana, M., Nur, M. M. A., & Hadiyanto, H. (2013). *Spirulina platensis*: Potensinya Sebagai bahan Pangan Fungsional. *Jurnal Aplikasi Teknologi Pangan*, 2(1).
- Ergina, S. N., & Pursitasari, I. D. (2014). Qualitative Test of Secondary Metabolites Compounds in Palado Leaves (*Agave Angustifolia*) Extracted With Water and Ethanol. *Jurnal Akademika Kimia*, 3(3), 165–172.
- Fithriani, D., Amini, S., Melanie, S., & Susilowati, R. (2015). Phytochemical Screening, Total Phenol Content and Antioxidant Activity of Microalgae *Spirulina* sp., *Chlorella* sp. and *Nannochloropsis* sp. *Jurnal Pascapanen Dan Bioteknologi Kelautan Dan Perikanan*, 10(2), 101.
- Hussain, G., Rasul, A., Anwar, H., Aziz, N., Razzaq, A., Wei, W., Ali, M., Li, J., & Li, X. (2018). Role of plant derived alkaloids and their mechanism in neurodegenerative disorders. *International Journal of Biological Sciences*, 14(3), 341–357.
- Khotimah, K., & Minarno, E. B. (2016). *Screening and Identification of Metabolites Secondary Phytochemical Compounds Carpain in Methanol Extract of Leaves of Carica pubescens Lenne & K. Koch with LC/MS (Liquid Chromatography -*

- tandem Mass Spectrometry). State Islamic University Maulana Malik Ibrahim.
- Man, S., Gao, W., Zhang, Y., Huang, L., & Liu, C. (2010). Chemical study and medical application of saponins as anti-cancer agents. *Fitoterapia*, 81(7), 703–714.
- Mane, R. S., & Chakraborty, B. (2018). Phytochemical screening of Spirulina platensis extracts from Rankala Lake. *Journal of Algal Biomass Utilization*, 9(4), 38–41.
- Mardani, Y. (2013). *Identifikasi Senyawa Metabolit Sekunder pada Tanaman Sirsak (Annona muricata L.), Srikaya (Annona squamosa L.), dan Mimba (Azadirachta indica A.Juss.)*. Universitas Muhammadiyah Malang.
- Notonegoro, H., Setyaningsih, I., & Tarman, K. (2018). The Active Compounds of Spirulina platensis Grown on Walne Media with Different NaNO₃ Concentrations. *Jurnal Pascapanen Dan Bioteknologi Kelautan Dan Perikanan*, 13(2), 111.
- Nurwidayati, A. (2012). *The phytochemical screening and thin layer chromatography results of Jatropha gossypifolia seeds*. 3(2), 27–31.
- Otlu, I., & Rudic, V. (2016). Antifungal activity of extracts from Arthrospira platensis against some pathogens, causing invasive mycoses. *Curierul Medical*, 59(6), 9–14.
- Rasyid, A. (2012). Identification of Secondary Metabolites Compounds, Antibacterial and Antioxidant Activities on the Methanol Extract of Sea Cucumber Stichopus hermanii. *Jurnal Ilmu Dan Teknologi Kelautan Tropis*, 4(2), 360–368.
- Saputra, H. R. (2018). *Screening of Phytochemicals and Antibacterial Activity Test of Ethanol Extract Spirulina platensis against Methicillin Resistant Staphylococcus aureus(MRSA) and Vancomycin Resistant Enterococci(VRE)*. Sriwijaya University.
- Saranraj, P., & Sivasakthi, S. (2014). Spirulina platensis - food for future: a review. *Asian Journal of Pharmaceutical Science & Technology*, 4(1), 26–33.
- Sotiroudis, T., & Sotiroudis, G. (2013). Health aspects of Spirulina (Arthrospira) microalga food supplement. *Journal of the Serbian Chemical Society*, 78(3), 395–405.
- Suratno. (2010). Phytochemical Screening of Spirulina Platensis Microalgae Etanol Extract Which Has the Potential as an Antibacterial Agent. *Jurnal Surya Medika*, 1(2), 27–33.
- Tamura, Y., Miyakoshi, M., & Yamamoto, M. (2012). Application of Saponin-Containing Plants in Foods and Cosmetics. In *Alternative Medicine*. InTech.
- Vonshak, A. (n.d.). *Spirulina Platensis Arthrospira : Physiology, Cell-Biology And Biotechnology*.
- Whitton, B. A. (2012). *Ecology of cyanobacteria II: their diversity in space and time*. Springer.
- Wollina, U., Voicu, C., Gianfaldoni, S., Lotti, T., França, K., & Tchernev, G. (2018a). Arthrospira Platensis - Potential in Dermatology and Beyond. *Open Access Macedonian Journal of Medical Sciences*, 6(1), 176–180.
- Wollina, U., Voicu, C., Gianfaldoni, S., Lotti, T., França, K., & Tchernev, G. (2018b). Arthrospira Platensis - Potential in Dermatology and Beyond. *Open Access Macedonian Journal of Medical Sciences*, 6(1),
- Yogianti, F., Kunisada, M., Nakano, E., Ono, R., Sakumi, K., Oka, S., Nakabeppu, Y., & Nishigori, C. (2014). Inhibitory Effects of Dietary Spirulina platensis on UVB-Induced Skin Inflammatory Responses and Carcinogenesis. *Journal of Investigative Dermatology*, 134(10),

Qualitative Tests of Secondary Metabolite Compounds in Ethanol

ORIGINALITY REPORT

9%

SIMILARITY INDEX

3%

INTERNET SOURCES

4%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to Sriwijaya University

Student Paper

3%

2

Dwiana Intan Pertiwi, Rifda Naufalin, Poppy Arsil, Erminawati, Rumpoko Wicaksono, Taslimatul Auliya. "QUALITY OF SIMPLICIAN BIOACTIVE COMPONENTS AND LIQUID EXTRACT OF KECOMBRANG FLOWER POWDER FROM TEMPERATURE AND TIME OPTIMIZATION RESULTS", IOP Conference Series: Earth and Environmental Science, 2019

Publication

1%

3

Natural Products, 2013.

Publication

1%

4

Wilis A. Setyati, Rini Pramesti, Muhamad Zainuddin, Maya Puspita, Person P Renta. "Cytotoxicity and Phytochemical Profiling of Extract As Anti-Mdr Bacteria ", IOP Conference Series: Earth and Environmental Science, 2018

Publication

1%

5	ejournal.undip.ac.id Internet Source	<1 %
6	Y H Teffu, A G O Kase, A Trianto, D P Wijayanti. " Secondary metabolites compound of sea plumes gorgonian () from Maumere Water- East Nusa Tenggara Indonesia ", IOP Conference Series: Materials Science and Engineering, 2020 Publication	<1 %
7	journal.unnes.ac.id Internet Source	<1 %
8	mafiadoc.com Internet Source	<1 %
9	bjbas.springeropen.com Internet Source	<1 %
10	downloads.hindawi.com Internet Source	<1 %
11	qascf.com Internet Source	<1 %
12	Mehdi Chaouachi, Sandrine Gautier, Yoann Carnot, Nicolas Bideau et al. " Provides a Small Advantage in Vertical Jump and Sprint Performance But Does Not Improve Elite Rugby Players' Body Composition ", Journal of Dietary Supplements, 2020 Publication	<1 %

13

Sariya Mapoung, Punnida Arjsri, Pilaiporn Thippraphan, Warathit Semmarath et al.
"Photochemoprotective effects of Spirulina platensis extract against UVB irradiated human skin fibroblasts", South African Journal of Botany, 2020

Publication

<1 %

14

www.freepatentsonline.com

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Qualitative Tests of Secondary Metabolite Compounds in Ethanol

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
