

Differences of maternal leukocyte count in premature rupture of membranes and preterm premature rupture of membranes

by Putri Sekar Wiyati

Submission date: 09-Dec-2021 11:47AM (UTC+0700)

Submission ID: 1725172855

File name: Differences_of_maternal_leukocyte_count_in_premature_rupture.pdf (229.57K)

Word count: 2312

Character count: 12431



Published by DiscoverSys

Differences of maternal leukocyte count in premature rupture of membranes and preterm premature rupture of membranes



CrossMark

Besari Adi Pramono^{1*}, Putri Sekar Wiyati², Julian Dewantiningrum¹

ABSTRACT

Background: Premature rupture of membranes (PROM) is rupture of membranes amniotic membrane prior to the labor. PROM is associated with an increased incidence of preterm labor and infection.

Aim: To compare maternal leukocyte count in premature rupture of membranes (PROM) and preterm premature rupture of membranes (PPROM).

Methods: This cross-sectional design was conducted at Dr. Kariadi General Hospital Medical Center and Kartini General Hospital Jepara from September 2019 to January 2020. The study sample was pregnant women with premature rupture of membranes that came

to Emergency Department. Samples were divided into two groups, PROM and PPRM. All samples were subjected to leukocyte count examination.

Results: This study has a total of 40 samples, with 20 samples each. In PROM, 8 (40%) leukocytosis and 12 (60%) were normal. In PPRM 7 (35%) leukocytosis and 13 (65%) were normal. There were no significant differences in maternal leukocyte levels in PROM and PPRM ($p=0.229$).

Conclusion: There were no significant differences in maternal leukocyte levels in PROM and PPRM.

Keywords: Premature rupture of membranes, aterm, preterm, leukocyte

Cite this Article: Pramono, B.A., Wiyati, P.S., Dewantiningrum, J. 2020. Differences of maternal leukocyte count in premature rupture of membranes and preterm premature rupture of membranes. *Bali Medical Journal* 9(3): 868-871. DOI: [10.15562/bmj.v9i3.2149](https://doi.org/10.15562/bmj.v9i3.2149)

¹Maternal Division, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Diponegoro/Dr Kariadi General Hospital Medical Center, Semarang

²Social Obstetrics and Gynecology Division, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Diponegoro/Dr Kariadi General Hospital Medical Center, Semarang

*Corresponding to:
Besari Adi Pramono; Fetomaternal Division, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Diponegoro/Dr Kariadi General Hospital Medical Center, Semarang; babas1504@yahoo.com

Received: 2020-10-23
Accepted: 2020-11-28
Published: 2020-12-01

INTRODUCTION

Premature rupture of membrane (PROM) is defined as the rupture of the amniotic membrane prior to the labor. Premature rupture of membranes can occur during or after 37 weeks of gestation (premature rupture of membranes / PROM), or before 37 weeks of gestation (preterm premature rupture of membranes / PPRM).¹ Under normal circumstances, 8-10% women with a term pregnancy will experience PROM.²

Currently, the problem of PROM needs a greater attention, because its prevalence is quite large and the incidence of infection tends to increase. The incidence of term PROM occurs in approximately 6.46-15.6% of term pregnancies and preterm PROM occurs in approximately 2-3% of all singleton pregnancies and 7.4% of multiple pregnancies. The incidence rate of PROM in Indonesia ranges from 4.5-6%. In Dr. Kariadi General Hospital Medical Center, the incidence rate of PROM in 2012 was 6.1%.³

Leukocytes are cells of the immune system that play a role in protecting the body from infection and foreign bodies. Leukocytes are produced by multipotent cells in the spinal cord. Leukocytes have a larger size than erythrocytes. The leukocyte

count is a test done to count the number of leukocytes in 1 ml of blood.⁴ Physiological levels of normal leukocyte counts in pregnancy tend to be higher, ranging from 5,600-13,800 mm.^{3,5} In PROM, leukocytes play a role in activating cytokines that stimulate prostaglandin production and metalloproteinase matrix (MMP).⁶

Leukocytosis is one of the sign of clinical chorioamnionitis or intraamniotic infection (IAI) or Triple I (Intrauterine infection or inflammation or both) which can occur in 40-70% of patients with PROM.⁷ In preterm PROM with maternal leukocytosis and managed as expectative management, poor infant outcomes can happen in the form of neurodevelopmental disorders during the first 2 years of life.⁸

Therefore, the authors intended to conduct a research on differences in levels of maternal leukocyte counts in patients with PROM and PPRM.

METHOD

This study was a cross-sectional study, which was conducted at Dr. Kariadi General Hospital Medical Center Semarang and Kartini General Hospital Jepara from September 2019 to January 2020.

12 sample of the study were pregnant women with premature rupture of membranes in the Emergency Room of both hospitals, which meet the inclusion 19 criteria, namely primigravida and multigravida with a gestational age of ≥ 20 weeks, and a single live intrauterine fetus. Patients with a history of cardiovascular disorders, a history of chronic hypertension, diabetes mellitus, kidney disease, chronic and systemic infections, smoking habits, drinking alcohol, use of drugs containing estrogen and progesterone during pregnancy, obesity, intrauterine fetal death, and antepartum hemorrhage were excluded from the study.

Sampling was done by block randomization. All samples were subjected to anamnesis examination, physical examination, and laboratory blood tests as indicated. Blood samples used vein blood as much as 5 cc and checked with a hematology analyzer machine.

Statistical 16 analysis was performed using the Shapiro-Wilk test to see the distribution of primary data distribution. Mann-Whitney test was used to analyze the levels of leukocyte counts.

Table 1. Characteristics of sample

Characteristics	Type of Rupture of Membrane		P
	PROM (n = 20)	PPR 18 (n = 20)	
Age (years) (Mean \pm SD; Median-min-max)	29.9 \pm 7.07; 30.0 (17-39)	27.4 \pm 4.68; 29.0 (17-35)	0.195
Parity (Mean \pm SD; Median-min-max)	2.5 \pm 1.36; 2.0 (1-5)	2.0 \pm 0.89; 2.0 (1-4)	0.297
History of rupture of membrane			
Yes	11 (55 %)	3 (15 %)	0.009
No	9 (45 %)	17 (85 %)	

* Age data distribution is normally distributed ($p>0.05$)

The distribution of parity and the history of PROM were not normally distributed ($p<0.05$)

Table 2. Analysis of the level of the leukocyte count

Levels of leukocytes	Type of rupture of membrane		P
	PROM (n = 20)	PPROM (n = 20)	
Value ($\times 10^3$)	10.885 \pm 3.885; 11.450 (4.500-17.800)	12.5350 \pm 3.119; 11.250 (7.800-18.300)	0.229
Leukocytosis			
Yes	8 (40 %)	7 (35 %)	0.3
No	12 (60 %)	13 (65 %)	

* The data is not normally distributed ($p<0.05$)

RESULTS

This research was conducted from September 2019 to January 2020, which obtained 40 research subjects consisting of 20 subjects in each PROM and PPROM.

Based on Table 1, there was no significant difference in the age variable (29.9 ± 7.07 vs 27.4 ± 4.68) and parity (2.5 ± 1.36 vs 2.0 ± 0.89) in the PROM groups at term and preterm ($p>0.05$). In this study, patients with a history of PROM were significantly different ($p<0.05$).

Differences in the levels of maternal leukocyte counts in PROM and PPROM

Table 2 shows PROM group there were 8 (40%) leukocytosis and 12 (60%) normal, while in the PPROM group obtained 7 (35%) leukocytosis and 13 (65%) normal. There was no significant difference between the leukocyte count levels in PROM and PPROM ($p=0.229$).

DISCUSSION

8 Premature rupture of membranes (PROM) is a rupture of the amniotic membrane prior to labor. Premature rupture of membranes can occur at or after 37 weeks' gestation (PROM or premature rupture of membranes), before 37 weeks' gestation (PROM or preterm premature rupture of membranes).¹

The etiology of PROM is still uncertain. A history of PROM in previous pregnancy, genital tract infection, antepartum hemorrhage, and smoking have a very close association with the incidence of PROM.⁹ In this study, the history of PROM was observed as much as 55% in PROM and 15% in PPROM. Research in Ethiopia stated that a previous history of PROM was a risk factor for the occurrence of PROM by 33.8%. The history of PROM is one of the risk factors for the incidence of PROM due to the presence of untreated/asymptomatic genital and urinary tract infections, short cervical length, as well as other obstetric problems underlying the cause of PROM.¹⁰

Leukocytosis is a sign of clinical chorioamnionitis or intraamniotic infection (IAI) or Triple I (Intrauterine infection or inflammation or both) which can occur in 40-70% of patients with PROM.⁷ In PPRM with maternal leukocytosis and managed as expectative management, poor infant outcomes can happen in the form of neurodevelopmental disorders during the first 2 years of life.⁸ Maternal leukocytosis in PROM can affect neonatal outcomes such as acute respiratory failure, sepsis, and death.¹¹ In this study, PROM obtained at term 8 (40%) leukocytosis and 12 (60%) normal, whereas in preterm PROM obtained 7 (35%) leukocytosis and 13 (65%) normal.

In this study, there was no significant difference in the levels of maternal leukocyte counts in PROM and PPRM ($p = 0.229$). This is the same results as the research conducted at Dr. H. Abdul Moeloek General Hospital at Lampung in 2018, which stated that there was no significant difference between the level of the patient's leukocyte count in PROM and PPRM.¹² Study in Czech Republic suggests that maternal leukocyte count levels are poor predictors of definitive diagnosis of IAI and microbial invasion of the amniotic cavity (MIAC), but can be used as a clinical sign of chorioamnionitis or IAI.¹³

During this study, there were no complications from venous blood sampling for routine blood laboratories. Examination of the maternal leukocyte count can be performed in PROM patients to determine the presence of leukocytosis as a clinical sign of chorioamnionitis.

The limitation of this study was that there was no assessment of other risk factors for the incidence of PROM in the form of body mass index data, the effect of genital infections, cervical length, and previous history of polyhydramnios. This study also did not have a limit on PROM occurrence for the first time, so that it can affect the level of leukocyte count in PROM and PPRM.

CONCLUSION

Statistically, there was no significant difference in leukocyte levels in PROM and PPRM. Examination of the maternal leukocyte count can be performed in PROM patients to determine the presence of leukocytosis as a clinical sign of chorioamnionitis.

CONFLICT OF INTEREST

The author declares there is no conflict of interest regarding publication of this study.

ACKNOWLEDGMENTS

We are grateful to all the participants. We also present our greatest gratitude to Dr. Kariadi General Hospital Medical Center Semarang and Kartini General Hospital Jepara as they helped to facilitate the study.

AUTHOR'S CONTRIBUTION

All of the authors developed the plan and design of the study together. BAP was the main investigator, he collected the data and conducted the analysis and first draft of the writing, as well as contributed in the editing and the approval of the final manuscript. PSW and JD were contributed in the editing and the approval of the final manuscript.

FUNDING

This study was self-funded by the corresponding author. Therefore, there isn't any conflict of interest presented to this study.

ETHICAL CONSIDERATION

This study has been approved by the Ethical Committee Faculty of Medicine Universitas Diponegoro/Dr. Kariadi Hospital, Semarang, Indonesia. All study protocol in accordance with Helsinki declaration of human rights.

REFERENCES

1. Maternal HKEM. National guide of medical service: premature rupture of membrane. Jakarta: Indonesian Society of Obstetric and Gynecologists; 2016.
2. Practice Bulletin No. 160: Premature Rupture of Membranes. *Obstet Gynecol*. 2016;127(1):e39-51.
3. Wulandari VE, Pramono BA. Factors associated with preterm labour in Dr. Kariadi Hospital Semarang. *JKD*. 2016;5(2):12-19.
4. Kacerovsky M, Musilova I, Khatibi A, Skogstrand K, Hougaard DM, Tambor V, et al. Intraamniotic inflammatory response to bacteria: analysis of multiple amniotic fluid proteins in women with preterm prelabor rupture of membranes. *J Matern Fetal Neonatal Med*. 2012;25(10):2014-9.
5. Tzur T, Weintraub AY, Sergienko R, Sheiner E. Can leukocyte count during the first trimester of pregnancy predict later gestational complications?. *Arch Gynecol Obstet*. 2013;287(3):421-427.
6. Tchirikov M, Schlabritz-Loutsevitch N, Maher J, et al. Mid-trimester preterm premature rupture of membranes (PPROM): etiology, diagnosis, classification, international recommendations of treatment options and outcome. *J Perinat Med*. 2018;46(5):465-488.
7. Peng CC, Chang JH, Lin HY, Cheng PJ, Su BH. Intrauterine inflammation, infection, or both (Triple I): A new concept for chorioamnionitis. *Pediatr Neonatol*. 2018;59(3):231-237.

8. Pasquier JC, Picaud JC, Rabilloud M, et al. Maternal leukocytosis after preterm premature rupture of membranes and infant neurodevelopmental outcome: a prospective, population-based study (Décrire l'ouverture des Membranes Inopinée le Nouveau-né et l'Organisation des Soins [DOMINOS] Study). *J Obstet Gynaecol Can.* 2007;29(1):20-26.
9. Greene MF, Creasy RK, Resnik R, Iams JD, Lockwood CJ, Moore T. *Creasy and Resnik's Maternal-Fetal Medicine: Principles and Practice E-Book*: Elsevier Health Sciences; 2008.
10. Assefa AN, Berhe H, Girma F, et al. Risk factors of premature rupture of membranes in public hospitals at Mekelle city, Tigray, a case-control study. *BMC Pregnancy Childbirth* 2018;18:386-92.
11. Pandey, Deeksha B, Saurav V, Vidhi VS, Binu P, Muralidhar K, Pratap L. Leukocyte count on admission as a predictor of clinical outcome in the expectant management for women with preterm prelabor rupture of membranes. *Sri Lanka Journal of Obstetrics and Gynaecology.* 2012;33:1-7.
12. Ratu N. Differences in leukocyte count in patient with premature rupture of membrane in Dr. H. Abdul Moeloek Hospital, Lampung. 2019. FK Universitas Lampung.
13. Musilova I, Pliskova L, Gerychova R, et al. Maternal white blood cell count cannot identify the presence of microbial invasion of the amniotic cavity or intra-amniotic inflammation in women with preterm prelabor rupture of membranes. *PLoS One.* 2017;12(12):e0189394.



This work is licensed under a Creative Commons Attribution

Differences of maternal leukocyte count in premature rupture of membranes and preterm premature rupture of membranes

ORIGINALITY REPORT

16%

SIMILARITY INDEX

13%

INTERNET SOURCES

11%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Udayana University Student Paper	3%
2	bmcmethics.biomedcentral.com Internet Source	2%
3	repository.usu.ac.id Internet Source	1%
4	Nicole O Alers, Graham Jenkin, Suzanne L Miller, Euan M Wallace. "Antenatal melatonin as an antioxidant in human pregnancies complicated by fetal growth restriction—a phase I pilot clinical trial: study protocol", BMJ Open, 2013 Publication	1%
5	Submitted to Universiti Kebangsaan Malaysia Student Paper	1%
6	hdl.handle.net Internet Source	1%
7	inabj.org Internet Source	1%

8	Submitted to St George's Hospital Medical School	1 %
Student Paper		
9	www.ncbi.nlm.nih.gov	1 %
Internet Source		
10	Céline Petit, Philippe Deruelle, Hélène Behal, Thameur Rakza et al. "Preterm premature rupture of membranes: Which criteria contraindicate home care management?", <i>Acta Obstetrica et Gynecologica Scandinavica</i> , 2018	1 %
Publication		
11	"The First China-Indonesia Joint International Symposium on Hepatobiliary Medicine and Surgery", <i>Hepatology International</i> , 2008	<1 %
Publication		
12	Christopher E. Ramsden, Maria Makrides, Zhi-Xin Yuan, Mark S. Horowitz et al. "Plasma oxylipins and unesterified precursor fatty acids are altered by DHA supplementation in pregnancy: Can they help predict risk of preterm birth?", <i>Prostaglandins, Leukotrienes and Essential Fatty Acids</i> , 2020	<1 %
Publication		
13	Ivana Musilova, Martina Kolackova, Ctirad Andrys, Marcela Drahosova et al. "Nicotinamide phosphoribosyltransferase and	<1 %

intra-amniotic inflammation in preterm
prelabor rupture of membranes", The Journal
of Maternal-Fetal & Neonatal Medicine, 2019

Publication

14

dspace.iua.edu.sd

Internet Source

<1 %

15

pt.scribd.com

Internet Source

<1 %

16

repository.unair.ac.id

Internet Source

<1 %

17

Eeva-Marja Rutanen, Tytti H. Kärkkäinen,
Johanna Lehtovirta, Jukka T. Uotila, Marianne
K. Hinkula, Anna-Liisa Hartikainen. "Evaluation
of a rapid strip test for insulin-like growth
factor binding protein-1 in the diagnosis of
ruptured fetal membranes", Clinica Chimica
Acta, 1996

Publication

<1 %

18

link.springer.com

Internet Source

<1 %

19

obgyn.pericles-prod.literatumonline.com

Internet Source

<1 %

20

doaj.org

Internet Source

<1 %

21

Başak Kaya, Uğur Turhan, Salim Sezer, Serdar
Kaya, İsmail Dağ, Ahmet Tayyar. "Maternal

<1 %

serum galectin-1 and galectin-3 levels in pregnancies complicated with preterm prelabor rupture of membranes", The Journal of Maternal-Fetal & Neonatal Medicine, 2019

Publication

22

Nuray İflazoğlu, Hasan Eroğlu, Harun Egemen Tolunay, Aykan Yücel. "Comparison of the maternal serum endocan levels in preterm premature rupture of membrane and normal pregnancy", Journal of Obstetrics and Gynaecology Research, 2021

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On