

anxiety in prisoner

by Nur Setiawati Dewi

Submission date: 14-Sep-2022 06:35PM (UTC+0700)

Submission ID: 1899566586

File name: anxiety_in_prisoner.pdf (168.42K)

Word count: 4203

Character count: 23235



Anxiety in prisoners who are currently serving a prison term: A descriptive study in a class IIA women's prison in Semarang

Maria Yoanita Bina^{1*}; Megah Andriany²; Nur Setiawati Dewi³

^{1, 2, 3} Universitas Diponegoro Semarang

ARTICLE INFO

Article history:

Received March 05, 2020
Accepted May 12, 2020
Published December 05, 2020

Keyword:

Anxiety
Women's correctional facility
Female inmates

ABSTRACT

Detention can cause problems and stressors for prisoners such as anxiety. Anxiety in female prisoners has an impact on psychological and emotional stress that causes violence in prisons. However, research related to anxiety in female prisoners is still limited to anxiety before going home. Therefore, researchers are interested in researching to determine the level of anxiety in prisoners who are serving a period of detention. The purpose of this study was to determine the level of anxiety in female prisoners who are serving a period of detention. The research design used in this study was a critical study to see a picture of the anxiety level of female prisoners. The sample in this study was 128 people who were selected using a purposive sampling technique. The results showed that the highest prisoners' anxiety level was in the moderate category as much as 44.5 percent and the lowest was in the panic category as much as 5.5 percent. Female prisoner anxiety arises because of negative emotions due to problems faced, stressors, feelings of failure, and feelings of insecurity. Most of the anxiety levels of female prisoners in prison are in the medium category. Therefore, nurses need to develop research related to interventions to overcome the anxiety of female prisoners in prison.

11

This open access article is under the CC-BY-SA license.



12 Kecemasan pada Warga Binaan Pemasyarakatan yang sedang menjalani masa tahanan: Studi deskriptif di lapas perempuan kelas IIA Semarang

ABSTRAK

Penahanan dapat menimbulkan masalah dan stressor bagi warga binaan pemasyarakatan (WBP) seperti kecemasan. Kecemasan pada WBP perempuan berdampak pada tekanan psikis dan emosional yang menimbulkan kekerasan dalam lapas. Namun, penelitian terkait kecemasan pada WBP perempuan masih terbatas pada kecemasan menjelang pulang. Oleh karena itu, peneliti tertarik untuk melakukan penelitian untuk mengetahui tingkat kecemasan pada WBP yang sedang menjalani masa tahanan. Tujuan dari penelitian ini adalah untuk mengetahui tingkat kecemasan pada WBP perempuan yang sedang menjalani masa tahanan. Desain penelitian yang digunakan dalam penelitian ini adalah penelitian kualitatif untuk melihat gambaran tingkatan kecemasan WBP perempuan. Sampel dalam penelitian ini sebesar 128 orang yang dipilih dengan menggunakan teknik purposive sampling. Hasil penelitian menunjukkan tingkat kecemasan WBP tertinggi berada pada kategori sedang sebanyak 44.5% dan terendah berada pada kategori panik sebanyak 5.5%. Kecemasan WBP perempuan muncul karena adanya emosi negatif karena adanya permasalahan yang dihadapi, adanya stressor, merasakan kegagalan, dan adanya perasaan tidak aman. Tingkat kecemasan WBP perempuan di Lapas sebagian besar berada pada kategori sedang. Oleh Karena itu, perawat perlu

Kata kunci:

Kecemasan
Lapas perempuan
Warga Binaan Perempuan

*) corresponding author

Departemen Ilmu Keperawatan Universitas
Diponegoro Semarang
Jl. Prof. Soedarto, SH, Tembalang,
Semarang/telp./fax: (024) 76408919/ (024)
76486849

Email: ns.yoanitabina@gmail.com

23
DOI: <https://doi.org/10.30604/jika.v5i2.294>

6

mengembangkan riset terkait intervensi untuk mengatasi kecemasan WBP perempuan di Lapas.

1
This open access article is under the [CC-BY-SA](https://creativecommons.org/licenses/by-sa/4.0/) license.



Introduction

Detention can cause problems and stressors for prisoners such as mental pressure, trauma, tendencies to self-isolate, social deviation, suspicion, and anxiety (Kartono, 2014). The results of research in several countries show the high incidence of mental disorders in prisons (Arnau, 2014; Iró et al., 2012; Greenberg & Rosenheck, 2014; Hassiotis et al., 2011; Kingston, Le Mesurier, Yorston, Wardle, & Heath, 2011; Kumar et al., 2014). Female prisoners' mental health status is mostly bad as much as 55% (Geitona & Milioni, 2016). Besides, 41.6% of female prisoners have severe mental disorders (Al-Rousan, Rubenstein, Sieleni, Deol, & Wallace, 2017). The problem of mental disorders that often occur in female assisted residents is anxiety.

The results showed that female prisoners' anxiety level who experienced severe anxiety was 38% (Utari, Fitria, & Rafiyah, 2012). Other studies also show that 53.83% of female prisoners experience moderate anxiety (Mandira, Marjohan, & Fernandes, 2019). Besides, the anxiety rate for female prisoners is greater (21.3%) compared to male prisoners (6.6%) (van den Brink, Harte, & Denzel, 2018). The high rate of anxiety is often associated with several factors.

Anxiety in female prisoners is influenced by the ability to adjust to a new place (Crewe, 2009; Værøy, 2011), family support and social support from WBP (Salim, Komariah, & Fitria, 2016), and personal relationships with family (Paul Swift, Eva Cyhlarova & O'Sullivan, 2014). Besides, women in prison have a great chance of experiencing economic difficulties and most of them are caring for their children alone, causing conflicts such as anxiety (Ferszt, Miller, Hickey, Maull, & Crisp, 2015). Anxiety has an adverse impact on female prisoners.

Anxiety in female prisoners has an impact on psychological and emotional stress that causes violence in prisons in the form of physical and sexual violence such as beatings, slaps, and being threatened with weapons (Bensimon, Einat, & Gilboa, 2015; Eleanor Taylor-Nicholson & Barry Krisberg, 2013). However, research related to anxiety in female prisoners is still limited to anxiety before going home (Salim et al., 2016; Utari et al., 2012) and mental disorders and has not been specific to female prisoners (Andreoli et al., 2014; Badru, Ogunlesi, Ogunwale, Abdulmalik, & Yusuf, 2018; Dadi, Dachew, Kisi, Yigzaw, & Azale, 2013; Malik, Singh, Beniwal, & Kumar, 2019). Therefore, the researcher is interested in conducting a study to determine the level of anxiety in prisoners who are serving a prison term, so the aim of this study is to determine the level of anxiety in female prisoners who are currently serving a prison term.

2 Method

The research design used in this study was a critical study to see a picture of female prisoners' anxiety level. This research was conducted at the Class IIA Women's Prison in Semarang. The population of female prisoners in this study was 302 people. The sample in this study was 128 people selected using a purposive sampling technique with inclusion criteria aged 25-55 years and a prison term of more than 1 year and exclusion criteria for female prisoners who were in isolation cells and who suffered from chronic disease.

Data were collected using a questionnaire. The questionnaire for the respondents' characteristics consisted of age, education, marital status, and period of imprisonment. The anxiety questionnaire uses Spielberger's The State Anxiety Inventory (STAI) which consists of 20 statement items (Spielberger, Gorsuch, Lushene, Vagg, & Jacobs, 1983). The State anxiety scale is designed to be sensitive to the situation when the measurement is made, the score on this scale can be influenced by the emotional state that is generally felt (Johnson & Spielberger, 1968). The STAI questionnaire's validity and reliability were tested at the Class IIA Women's Prison in Semarang. Respondents who have been used to test the validity and reliability will not be used again in this study. The results of the validity test on 30 respondents showed rcount of 0.430-0.790 and the reliability test showed Cronbach's alpha of 0.892. Data analysis in this study used univariate analysis in the form of frequency and distribution to describe the level of anxiety of female prisoners.

Results and Discussion

The results showed that most female prisoners were in early adulthood amounting to 66.4% (Table 1). The highest education level of female prisoners was Senior High School as much as 49.2% and the lowest is not in school as much as 1.6% (Table 1). The highest female prisoner status was married 48.4% and the lowest was single 22.7% (Table 1). The highest prison term for female prisoners was 1-5 years as much as 42.3% and the lowest in over 5 years as much as 10.95% (Table 1).

All humans can experience anxiety at various age levels. However, anxiety is more common in adulthood. The results showed that most of the respondents were in early adulthood. Medley argues that early adulthood is in the age range of 22-34 years (Medley, 1980). Early adulthood is a time when a person adapts to a new style of life and new community expectations so that prisoners in early adulthood

5
 need the ability to be able to adapt to the various changes that occur. Changes that occur in early adulthood include work by engaging in adult duties to confirm identity, marriage or partnership in intimacy duties, and parenting to ensure the next generation's welfare (Srivastava, John, Gosling, & Potter, 2003). Female prisoners who cannot adapt to change will experience anxiety (Medley, 1980; Salim et al., 2016). The results showed that most prisoners experienced anxiety in early adulthood (Osasona & Koleoso, 2015., Harlianty, R., & Paramastri, I. 2020). Therefore, it can be concluded that female prisoners in early adulthood are prone to experience anxiety caused by changes in society's roles and demands.

Table 1
Characteristics of female prisoners in Class IIA Women's Prison in Semarang (n = 128)

Variable	Frequency (n)	Percentage (%)
Age		
Early adulthood	85	66.4
Late adulthood	43	33.6
Education level		
Uneducated	2	1.6
Elementary School	15	11.7
Junior High School	24	18.8
Senior High School	63	49.2
College	24	18.8
Marital Statues		
Unmarried	29	22.7
Married	62	48.4
Widow	37	28.9
Criminal period		
1-5 years	55	42.3
6-10 years	41	32
11-15 years	18	14.1
>15 years	14	10.95

The results showed that most female prisoners had senior high school education levels. The results of this study are supported by previous research which shows that most of the prisoners have senior high school education (58.8%) (Mandira et al., 2019). The education level of female prisoners will affect the level of their understanding so that it helps them to understand the information obtained and use it as an alternative problem solving so as to form an adaptive coping mechanism in overcoming various problems in prison such as anxiety, feelings of guilt and regret, disappointment with family attitudes, and feeling that life is not meaningful (Hayati, 2015; Mutoharoh, 2009; Putri, Ira, & Hilma, 2014). Therefore, female prisoners who have the elementary school education level have the opportunity to understand the use of a positive defense system to solve problems.

The results showed that most of the marital status of female prisoners was already married at 48.4% and 28.9% were widows. Marital status is one of the stressors for female prisoners due to limitations in maintaining family integrity, the role of mother in caring for children, and conflicts in marriage with husbands (Aini, 2015). In addition, most prisoners who are serving a period of detention carry out their role as sole caregivers so that they become stressors for female prisoners because they think about their children's future (Ferszt et al., 2015). Therefore, the risk of marital status creates anxiety for female prisoners.

The results showed that most female prisoners in prison have a prison term of 1-5 years. The results of research in the UK show that long incarceration with little mental stimulation contributes to deteriorating mental health and is a cause of anger, frustration, and anxiety (Martha & Annatagia, 2014; Nurse, Woodcock, & Ormsby, 2003). Mental health services in prisons are so limited that they cannot ensure the mental health of all prisoners. Therefore, female prisoners who have a long prison term are at risk of experiencing anxiety.

Table. 2
Anxiety Level of Female Prisoners in Class IIA Prison in Semarang (n = 128)

Anxiety Level	Frequency (n)	Percentage (%)
Mild anxiety	36	28.1
Moderate anxiety	57	44.5
Sever anxiety	28	21.9
Panic	7	5.5

21
 22
 The results showed that the highest female prisoners' anxiety level was in the moderate category as much as 44.5% and the lowest was in the panic category as much as 5.5% (Table 1). Most of the female prisoners who served a period of detention had an anxiety level in the moderate anxiety category. Most of the female prisoners who were in prison had moderate levels of anxiety. The results of this study are supported by research conducted by Mandira which shows that female prisoners experience anxiety in the moderate category of 53.8% (Mandira et al., 2019). These results indicate that female prisoners are not only feeling anxiety before the time of release

Prison conditions play an important role in prisoners' anxiety because prisoners have to adapt to new places and roles (Værøy, 2011). Class IIA Women's Prison in Semarang has mandatory regulations and activities that must be followed so that WBP feels that their freedom is very limited. This is what makes it difficult for WBP to adapt. The prison environment has a deep and negative impact on the physical, mental, and emotional well-being of prisoners. Losing freedom, especially limitation in making decisions to carry out the role, causes feelings of frustration, anxiety, and stress. In addition, excess capacity creates female prisoners' problems because they cannot maintain their privacy and creates a situation that creates hostility (Bradford, 2006; Goomany & Dickinson, 2015).

The results showed that female prisoners' anxiety arises because of negative emotions due to problems faced, stressors, feelings of failure, and feelings of insecurity (Martha & Annatagia, 2014). This is in line with female prisoners' statement that the anxiety that is felt arises because of negative feelings caused by feelings of depression, insecurity, and feelings of distrust of oneself. The results showed that they experienced psychological pressure due to forced separation (Harner & Riley, 2013). Separation from children is a stressor because prisoners cannot confirm the condition and whereabouts of their children. This was also conveyed by the female prisoners at the Class IIA Women's Prison in Semarang that the inability to ensure the development and fulfillment of children's needs is one of the causes of feelings of anxiety. In addition, close supervision during visits causes the relationship between prisoners and their families and children to become obstructed (Goomany & Dickinson, 2015; Prison Reform Trust, 2019). Other studies have shown that access to information on the family by the telephone is sometimes difficult to do because of many

requests and long queues so that the relationship with the family is cut off causing feelings of fear, worry, and triggers anxiety (Harner & Riley, 2013; Nurse et al., 2003). The female prisoners said that the time for using telephones was very limited and sometimes the call was not answered by the family so that the prisoners could not get information related to the family, this triggered feelings of anxiety. However, mental health services in prisons are very limited.

The results of the literature review show that a low of mental health services in prisons starts from screening, triage, assessment, intervention, and reintegration (Forrester, Till, Simpson, & Shaw, 2018). However, research results show that mental health services in prisons have not been carried out optimally (Birmingham, 2003). These results were supported by interviews with health workers in prisons who said that there was no special program related to mental

health in prisons so that the results of mental health screening carried out at prison were not followed up. In addition, follow-up on mental health problems can only be carried out if the prisoner arrives with a clinical diagnosis that he has a mental disorder. Therefore, it is necessary to have a mental health service program in prisons including interventions to overcome prisoner anxiety.

Conclusions and Recommendations

Most of the anxiety levels of female prisoners are in the moderate category. Therefore, nurses need to develop research related to interventions to overcome the anxiety of female prisoners.

References

- Aini, R. (2015). Pengaruh cognitive behavior therapy (CBT) terhadap penurunan derajat kecemasan pada narapidana yang akan bebas di Lembaga Pemasyarakatan Wanita kelas II A Bandung. *Scientica*, 2(1), 58–71. <https://doi.org/10.29313/SCIENTICA.V2I1.2312>
- Al-Rousan, T., Rubenstein, L., Sieleni, B., Deol, H., & Wallace, R. B. (2017). Inside the nation's largest mental health institution: A prevalence study in a state prison system. *BMC Public Health*, 17(1), 342. <https://doi.org/10.1186/s12889-017-4257-0>
- Andreoli, S. B., Dos Santos, M. M., Quintana, M. I., Ribeiro, W. S., Blay, S. L., Taborda, J. G. V., & De Jesus Mari, J. (2014). Prevalence of mental disorders among prisoners in the state of Sao Paulo, Brazil. *PLoS ONE*, 9(2). <https://doi.org/10.1371/journal.pone.0088836>
- Amáu-Peiró, F., García-Guerrero, J., Herrero-Matías, A., Castellano-Cervera, J. C., Vera-Remartínez, E. J., Jorge-Vidal, V., ... Iranzo-Tatay, C. (2012). Description of the psychiatric unit in prisons in the autonomous community of Valencia. *Revista Espanola de Sanidad Penitenciaria*, 14(2), 50–60. <https://doi.org/10.1590/S1575-06202012000200003>
- Badru, O. A., Ogunlesi, A. O., Ogunwale, A., Abdulmalik J. O., & Yusuf, O. B. (2018). Prevalence of generalized anxiety disorder and major depression among correctional officers in a Nigerian prison. *Journal of Forensic Psychiatry and Psychology*, 29(4), 509–526. <https://doi.org/10.1080/14789949.2017.1421250>
- Bensimon, M., Einat, T., & Gilboa, A. (2015). The impact of relaxing music on prisoner's levels of anxiety and anger. *International Journal of Offender Therapy and Comparative Criminology*, 59(4), 406–423. <https://doi.org/10.1177/0306624X13511587>
- Birmingham, L. (2003). The mental health of prisoners. *Advances in Psychiatric Treatment*, 9(3), 191–201. <https://doi.org/10.1192/apt.9.3.191>
- Bradford, A. R. (2006). *An examination of the prison environment: An analysis of Inmate concerns across eight environmental dimensions*. East Tennessee State University, Tennessee. Retrieved from <http://dc.etsu.edu/etd/2216>
- Crewe, B. (2009). The prisoner society power, adaptation and social life in an English prison. <https://doi.org/10.1093/acprof:oso/9780199577965.001.0001>
- Dadi, A. F., Dachew, B. A., Kisi, T., Yigzaw, N., & Azale, T. (2016). Anxiety and associated factors among prisoners in North West of Amhara Regional State, Ethiopia. *BMC Psychiatry*, 16(1), 1–6. <https://doi.org/10.1186/s12888-016-0792-y>
- Eleanor Taylor-Nicholson, & Barry Krisberg. (2013, February 6). Contagion of violence. Retrieved April 27, 2020, from <https://www.ncbi.nlm.nih.gov/books/NBK207242/>
- Ferszt, G., Miller, R., Hickey, J., Maull, F., & Crisp, K. (2015). The impact of a mindfulness based program on perceived stress, anxiety, depression and sleep of incarcerated women. *International Journal of Environmental Research and Public Health*, 12(9), 11594–11607. <https://doi.org/10.3390/ijerph120911594>
- Forrester, A., Till, A., Simpson, A., & Shaw, J. (2018). Mental illness and the provision of mental health services in prisons. *British Medical Bulletin*, 127(1), 101–109. <https://doi.org/10.1093/bmb/ldy027>
- Geitona, M., & Milioni, S.-O. (2016). Health status and access to health services of female prisoners in Greece: a cross-sectional survey. *BMC Health Services Research*, 16(1), 243. <https://doi.org/10.1186/s12913-016-1506-3>
- Goomany, A., & Dickinson, T. (2015, August 1). The influence of prison climate on the mental health of adult prisoners: A literature review. *Journal of Psychiatric and Mental Health Nursing*. Blackwell Publishing Ltd. <https://doi.org/10.1111/jpm.12231>
- Greenberg, G. A., & Rosenheck, R. A. (2014). Psychiatric correlates of past incarceration in the national co-morbidity study replication. *Criminal Behaviour and Mental Health*, 24(1), 18–35. <https://doi.org/10.1002/cbm.1875>
- Hadi W, R. (2013). Pengaruh intervensi musik gamelan terhadap depresi pada lansia di panti wreda harapan ibu, Semarang. *Keperawatan Komunitas*, 1, 135–140.
- Harlianty, R., & Paramastri, I. (2020). The role of religious coping as a moderator of the relationship between psychological burden and quality of life among caregiver of women with breast cancer. *Journal of Psychological Perspective*, 2(2), 71–78. Retrieved from <https://ukinstitute.org/journals/jopp/article/view/2206>
- Hamer, H. M., & Riley, S. (2013). The impact of incarceration on women's mental health: Responses from women in a maximum-security prison. *Qualitative Health Research*, 23(1), 26–42. <https://doi.org/10.1177/1049732312461452>

- Hassiotis, A., Gazizova, D., Akinlonu, L., Bebbington, P., Meltzer, H., & Strydom, A. (2011). Psychiatric morbidity in prisoners with intellectual disabilities: analysis of prison survey data for England and Wales. *British Journal of Psychiatry*, 199(02), 156–157. <https://doi.org/10.1192/bjp.bp.110.088039>
- Hayati, N. (2015). *Penanganan terhadap problematika psikologis kehidupan rumah tangga warga binaan Lapas Wanita kelas II A SEMARANG (Prespektif bimbingan konseling keluarga Islami)*. Universitas Negeri Walisongo, Semarang. Retrieved from <http://eprints.walisongo.ac.id/4937/1/101111080.pdf>
- Johnson, D. T., & Spielberger, C. D. (1968). The effects of relaxation training and the passage of time on measures of state- and trait-anxiety. *Journal of Clinical Psychology*, 24(1), 20–23. [https://doi.org/10.1002/1097-4679\(196801\)24:1<20::AID-JCLP2270240105>3.0.CO;2-9](https://doi.org/10.1002/1097-4679(196801)24:1<20::AID-JCLP2270240105>3.0.CO;2-9)
- Kartono, K. (2014). *Patologi sosial*. Jakarta: Raja Grafindo.
- Kingston, P., Le Mesurier, N., Yorston, G., Wardle, S., & Heath, L. (2011). Psychiatric morbidity in older prisoners: unrecognized and undertreated. *International Psychogeriatrics*, 23(08), 1354–1360. <https://doi.org/10.1017/S1041610211000378>
- Kumar, D., Viswanath, B., Sebastian, A., Holla, B., Konduru, R., Chandrashekar, C. R., & Math, S. B. (2014). Profile of male forensic psychiatric inpatients in South India. *International Journal of Social Psychiatry*, 60(1), 55–62. <https://doi.org/10.1177/0020764012461334>
- Malik, J. S., Singh, P., Beniwal, M., & Kumar, T. (2019). Prevalence of depression, anxiety and stress among jail inmates. *International Journal of Community Medicine And Public Health*, 6(3), 1306. <https://doi.org/10.18203/2394-6040.ijcmph20190631>
- Mandira, T. M., Marjohan, & Fernandes, F. (2019). Gambaran kecemasan narapidana wanita di Sumatera Barat, 13(3), 213–217. <https://doi.org/10.33024/hjk.v13i3.1665>
- Martha, S. I., & Annatagia, L. (2014). Hubungan kecerdasan emosi dengan kecemasan menghadapi masa pembebasan pada narapidana. *Jurnal Psikologi Integratif*, 2(2). <https://doi.org/10.14421/jpsi.2014.2x>
- Medley, M. L. (1980). Life satisfaction across four stages of adult life. *The International Journal of Aging and Human Development*, 11(3), 193–209. <https://doi.org/10.2190/D41G-A1JQ-8850-GYDV>
- Mutoharoh, I. (2009). *Faktor-faktor yang berhubungan dengan mekanisme koping klien gagal ginjal kronik yang menjalani terapi hemodialisis di RSUP Fatmawati*. Universitas Islam Negeri Syarif Hidayatullah Jakarta, Jakarta. Retrieved from http://repository.uinjkt.ac.id/dspace/bitstream/123456789/2340/1/ITOH_MUTOHAROH-FKIK.PDF
- Nurse, J., Woodcock, P., & Ormsby, J. (2003). Influence of environmental factors on mental health within prisons: focus group study. *BMJ (Clinical Research Ed.)*, 327(7413), 480. <https://doi.org/10.1136/bmj.327.7413.480>
- Oasona, S. O., & Koleoso, O. N. (2015). Prevalence and correlates of depression and anxiety disorder in a sample of inmates in a Nigerian prison. *The International Journal of Psychiatry in Medicine*, 50(2), 203–218. <https://doi.org/10.1177/0091217415605038>
- Paul Swift, Eva Cyhlarova, I., & O'Sullivan, G. dan C. (2014). Living with anxiety. Retrieved September 7, 2018, from <https://www.mentalhealth.org.nz/assets/A-Z/Downloads/Living-with-anxiety-report-MHF-UK-2014.pdf>
- Prison Reform Trust. (2019). *Prison: the facts*. Retrieved from <http://www.statcan.gc.ca/pub/11-630-x/11-630-x2015001-eng.htm>
- Putri, D. E., Ira, E., & Hilma, A. (2014). Hubungan dukungan sosial dengan tingkat kecemasan narapidana di lembaga pemasyarakatan kelas IIA Muaro Padang. *Keperawatan*, 10(1). <https://doi.org/10.25077/njk.10.2.118-135.2014>
- Salim, shalha U., Komariah, M., & Fitria, N. (2016). Gambaran faktor yang mempengaruhi kecemasan WBP menjelang bebas di LP Wanita IIA Bandung, 4(1), 32–42. <https://doi.org/10.31311/V4I1.402>
- Spielberger, C. D., Gorsuch, R. L., Lushene, R., Vagg, P. R., & Jacobs, G. A. (1983). *Manual for the State-Trait Anxiety Inventory*. <https://doi.org/10.1007/978-1-4419-9893-4>
- Srivastava, S., John, O. P., Gosling, S. D., & Potter, J. (2003). Development of personality in early and middle adulthood: Set like plaster or persistent change? *Journal of Personality and Social Psychology*, 84(5), 1041–1053. <https://doi.org/https://doi.org/10.1037/0022-3514.84.5.1041>
- Utari, D. I., Fitria, N., & Rafiyah, I. (2012). Gambaran tingkat kecemasan pada warga binaan wanita menjelang bebas di lembaga pemasyarakatan wanita Kelas II A Bandung, 1(1), 33. Retrieved from <http://journals.unpad.ac.id/index.php/ejournal/article/view/777>
- Værøy, H. (2011). Depression, anxiety, and history of substance abuse among Norwegian inmates in preventive detention: Reasons to worry? *BMC Psychiatry*, 11(1), 40. <https://doi.org/10.1186/1471-244X-11-40>
- van den Brink, C., Harte, J. M., & Denzel, A. D. (2018). Men and women with borderline personality disorder resident in Dutch special psychiatric units in prisons: A descriptive and comparative study. *Criminal Behaviour and Mental Health: CBMH*, 28(4), 324–334. <https://doi.org/10.1002/cbm.2084>

anxiety in prisoner

ORIGINALITY REPORT

16%

SIMILARITY INDEX

13%

INTERNET SOURCES

11%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	storage-imelda.s3.ap-southeast-1.amazonaws.com Internet Source	2%
2	simdos.unud.ac.id Internet Source	2%
3	journal.walisongo.ac.id Internet Source	2%
4	Tria Monja Mandira, Marjohan Marjohan, Feri Fernandes. "Gambaran kecemasan narapidana wanita di Sumatera Barat", <i>Holistik Jurnal Kesehatan</i> , 2019 Publication	1%
5	www.ssoar.info Internet Source	1%
6	Submitted to Universitas Diponegoro Student Paper	1%
7	dc.etsu.edu Internet Source	1%

8	Norman, Ian, Rylie, Iain. "The Art and Science of Mental Health Nursing: Principles and Practice", The Art and Science of Mental Health Nursing: Principles and Practice, 2018	1 %
Publication		
9	Garuda.Kemdikbud.Go.Id	<1 %
Internet Source		
10	Iwan Shalahuddin, Indra Maulana, Sandra Pebrianti, Theresia Eriyani. "Blood Sugar Levels Regulation in Diabetes Mellitus Type 2 Patients Through Diet Management", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2022	<1 %
Publication		
11	R Rokhmat, Achmad Zulfa Juniarto, Chandra Bagus Ropyanto. "Descriptive study of anxiety among chronic kidney disease patients underwent hemodialysis: Pilot study", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2020	<1 %
Publication		
12	eprints.walisongo.ac.id	<1 %
Internet Source		
13	repository.unisba.ac.id:8080	<1 %
Internet Source		
14	biblio.ugent.be	<1 %
Internet Source		
15	worldwidescience.org	<1 %
Internet Source		

<1 %

16

Scholar.ufs.ac.za

Internet Source

<1 %

17

docksci.com

Internet Source

<1 %

18

ejournal.stikesmuhgombong.ac.id

Internet Source

<1 %

19

jurnalmadanimedika.ac.id

Internet Source

<1 %

20

www.scielo.org.za

Internet Source

<1 %

21

David Onchonga, Enoch Ngetich, Wilbroda Makunda, Pius Wainaina, Diana Wangeshi, Prémusz viktoria. "Anxiety and Depression due to 2019 SARS-CoV-2 among Frontier Healthcare Workers in Kenya", Heliyon, 2021

Publication

<1 %

22

garuda.ristekdikti.go.id

Internet Source

<1 %

23

Tetti Solehati, Bhekti Imansari, Devita Madiu, Cecep Eli Kosasih, Yanti Hermayanti, Henny Suzana Mediani. "Effect of Cervical Cancer Education and Provider Recommendation for Screening: A Systematic Review", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2020

<1 %

24 digilib.unila.ac.id <1 %
Internet Source

25 ejournalmalahayati.ac.id <1 %
Internet Source

26 scholarworks.waldenu.edu <1 %
Internet Source

27 Kinga Kołodziej, Anna Kurowska, Anna Majda. "Intensity of perceived stress and control of anger, anxiety and depression of people staying in Polish penitentiary institutions", International Journal of Prisoner Health, 2021
Publication

28 Cory Gerritsen, Vito Adamo, Lindsay Fulham, Roland Jones, Stephanie Penney, Abanti Tagore, Alexander Simpson. "Evidence for the ecological validity of the Brief Jail Mental Health Screen: Positive predictive value among remanded men and women", Criminal Behaviour and Mental Health, 2022
Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On