

2024 Brazil London Atlas

M3.pdf

by Rizky Merdietio Boedi

Submission date: 19-Oct-2024 06:59PM (UTC+0700)

Submission ID: 2490166189

File name: 2024_Brazil_London_Atlas_M3.pdf (850.27K)

Word count: 5323

Character count: 25055

Accuracy of the London Atlas to estimate the age of legal majority in a sample of the Amazon Region

Flávia Afonso ¹, Debora Moreira ², Rizky Boedi ³, Izabella Goetten ¹, Ernesto Lourenço-Junior ⁴, Vanessa Sartori ⁵, João-Paulo de Carli ⁵, Ademir Franco ^{1,6}

¹ Division of Forensic Dentistry, Faculdade São Leopoldo Mandic, Brazil

² Division of Oral Radiology, Faculdade São Leopoldo Mandic, Brazil

³ Department of Dentistry, Faculty of Medicine, Universitas Diponegoro, Indonesia

⁴ Medical and Odontological Imaging Clinic (CIMO), Amazonas, Brazil

⁵ Department of Dentistry, Federal University of Passo Fundo, Brazil

⁶ Department of Therapeutic Dentistry, Institute of Dentistry, Sechenov University, Moscow, Russia

Correspondence:

Division of Forensic Dentistry

Faculdade São Leopoldo Mandic

R. Dr. José Rocha Junqueira 13

Swift, Campinas, São Paulo, Brazil

Postal code: 13.045-755

ademirjunior@slmandic.edu.br

Afonso F, Moreira D, Boedi R, Goetten I, Lourenço-Junior E, Sartori V, de Carli JP, Franco A. Accuracy of the London Atlas to estimate the age of legal majority in a sample of the Amazon Regio. J Clin Exp Dent. 2024;16(4):e472-9.

Received: 01/12/2023

Accepted: 26/02/2024

Article Number: 61263 <http://www.medicinaoral.com/odo/indice.htm>

© Medicina Oral S. L. C.I.F. B 96689336 - eISSN: 1989-5488

eMail: jced@jced.es

Indexed in:

Pubmed
 Pubmed Central® (PMC)
 Scopus
 DOI® System

Abstract

Background: Estimating the age of majority is a challenging task in forensic odontology, especially because the third molars are usually the only developing teeth between the ages of 16 and 21 years. The London Atlas emerged as an alternative to estimate age using dental development, eruption and deciduous root resorption as parameters. The method has performed well in young age categories, while its performance for age estimation via third molars has been disputed. The present study tested the performance of the London Atlas to estimate the age of legal majority in a sample of individuals from the Amazon Region.

Material and Methods: The sample consisted of 1.256 panoramic radiographs of women (n = 694) and men (n = 562) between 16 and 22.9 years. The method was applied to the maxillary (#28) and mandibular (#38) left third molars. For comparative purposes, the sample was divided into seven age groups: 16 | 16.99; 17 | 17.99; 18 | 18.99; 19 | 19.99; 20 | 20.99; 21 | 21.99; and 22 | 22.99 years. Chronological and estimated ages were compared respectively by means of mean absolute errors (MAE) and root mean squared errors (RMSE), as well as through Receiver Operating Characteristic (ROC) curves and their area under the curve (AUC).

Results: The MAE of the age estimates using tooth #28 was 1.76 years for females and 1.52 years for males. When the tooth #38 was used, the MAE for the females and males were 1.68 and 1.51 years, respectively. The MAE and RMSE increased in ascending order between age categories. Tooth #28 led to 74% of correct classifications around the age of legal majority, while tooth #38 reached 77%. The area under the curve was 0.75 for tooth #28 and 0.73 for tooth #38.

Conclusions: The London Atlas should be used carefully to estimate the age of legal majority and not as a single method when the age threshold is 18 years.

Key words: Age determination by teeth, forensic dentistry, forensic science, third molar.

Introduction

Testing the performance of dental age estimation methods in samples with different geographic origin has been the objective of many studies in forensic odontology (1-5). When the tests confirm the applicability and validity of a method originally developed from an international sample, they pave the way with scientific evidence to support country-specific forensic practices with a larger armamentarium. The London Atlas is an age estimation method based on human dental development and eruption timing (6). This method is known for the intuitive user-interface and for the robust scientific background. Several studies have tested the performance of the London Atlas worldwide, including in populations from Brazil (7), China (8), Korea (9), Portugal (10), South Africa (11) and Thailand (12). A recent systematic review claimed that the method has an acceptable error rate and proper accuracy for dental age estimation (13). This conclusion, however, might be disputable when it comes to the Brazilian population.

A population-specific meta-analysis ranked the best radiographic dental age estimation methods applied to Brazilian children (14). Out of 2,527 entries detected with a dedicated search string, the London Atlas has not been included amongst the top ranked methods (14). Regarding the performance of the method, previous studies have found an overall accuracy of 79.9% (15), while others have demonstrated concern especially when the method is applied solely based on third molars (7,16). Most of the performance tests that have been accomplished with the London Atlas have followed the common methodological structure of dental age estimation studies and quantified the difference between chronological and estimated ages (7,16). Moreover, all the studies performed in the Brazilian population sampled participants from the Southeastern, Central-Western and Northeastern regions of Brazil (7,13-16). Hence, the scientific literature has no study with a population from the Amazon region.

The importance of studying populations from this region relies on the broad frontier between Brazil, Colombia, Peru and Venezuela. Regions of geographic frontiers might facilitate clandestine migration, human trafficking, and exploitation, and even body concealment – situations that usually involve undocumented persons. By estimating a person's age, experts might be able to contribute to the identification process, and also help supporting the legal systems regarding Court decisions about the age of majority and sexual consent. The scientific literature about dental age estimation methods applied to populations of the Amazon region is scarce.

This study aimed to apply and test the performance of the London Atlas in a population of the Amazon region.

2

Material and methods

Study design and ethical aspects

This was an observational, analytical, cross-sectional study formed. Hence, the checklist for cross-sectional studies established by the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) was followed (17). Ethical approval was obtained from the institutional review board (protocol: 44953421.2.0000.5374).

Sample and participants

The sample consisted of 1,256 panoramic radiographs of women (n = 694) and men (n = 562) between 13 and 22.9 years. The radiographs were retrospectively collected from an existing radiological database in Manaus (capital city of the State of Amazonas, Brazil), which means that no patient was exposed to ionizing radiation for research purposes. The inclusion criteria consisted of radiographs from participants of Brazilian nationality and originally from the Amazon region, with age between 16 and 22.9 years, and with a least one panoramic radiograph stored at the radiological database. The exclusion criteria consisted of radiographs with missing maxillary (#28) and mandibular (#38) left third molars, presence of teeth #28 and #38 with extensive decay, restorations and root canal treatment; presence of visible bone lesions associated with teeth #28 and #38; surgical appliances in the mandible, deformation of maxillofacial bones, and visible third molar anomalies; poor image quality; and missing data about the date of image acquisition and patient's date of birth and sex.

Sample collection was performed between January and June, 2022. The sample size established in the current section was based on a systematic review and meta-analysis (13) that compiled dental age estimation studies with the London Atlas. Hence, we targeted a sample size (n) as close as possible to the one used by the developers of the London Atlas in a comparative study published in 2014 (18). Apart from the study published by the London Atlas' authors, most of the studies worldwide (96%) that tested the London Atlas used smaller samples (13). This is the first study with a sample from the Amazon region. The sample of panoramic radiographs was imported to a personal computer equipped with a 15" screen and Adobe Photoshop CS6™ image viewer (Adobe Inc. San Jose, CA, USA) for magnification of 100% and eventual adjustments of brightness and contrast prior to analysis.

Variables and data sources

The difference between chronological and estimated ages was the main outcome of this study. This outcome was assessed separately for females and males. Hence, sex was the first variable to be considered. The second variable consisted of the chronological age, which calculated as the difference between date of radiographic

image acquisition and data of birth of each participant. Sex and chronological age were obtained from the headings of the digital panoramic radiographs before cropping them for anonymization. The third variable was the estimated age quantified by means of London Atlas method. The comparisons between chronological and estimated age were performed for based on age categories (meaning that age as a continuous variable was converted into a categorical variable): 16-16.99 years, 17-17.99 years, 18-18.99 years, 19-19.99 years, 20-20.99 years, 21-21.99 years, and 22-22.99 years.

-Assessment of reproducibility

Following the protocol for the assessment of reproducibility previously published, the main observer revisited 100 panoramic radiographs (n = 200 third molars) of the sample, 30 days from the analysis of the full sample. The comparison between observer classification of third molar development enabled the assessment of the intra-observer agreement. For the assessment of the inter-observer reproducibility, an additional observer was recruited to assess the same 100 radiographs. The comparisons between the main and the second observers was accomplished. Intra- and inter-observer reproducibility were calculated between estimated ages using the London Atlas, and the Intraclass Correlation Coefficient (ICC) was applied.

-Statistical analyses

Descriptive assessment was applied to observer first the distribution of participants based on sex and age categories. Additionally, the descriptive approach was applied to count the number of correct classifications per sex, age category and third molar (#28, #38). Descriptive statistics consisted of measures of central tendency and dispersion (such as means and standard deviation, respectively) as well as absolute (n) and relative (%) frequencies of distribution. The difference between chronological and estimated ages was expressed as mean absolute error (MAE) and root mean square error (RMSE). Spearman correlation test was applied to assess the correlation between third molar development and chronological age. T-tests were performed to verify the statistical significance of the correlation tests (statistical significance set at 0.05). The accuracy of third molar development based on the London Atlas as age estimation method was measured by means of Receiver Operating Characteristic (ROC) curves and the inherent area under the curve (AUC). The curves considered the legal age threshold of 18 years as cutting point to distinguish individuals (females and males) below or above the age of majority. The tests were performed separately for the teeth #28 and #38. Performance metrics were accuracy, sensitivity, specificity, positive and negative predictive values, and AUC. Statistical analyses were performed with R (R foundation, Vienna, Austria).

Results

The descriptive analysis of our data revealed that the distribution of participants (n = 1.256) per age and sex was nearly optimal since the mean age in each age category and sex was exactly between the minimum and maximum bounds of the age categories (i.e. in the age category of 16-16.99 years, the mean age of both the females and males was 16.5 years) (Table 1).

Table 1: Sample distribution based on sex and age category.

Age	Sex	n	Mean age
16-16.99	F	95	16.5
	M	86	16.5
17-17.99	F	78	17.5
	M	99	17.5
18-18.99	F	89	18.5
	M	80	18.4
19-19.99	F	112	19.5
	M	80	19.5
20-20.99	F	105	20.5
	M	80	20.5
21-21.99	F	110	21.5
	M	72	21.5
22-22.99	F	105	22.5
	M	65	22.5

Age expressed in years; F: females, M: males; n: number of individuals.

Age estimation using tooth #28 showed that most of the individuals were better classified into their age category if their age was in the younger spectrum of the sample's age interval. In this spectrum, age classifications were floating between correct and underestimations. Individuals in the age category of 16-16.99 years, for instance, were correctly estimated in most of the cases (as 16.5 years) or were estimated as 15.5 years. As age increases, wrongful classifications increased as well, revealing a tendency of less accurate performances of the method. A balanced distribution occurred, for example, among the females that were in the age category of 21-21.99 years – they were mostly and similarly classified between 17.5, 18.5, 19.5, 20.5 and 21.5 years (Table 2). The analyses based on the mandibular left third molar (#38) seemed sparser across the age categories, with a mix of underestimations and overestimations. Some of the estimates were distributed with a large number of individuals spread into four age categories. For instance, the females with age 16-16.99 years were estimated as 15.5 (n = 31), 16.5 (n = 19), 17.5 (n = 17) and 18.5 (n = 19) years (Table 2).

Table 2: Age predictions using the maxillary (#28) and mandibular (#38) left third molars for each sex and age category.

#	Age	Sex	Age estimated with the maxillary left third molar									
			15.5	16.5	17.5	18.5	19.5	20.5	21.5	22.5	23.5	n/a
28	16-16.99	F	27	28	16	9	6	1	2	0	0	6
		M	20	16	19	13	7	5	0	0	0	6
	17-17.99	F	12	24	19	9	8	2	0	0	0	4
		M	12	18	24	14	15	4	3	2	0	7
	18-18.99	F	9	16	15	14	14	3	6	3	2	7
		M	9	7	10	19	17	2	4	5	3	4
	19-19.99	F	7	14	23	13	19	10	6	8	1	11
		M	1	11	5	8	15	12	11	8	1	8
	20-20.99	F	3	13	10	16	24	3	11	10	2	13
		M	2	6	5	5	15	11	14	12	0	10
	21-21.99	F	2	7	11	12	19	14	18	15	4	8
		M	0	1	4	1	16	5	12	20	6	7
22-22.99	F	4	4	7	10	17	11	15	13	6	18	
	M	0	2	1	3	7	9	13	11	6	13	
38	16-16.99	F	31	19	17	19	0	4	1	0	0	4
		M	13	8	21	26	8	1	1	0	0	8
	17-17.99	F	14	13	16	22	7	4	0	1	0	1
		M	13	12	23	24	13	4	2	4	1	3
	18-18.99	F	6	8	10	32	12	3	1	8	3	6
		M	7	2	8	20	9	14	5	4	4	7
	19-19.99	F	8	10	12	35	12	8	6	12	4	5
		M	2	5	1	8	21	16	14	7	3	3
	20-20.99	F	2	10	8	23	14	10	13	13	5	7
		M	1	5	1	7	19	17	9	14	5	2
	21-21.99	F	1	1	8	17	22	18	16	12	9	6
		M	2	0	0	3	9	14	11	20	11	2
22-22.99	F	2	4	1	13	19	19	15	14	7	11	
	M	0	1	3	4	6	10	12	20	5	4	

Age expressed in years; F: females, M: males; n: number of individuals; n/a: cases in which age estimation was not applicable.

The mean absolute error (MAE) of the age predictions using tooth #28 was 1.66 years, and the root mean squared error (RMSE) was 2.12 years, for the general sample. In females and males the values of MAE were 1.76 years (RMSE = 2.25 years) and 1.52 years (RMSE = 1.95 years), respectively. The performance of the method was better in the first two age categories (16-16.99 years and 17-17.99 years). In the remaining age categories, the MAE was above 1.5 years and the RMSE was above 2 years (Table 3, Fig. 1). When the tooth #38 was used, the MAE for the general sample was 1.68 years (RMSE = 2.04 years) – 1.68 years (RMSE = 2.12 years) for females and 1.51 years (RMSE = 1.94 years) for males. The same tendency of mean overestimations above

1.5 years in the age groups above 17.99 years was detected, as well the increase of the RMSE above 2 years (Table 3, Fig. 2).

Strong correlations were observed between the development of teeth #28 (0.53, $p < 0.01$) and #38 (0.54, $p < 0.01$) and the chronological age. Among females, the correlation was 0.51 ($p < 0.01$) and 0.52 ($p < 0.01$) for the same teeth, respectively. For males, the correlation was 0.58 ($p < 0.01$) for both teeth.

When the age threshold of legal interest of 18 years was considered, tooth #28 led to 74% of correct classifications of individuals that were younger or older (or equal) than 18 years. Tooth #38 led to 77% of correct classifications. Despite the sensitivity of 0.91 and 0.84 for teeth

Table 3: Distribution of mean absolute errors (MAE) and root mean squared errors (RMSE) per sex and age using the maxillary (#28) and mandibular (#38) left third molars.

#	Age	Sex	Age group-based		Sex-based	
			RMSE	MAE	RMSE	MAE
28	16-16.99	F	1.57	1.17	1.49	1.07
		M			1.65	1.28
	17-17.99	F	1.55	1.23	1.32	1.09
		M			1.71	1.35
	18-18.99	F	2.08	1.64	2.06	1.7
		M			2.11	1.58
	19-19.99	F	2.11	1.76	2.15	1.79
		M			2.04	1.7
	20-20.99	F	2.22	1.84	2.35	2.02
		M			2.03	1.61
	21-21.99	F	2.26	1.78	2.49	1.96
		M			1.85	1.49
	22-22.99	F	2.91	2.27	3.19	2.56
		M			2.35	1.79
38	16-16.99	F	1.61	1.31	1.55	1.21
		M			1.68	1.43
	17-17.99	F	1.74	1.36	1.53	1.26
		M			1.9	1.45
	18-18.99	F	2.09	1.55	2	1.44
		M			2.18	1.68
	19-19.99	F	2.06	1.67	2.17	1.79
		M			1.89	1.51
	20-20.99	F	2.08	1.73	2.22	1.89
		M			1.91	1.52
	21-21.99	F	2	1.6	2.15	1.75
		M			1.76	1.38
	22-22.99	F	2.61	2.02	2.82	2.28
		M			2.25	1.62

F: females; M: males; age expressed in years.

#28 and #38, specificity was below 0.6 for both teeth (Table 4). The area under the curve was 0.75 and 0.73 for teeth #28 and #38, respectively (Fig. 3).

Discussion

In 2022, Goetten *et al.* (19) highlighted the importance of testing dental age estimation method in the Amazon region. The authors focused on three main points: I) the Brazilian border with Venezuela has enabled an increase in the number of clandestine migrations (given the economic challenges in the neighbor country); II) the Northern Brazilian population has scarcer European ancestry compared to other geographic regions of Brazil; and III) the few dental age estimation methods that have

been tested in this population. The London Atlas, for instance, never has been tested in the population of the Amazon Region.

The first set of outcomes obtained with this study confirmed the rationale that better age predictions are obtained in younger ages. When the participants were distributed in age categories, the smallest differences between chronological and estimated ages were observed amongst individuals in age groups 16-16.99 and 17-17.99 years. This finding may be explained by the fact that important variations occur when third molars are closing their apices. Consequently, when age-related biological information becomes restricted to the late stages of root and apex formation, error rates increase (15). This phenome-




Fig. 1: Mean absolute error (MAE) and root mean squared error (RMSE) calculated for age group 1 (16-16.99 years), 2 (17-17.99 years), 3 (18-18.99 years), 4 (19-19.99 years), 5 (20-20.99 years), 6 (21-21.99 years), and 7 (22-22.99 years), using the maxillary left third molar (tooth #28) for females (F) and males (M). The results are presented for the combined (A) sex (F + M), and separately based on sex (B).




Fig. 2: Mean absolute error (MAE) and root mean squared error (RMSE) calculated for age group 1 (16-16.99 years), 2 (17-17.99 years), 3 (18-18.99 years), 4 (19-19.99 years), 5 (20-20.99 years), 6 (21-21.99 years), and 7 (22-22.99 years), using the mandibular left third molar (tooth #38) for females (F) and males (M). The results are presented for the combined (A) sex (F + M), and separately based on sex (B).

non has been observed by Correia *et al.* (15) when they noticed worse specificity values by gradually narrowing their sample (from 16 to 21.99 years) closer to the age of 18 years. This scenario proposes a real challenge to experts because it suggests that some methods might have their performance jeopardized especially when a better performance is needed: around the age of 18 years (for most legal systems).

Contrasting, when The London Atlas was applied based on the developing permanent dentition of Brazilian

individuals aged 6-15.99 years (7), the MAE was only 0.56 and 0.60 years. The current results, based solely on third molar for age estimation in the age group 16-22.99 years, showed MAE of 1.76 and 1.52 years for females and males, respectively. These outcomes seem to point out to a potentially acceptable performance of the method since third molar age estimation often leads to MAE rates that could reach over 2.0 years (20, 21). An example of a study with dental age estimation staging system that have found a higher MAE error rate in Brazilians inclu-

Table 4: Diagnostic accuracy metrics for the maxillary left (#28) and mandibular left (#38) third molars.

Metrics	#28	#38
Accuracy	0.7443	0.7717
Sensitivity	0.9199	0.8473
Specificity	0.3254	0.5848
Positive Predictive Value	0.7648	0.8345
Negative Predictive Value	0.6301	0.6079
Area under the curve	0.75	0.73

^a: 95% CI = 0.7178; 0.7694; ^b: 95% CI = 0.7467; 0.7953; Tooth #28 predicted as true 735 true cases and 226 false cases. Additionally, the same tooth predicted as false 64 true cases and 109 false cases. Tooth #38 predicted as true 716 true cases and 142 false cases. Additionally, the same tooth predicted as false 129 true cases and 200 false cases.




Fig. 3: Receiver operating characteristic (ROC) curves for the predictions of the maxillary (A) and mandibular (B) left third molars leading to areas under the curve (AUC) of 0.75 and 0.73, respectively.

de the one by Sartori *et al.* (20). Comparisons, however, are hampered because the authors used a different age estimation method – namely the method proposed by Gunst *et al.* (22) and sampled a Brazilian population from the opposite geographic region (South).

When accuracy was considered, we noticed a slightly better performance of tooth #38. This tooth is known to be the reference source for dental age estimation analyses when it comes to the metric approach proposed by Cameriere *et al.* 2008 (23), for example. The method based on ratios of third molar length and apical width has led to accuracy rates around 80% in the Northeastern Brazilian population (15). Our outcomes, both for tooth #38 and tooth #28, were similar as the teeth reached accuracy rates of 77% and 74%, respectively. The similarity was even more evident when the comparisons were restricted to the previous applications of The London Atlas in the Brazilian population (15). It must be

noted that the present sample was from the Northern region of Brazil, while the previous application of The London Atlas was in the Northeastern region. These results converge to suggest that third molar development timing might be similarly depicted by The London Atlas between these populations. This finding, however, must not be misinterpreted. Being similar between different populations of the same country does not mean that the method is fully applicable without restrictions. Accuracy rates around 77% mean that almost one in every four classifications of age (below or above the age of majority of 18 years) based on third molar development could be wrong.

The forensic value of The London Atlas as a method should not be diminished based on inherent limitations

for third molar age estimation (common to most third molar age estimation methods). The London Atlas covers a much broader range of applications, including children and young adolescents (24) with evidence proving the method's efficiency (7). Future studies in the field should be planned to test the applicability of the method in other samples, especially when it comes to third molar development. The rationale behind the future studies needed is the scarce literature focusing on the performance of The London Atlas exclusively for third molar age estimation.

Conclusions

The London Atlas showed proper applicability for dental age estimation of young adolescents, especially in the lower age limit of the present sample. The differences between chronological and estimated dental ages increased progressively with time. Slightly better performan-

ces were observed among males compared to females and using the mandibular left third molar compared to the maxillary left third molar. The overall accuracy rates indicated that one in every four classifications of individuals below or above the age of majority (18 years) could be wrong based on The London Atlas – which makes this method a proper tool for age estimation except when used as the sole resource for classifications around the age of 18 years.

References

1. Franco A, Thevissen P, Fieuws S, Souza PH, Willems G. Applicability of Willems model for dental age estimations in Brazilian children. *Forensic Sci Int.* 2013;231:401.
2. Nóbrega JBM, Protasio APL, Ribeiro ILA, Valença AMG, Santiago BM, Cameriere R. Validation of the Third Molar Maturation Index to estimate the age of criminal responsibility in Northeastern Brazil. *Forensic Sci Int.* 2019;304:109917.
3. Deitos AR, Costa C, Michel-Crosato E, Gali I, Cameriere R, Biazevic MG. Age estimation among Brazilians: Younger or older than 18? *J Forensic Leg Med.* 2015;33:111-5.
4. Angelakopoulos N, Franco A, Mula AP, Moukarzel M, Sharma S, Balla SB. Effect of impaction on third molar development and age estimation - A study in a Lebanese population. *Morphologie.* 2023;107:100607.
5. Murray J, Heng D, Lygate A, Porto L, Abade A, Mânica S, et al. Applying artificial intelligence to determination of legal age of majority from radiographic. *Morphologie.* 2023;108:100723.
6. AlQahtani SJ, Hector MP, Liversidge HM. Brief communication: The London atlas of human tooth development and eruption. *Am J Phys Anthropol.* 2010;142:481-90.
7. Dalben E, Bueno J, Fonseca Y, Mânica S, Nascimento M, Franco A. Application of the London Atlas Software App 2nd edition for the radiographic assessment of dental development. *J Clin Exp Dent.* 2022;14:944-52.
8. Zhou J, Qu D, Fan L, Yuan X, Wu Y, Sui M, et al. Applicability of the London Atlas method in the East China population. *Pediatr Radiol.* 2023;53:256-64.
9. Cheong H, Roh BY, Kumagai A, Oh S, Lee SS. Validation of London Atlas for forensic age estimation in Koreans by comparing with Lee's and Willems' methods. *Heliyon.* 2023;9:e19957.
10. Pavlović S, Palmela Pereira C, Santos RFVS. Age estimation in Portuguese population: The application of the London atlas of tooth development and eruption. *Forensic Sci Int.* 2017;272:97-103.
11. Ishwarkumar S, Pillay P, Chetty M, Satyapal KS. Employing the London Atlas in the age estimation of a select South African population. *Dent J (Basel).* 2022;10:171.
12. Namwong W, Mânica S. Testing the London atlas for age estimation in Thai population. *Acta Odont Scand.* 2020;78:161-4.
13. Jacometti V, Sato CM, Meireles DA, Silva RHAD. Age estimation using London Atlas methodology: A systematic review and meta-analysis. *Forensic Sci Int.* 2023;342:111532.
14. Franco A, de Oliveira MN, Campos Vidigal MT, Blumenberg C, Pinheiro AA, Paranhos LR. Assessment of dental age estimation methods applied to Brazilian children: a systematic review and meta-analysis. *Dentomaxillofac Radiol.* 2021;50:20200128.
15. Correia AM, Barbosa DDS, Alcantara JADS, Oliveira PMDC, Silva PGB, Franco A, et al. Performance and comparison of the London Atlas technique and Cameriere's third molar maturity index (I3M) for allocating individuals below or above the threshold of 18 years. *Forensic Sci Int.* 2020;317:110512.
16. Sousa AMDS, Jacometti V, AlQahtani S, Silva RHAD. Age estimation of Brazilian individuals using the London Atlas. *Arch Oral Biol.* 2020;113:104705.
17. Vandembroucke JP, von Elm E, Altman DG, Gøtzsche PC, Mulrow CD, Pocock SJ, et al; STROBE Initiative. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *PLoS Med.* 2007;4:e297.
18. AlQahtani SJ, Hector MP, Liversidge HM. Accuracy of dental age estimation charts: Schour and Massler, Ubelaker and the London Atlas. *Am J Phys Anthropol.* 2014;154:70-8.
19. Goetten IFS, Oenning ACC, Silva RF, Nuzzolese E, Lourenço-Junior E, Franco A. Diagnostic accuracy of the third molar maturity index (I3M) to assess the age of legal majority in Northern Brazil-population-specific cut-off values. *Int J Legal Med.* 2022;136:1507-14.
20. Sartori VK, Franco A, Linden MS, Cardoso M, Castro D, Sartori A, et al. Testing international techniques for the radiographic assessment of third molar maturation. *J Clin Exp Dent.* 2021;13:1182-8.
21. Kanchan T, Chugh V, Chugh A, Meshram V, Shedge R, Patnana AK. Age estimation using third molar maturation based on Demirjian's criteria. *Legal Med.* 2021;53:101959.
22. Gunst K, Mesotten K, Carbonez A, Willems G. Third molar root development in relation to chronological age: a large sample sized retrospective study. *Forensic Sci Int.* 2003;136:52-7.
23. Cameriere R, Ferrante L, De Angelis D, Scarpino F, Galli F. The comparison between measurement of open apices of third molars and Demirjian stages to test chronological age of over 18 year olds in living subjects. *Int J Legal Med.* 2008;122:493-7.
24. McCloe D, Marion I, da Fonseca MA, Colvard M, AlQahtani S. Age estimation of Hispanic children using the London Atlas. *Forensic Sci Int.* 2018;288:332.

Conflict of interest

None declared.

ORIGINALITY REPORT

22%

SIMILARITY INDEX

14%

INTERNET SOURCES

19%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Marcos Vinicius Fernandes Machado, Mariana Quirino Silveira Soares, Andrea Maia Sampaio Alonso Baz, Jose Luiz Cintra Junqueira et al. "A large sample-sized study on dental development of children treated at the Central Dental Clinic (OCEx) of the Brazilian Army", *Clinical Oral Investigations*, 2022
Publication 2%
- 2** www.researchsquare.com
Internet Source 1%
- 3** Sukran Senyurek, Efsun Somay, Nilufer Kilic Durankus, Sibel Bascil, Duriye Ozturk, Ugur Selek, Erkan Topkan. "The unique CARWL score stratifies locally advanced nasopharyngeal cancer patients receiving concurrent chemoradiotherapy into risk groups for radiation-induced trismus", *Discover Oncology*, 2024
Publication 1%
- 4** Izabella Ferreira dos Santos Goetten, Anne Caroline Costa Oenning, Rhonan Ferreira 1%

Silva, Emilio Nuzzolese et al. "Diagnostic accuracy of the third molar maturity index (I3M) to assess the age of legal majority in Northern Brazil—population-specific cut-off values", *International Journal of Legal Medicine*, 2022

Publication

5

A. Franco, A.P. Cornacchia, D. Moreira, P. Miamoto, J. Bueno, J. Murray, D. Heng, S. Mânica, L. Porto, A. Abade. "Radiographic morphology of canines tested for sexual dimorphism via convolutional-neural-network-based artificial intelligence", *Morphologie*, 2024

Publication

1 %

6

ejfs.springeropen.com

Internet Source

1 %

7

roderic.uv.es

Internet Source

1 %

8

Rafael Cidade, Marcelo dos Santos, Thaís Cássia Alves, Juliano Martins Bueno et al. "Radiographic dental age estimation applying and comparing Demirjian's seven (1973) and four (1976) teeth methods", *Forensic Science, Medicine and Pathology*, 2023

Publication

1 %

9

Nikolaos Angelakopoulos, Nallan CSK Chaitanya, Sudheer B. Balla. "Effect of

1 %

impaction on third molar development and age estimation–A study in a female black South African population", Archives of Legal Medicine, 2024

Publication

10

pubmed.ncbi.nlm.nih.gov

Internet Source

1 %

11

Gretel González-Colmenares, María del Socorro Barraza Salcedo, Natalia Bernal-Ortiz, Jeison Nicolás Cepeda-Rojas et al. "Estimation of Dental Age in a Sample of Colombian Population Using the London Atlas", Forensic Science International: Reports, 2022

Publication

1 %

12

www.ricardohenrique.com.br

Internet Source

1 %

13

www.medoraljced.com

Internet Source

1 %

14

Adriana de Moraes Correia, Debora da Silva Barbosa, Jeisiana Alves da Silva Alcantara, Patrícia Maria da Costa Oliveira et al. "Performance and comparison of the London Atlas technique and Cameriere's third molar maturity index (I3M) for allocating individuals below or above the threshold of 18 years", Forensic Science International, 2020

Publication

1 %

15

Carolina Malschitzky, Maria Tereza Campos Vidigal, Debora Duarte Moreira, Rhonan Ferreira Silva et al. "How reliable is stature estimation by dental means? Systematic review and meta-analysis", *Forensic Science International*, 2024

Publication

1 %

16

Raquel Porto Alegre Valente Franco, Ademir Franco, Anna Turkina, Marianna Arakelyan et al. "Radiographic assessment of third molar development in a Russian population to determine the age of majority", *Archives of Oral Biology*, 2021

Publication

1 %

17

A. Pinheiro, R. Franco, I. Makeeva, J. Bueno, P. Miamoto, A. Franco. "30years of the ABFO study: Reproduction in a Brazilian sample", *Morphologie*, 2023

Publication

<1 %

18

link.springer.com

Internet Source

<1 %

19

N. Angelakopoulos, A. Franco, A.P. Mula, M. Moukarzel, S. Sharma, S.B. Balla. "Effect of impaction on third molar development and age estimation—A study in a Lebanese population", *Morphologie*, 2023

Publication

<1 %

20	Submitted to Temple University Student Paper	<1 %
21	Heba H. Bakhsh, Nada A. Al-shehri, Alanoud Shahwan, Rabab Altuwairqi, Faten J. Mojaleed, Ghaida Alwaalan, Shahad Asaad. "A Comparison of Two Methods of Dental Age Estimation in a Population of Saudi Children and Adolescents", Diagnostics, 2024 Publication	<1 %
22	acikerisim.hacettepe.edu.tr Internet Source	<1 %
23	boris.unibe.ch Internet Source	<1 %
24	ejmcm.com Internet Source	<1 %
25	Jun Wang, Andrew Kusiak. "Computational Intelligence In Manufacturing Handbook", CRC Press, 2019 Publication	<1 %
26	Submitted to University of Dundee Student Paper	<1 %
27	www.journals.usp.br Internet Source	<1 %
28	www.mdpi.com Internet Source	<1 %

29 www.ncbi.nlm.nih.gov <1 %
Internet Source

30 www.tandfonline.com <1 %
Internet Source

31 www2.mdpi.com <1 %
Internet Source

32 Dara M. Alsudairi, Sakher J. AlQahtani.
"Testing and Comparing the Accuracy of Two
Dental Age Estimation Methods on Saudi
Children: Measurements of Open Apices in
Teeth and The London Atlas of Tooth
Development", Forensic Science International,
2018
Publication

33 Dimitrios A. Karras, Srinesh Thakur, Sai Kiran
Oruganti. "Advancements in Science and
Technology for Healthcare, Agriculture, and
Environmental Sustainability: A Review of the
Latest Research and Innovations", CRC Press,
2024
Publication

34 academic.oup.com <1 %
Internet Source

35 www.science.gov <1 %
Internet Source

36

"Fundamentals of Forensic Biology", Springer
Science and Business Media LLC, 2024

Publication

<1 %

37

Marco Cummaudo, Danilo De Angelis,
Francesca Magli, Giulia Minà, Vera Merelli,
Cristina Cattaneo. "Age estimation in the
living: A scoping review of population data for
skeletal and dental methods", Forensic
Science International, 2021

Publication

<1 %

38

Ron D. Hays, Phyllis L. Ellickson.
"COMPARISON OF THE ROST AND THE CAGE
ALCOHOL SCREENING INSTRUMENTS IN
YOUNG ADULTS", Substance Use & Misuse,
2001

Publication

<1 %

39

acikerisimarsiv.selcuk.edu.tr:8080

Internet Source

<1 %

40

biblio.ugent.be

Internet Source

<1 %

41

doi.org

Internet Source

<1 %

42

www.iofos.eu

Internet Source

<1 %

43

www.researchgate.net

Internet Source

<1 %

44

Ademir Franco, Débora Duarte Moreira, Rafael Cidade, Marcos Machado et al. "The Brazilian (FRANCO) method for dental age estimation: Willems' model revisited", *Clinical Oral Investigations*, 2024

Publication

<1 %

45

Akiko Kumagai, Guy Willems, Ademir Franco, Patrick Thevissen. "Age estimation combining radiographic information of two dental and four skeletal predictors in children and subadults", *International Journal of Legal Medicine*, 2018

Publication

<1 %

46

Angelines Cruz-Landeira. "Dental age estimation in Spanish and Venezuelan children. Comparison of Demirjian and Chaillet's scores", *International Journal of Legal Medicine*, 10/28/2009

Publication

<1 %

47

Nicolás Vila-Blanco, Paulina Varas-Quintana, Inmaculada Tomás, María J. Carreira. "A systematic overview of dental methods for age assessment in living individuals: from traditional to artificial intelligence-based approaches", *International Journal of Legal Medicine*, 2023

Publication

<1 %

48

Sehyun Oh, Akiko Kumagai, Sin-Young Kim, Sang-Seob Lee. "Accuracy of age estimation and assessment of the 18-year threshold based on second and third molar maturity in Koreans and Japanese", PLOS ONE, 2022

Publication

<1 %

49

[sites.google.com](https://www.google.com)

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

2024 Brazil London Atlas M3.pdf

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8